



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12Q 1/68, C12N 15/00, 15/09, 15/63, 15/86</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/55375</b> <b>(43) International Publication Date:</b> 21 September 2000 (21.09.00)																		
<b>(21) International Application Number:</b> PCT/US00/07285 <b>(22) International Filing Date:</b> 17 March 2000 (17.03.00)  <b>(30) Priority Data:</b> <table border="0"><tr><td>60/124,916</td><td>17 March 1999 (17.03.99)</td><td>US</td></tr><tr><td>60/124,808</td><td>17 March 1999 (17.03.99)</td><td>US</td></tr><tr><td>60/149,639</td><td>17 August 1999 (17.08.99)</td><td>US</td></tr><tr><td>60/157,247</td><td>1 October 1999 (01.10.99)</td><td>US</td></tr><tr><td>60/167,824</td><td>29 November 1999 (29.11.99)</td><td>US</td></tr><tr><td>60/182,711</td><td>15 February 2000 (15.02.00)</td><td>US</td></tr></table> <b>(71) Applicant:</b> ALPHAGENE, INC. [US/US]; 260 West Cummings Park, Woburn, MA 01801 (US).  <b>(72) Inventors:</b> VALENZUELA, Dario; 1081 Hill Road, Boxborough, MA 01719-1010 (US). YUAN, Olive; 292 Mystic Street, Arlington, MA 02174 (US). HOFFMAN, Heidi; 90 Houghton Mill Road, Lunenburg, MA 01462 (US). HALL, Jeff; 4 Alderwood Drive, Stratham, NH 03885 (US). RAPIEJKO, Peter; 63 Old Grafton Road, Upton, MA 01568 (US).  <b>(74) Agent:</b> SPRUNGER, Suzanne, A.; American Home Products Corporation, Patent & Trademark Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US).		60/124,916	17 March 1999 (17.03.99)	US	60/124,808	17 March 1999 (17.03.99)	US	60/149,639	17 August 1999 (17.08.99)	US	60/157,247	1 October 1999 (01.10.99)	US	60/167,824	29 November 1999 (29.11.99)	US	60/182,711	15 February 2000 (15.02.00)	US	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
60/124,916	17 March 1999 (17.03.99)	US																		
60/124,808	17 March 1999 (17.03.99)	US																		
60/149,639	17 August 1999 (17.08.99)	US																		
60/157,247	1 October 1999 (01.10.99)	US																		
60/167,824	29 November 1999 (29.11.99)	US																		
60/182,711	15 February 2000 (15.02.00)	US																		
<b>(54) Title:</b> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM  <b>(57) Abstract</b>  Novel polynucleotides and the proteins encoded thereby are disclosed.																				

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



## SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

5                   This application is a continuation-in-part of the following applications:

- (1)   provisional application Ser. No. 60/124,916, filed March 17, 1999;
  - (2)   provisional application Ser. No. 60/124,808, filed March 17, 1999;
  - (3)   provisional application Ser. No. 60/149,639, filed August 17, 1999;
  - (4)   provisional application Ser. No. 60/157,247, filed October 1, 1999;
  - 10 (5)   provisional application Ser. No. 60/167,824, filed November 29, 1999;
  - (6)   provisional application Ser. No. 60/182,711, filed February 15, 2000;
- all of which are incorporated by reference herein.

FIELD OF THE INVENTION

15           The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these polynucleotides and proteins.

BACKGROUND OF THE INVENTION

20           Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past decade. The now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of hybridization cloning; activity of the protein in the case of expression cloning). More recent "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, have advanced the state of the art by making available large numbers of DNA/amino acid sequences for proteins that are known to have biological activity by virtue of their secreted nature in the case of leader sequence cloning, or by virtue of the cell or tissue source in the case of PCR-based techniques. It is to these proteins and the polynucleotides encoding them that the present invention is directed.

25

30

SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 27 to nucleotide 260;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 72 to nucleotide 260;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc62\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number  
15 207114;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc62\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA  
20 insert of clone vc62\_1 deposited with the ATCC under accession number 207114;
- (h) ----- a polynucleotide encoding a protein comprising the amino acid  
sequence of SEQ ID NO:2;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment  
25 comprising eight contiguous amino acids of SEQ ID NO:2;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:1.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 27 to nucleotide 260; the nucleotide sequence of SEQ ID NO:1 from nucleotide 72 to nucleotide 260; the nucleotide sequence of the full-length protein coding sequence of clone vc62\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vc62\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:2.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:1.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:1 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1 from nucleotide 27 to nucleotide 260, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:1 from nucleotide 27 to nucleotide 260, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 27 to nucleotide 260. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1 from nucleotide 72 to nucleotide 260, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:1 from nucleotide 72 to nucleotide 260, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 72 to nucleotide 260.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins. Preferably such
- 10 protein comprises the amino acid sequence of SEQ ID NO:2. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a protein comprising a fragment of the amino acid sequence of SEQ
- 15 ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:2.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:3;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 6 to nucleotide 1325;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 99 to nucleotide 1325;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp10\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number
- 30 207114;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp10\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA  
5 insert of clone vp10\_1 deposited with the ATCC under accession number 207114;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment  
10 comprising eight contiguous amino acids of SEQ ID NO:4;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any  
15 one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:3.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:3 from nucleotide 6 to nucleotide 1325; the nucleotide sequence of SEQ ID NO:3 from nucleotide 99 to nucleotide 1325; the nucleotide sequence of the full-length protein coding sequence of clone vp10\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vp10\_1 deposited  
25 with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological  
30 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having

biological activity, the fragment comprising the amino acid sequence from amino acid 215 to amino acid 224 of SEQ ID NO:4.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:3.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(ab) the nucleotide sequence of the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number 207114;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(bb) the nucleotide sequence of the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:3 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 6 to nucleotide 1325, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 6 to nucleotide 1325, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 6 to nucleotide 1325. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 99 to nucleotide 1325, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 99 to nucleotide 1325, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 99 to nucleotide 1325.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:4. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising the amino acid sequence from amino acid 215 to amino acid 224 of SEQ ID NO:4.



In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 149 to nucleotide 322;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 200 to nucleotide 322;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp11\_1 deposited with the ATCC under  
10 accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp11\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any  
30 one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:5.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:5 from nucleotide 149 to nucleotide 322; the nucleotide sequence of SEQ ID NO:5 from nucleotide 200 to nucleotide 322; the nucleotide sequence of the full-length protein coding sequence of clone vp11\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vp11\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising the amino acid sequence from amino acid 24 to amino acid 33 of SEQ ID NO:6.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:5.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:5 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5 from nucleotide 149 to nucleotide 322, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:5 from nucleotide 149 to nucleotide 322, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:5 from nucleotide 149 to nucleotide 322. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5 from nucleotide 200 to nucleotide 322, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:5 from nucleotide 200 to nucleotide 322, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:5 from nucleotide 200 to nucleotide 322.

20

25

30

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins. Preferably such
- 10 protein comprises the amino acid sequence of SEQ ID NO:6. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a protein comprising a fragment of the amino acid sequence of SEQ
- 15 ID NO:6 having biological activity, the fragment comprising the amino acid sequence from amino acid 24 to amino acid 33 of SEQ ID NO:6.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:7;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 288 to nucleotide 629;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 363 to nucleotide 629;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp13\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number
- 30 207114;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp13\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:8;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:7.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:7 from nucleotide 288 to nucleotide 629; the nucleotide sequence of SEQ ID NO:7 from nucleotide 363 to nucleotide 629; the nucleotide sequence of the full-length protein coding sequence of clone vp13\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vp13\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of

SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 52 to amino acid 61 of SEQ ID NO:8.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:7.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and

(ab) the nucleotide sequence of the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and

(bb) the nucleotide sequence of the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:7 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 288 to nucleotide 629, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 288 to nucleotide 629, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 288 to nucleotide 629. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 363 to nucleotide 629, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 363 to nucleotide 629, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 363 to nucleotide 629.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:8. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 52 to amino acid 61 of SEQ ID NO:8.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 11 to nucleotide 298;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 149 to nucleotide 298;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp16\_1 deposited with the ATCC under  
10 accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp16\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any  
30 one of the polynucleotides specified in (a)-(i); and



(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:9.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:9 from nucleotide 11 to nucleotide 298; the nucleotide sequence of SEQ ID NO:9 from nucleotide 149 to nucleotide 298; the nucleotide sequence of the full-length protein coding sequence of clone vp16\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vp16\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:10.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:9.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:9 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 11 to nucleotide 298, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:9 from nucleotide 11 to nucleotide 298, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 11 to nucleotide 298. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 149 to nucleotide 298, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:9 from nucleotide 149 to nucleotide 298, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 149 to nucleotide 298.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
- 5 (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins. Preferably such  
10 protein comprises the amino acid sequence of SEQ ID NO:10. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a protein comprising a fragment of the amino acid sequence of SEQ  
15 ID NO:10 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:10.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 20 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 257 to nucleotide 607;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 479 to nucleotide 607;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp21\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number  
30 207114;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp21\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:11.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:11 from nucleotide 257 to nucleotide 607; the nucleotide sequence of SEQ ID NO:11 from nucleotide 479 to nucleotide 607; the nucleotide sequence of the full-length protein coding sequence of clone vp21\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vp21\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of

SEQ ID NO:12 having biological activity, the fragment comprising the amino acid sequence from amino acid 53 to amino acid 62 of SEQ ID NO:12.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:11.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(ab) the nucleotide sequence of the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(bb) the nucleotide sequence of the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:11 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:11, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:11. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 257 to nucleotide 607, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 257 to nucleotide 607, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 257 to nucleotide 607. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 479 to nucleotide 607, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from  
15 nucleotide 479 to nucleotide 607, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 479 to nucleotide 607.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:12;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:12. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:12, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising the amino acid sequence from amino acid 53 to amino acid 62 of SEQ ID NO:12.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13 from nucleotide 163 to nucleotide 477;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13 from nucleotide 238 to nucleotide 477;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp22\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp22\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:14;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:13.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:13 from nucleotide 163 to nucleotide 477; the nucleotide sequence of SEQ ID NO:13 from nucleotide 238 to nucleotide 477; the nucleotide sequence of the full-length protein coding sequence of clone vp22\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vp22\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:14.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:13.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and



(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:13 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:13, but  
20 excluding the poly(A) tail at the 3' end of SEQ ID NO:13. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 163 to nucleotide 477, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:13 from nucleotide 163 to nucleotide 477, to a nucleotide  
25 sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 163 to nucleotide 477. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 238 to nucleotide 477, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:13 from  
30 nucleotide 238 to nucleotide 477, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 238 to nucleotide 477.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:14;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins. Preferably such
- 10 protein comprises the amino acid sequence of SEQ ID NO:14. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a protein comprising a fragment of the amino acid sequence of SEQ
- 15 ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:14.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:15;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15 from nucleotide 58 to nucleotide 624;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15 from nucleotide 106 to nucleotide 624;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq2\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number
- 30 207114;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq2\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:15.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:15 from nucleotide 58 to nucleotide 624; the nucleotide sequence of SEQ ID NO:15 from nucleotide 106 to nucleotide 624; the nucleotide sequence of the full-length protein coding sequence of clone vq2\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vq2\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of

SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 89 to amino acid 98 of SEQ ID NO:16.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:15.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(ab) the nucleotide sequence of the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(bb) the nucleotide sequence of the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:15 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:15, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:15. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 58 to nucleotide 624, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from nucleotide 58 to nucleotide 624, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide 58 to nucleotide 624. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 106 to nucleotide 624, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from  
15 nucleotide 106 to nucleotide 624, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide 106 to nucleotide 624.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:16;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:16. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:16, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 89 to amino acid 98 of SEQ ID NO:16.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 773 to nucleotide 1090;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 842 to nucleotide 1090;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq3\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq3\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:18;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:17.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:17 from nucleotide 773 to nucleotide 1090; the nucleotide sequence of SEQ ID NO:17 from nucleotide 842 to nucleotide 1090; the nucleotide sequence of the full-length protein coding sequence of clone vq3\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vq3\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:18.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:17.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:17 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:17, but  
20 excluding the poly(A) tail at the 3' end of SEQ ID NO:17. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 773 to nucleotide 1090, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from nucleotide 773 to nucleotide 1090, to a nucleotide  
25 sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 773 to nucleotide 1090. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 842 to nucleotide 1090, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from  
30 nucleotide 842 to nucleotide 1090, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 842 to nucleotide 1090.



In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:18;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins. Preferably such
- 10 protein comprises the amino acid sequence of SEQ ID NO:18. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a protein comprising a fragment of the amino acid sequence of SEQ
- 15 ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:18.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:19;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19 from nucleotide 96 to nucleotide 275;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19 from nucleotide 159 to nucleotide 275;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq5\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number
- 30 207114;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq5\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:19.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:19 from nucleotide 96 to nucleotide 275; the nucleotide sequence of SEQ ID NO:19 from nucleotide 159 to nucleotide 275; the nucleotide sequence of the full-length protein coding sequence of clone vq5\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vq5\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:20, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of

SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:20.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:19.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (aa) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and

(ab) the nucleotide sequence of the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and

(bb) the nucleotide sequence of the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:19 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:19, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:19. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19 from nucleotide 96 to nucleotide 275, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:19 from nucleotide 96 to nucleotide 275, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:19 from nucleotide 96 to nucleotide 275. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19 from nucleotide 159 to nucleotide 275, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:19 from  
15 nucleotide 159 to nucleotide 275, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:19 from nucleotide 159 to nucleotide 275.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:20;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:20. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:20, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:20.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 176 to nucleotide 340;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 230 to nucleotide 340;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq6\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq6\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:21.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:21 from nucleotide 176 to nucleotide 340; the nucleotide sequence of SEQ ID NO:21 from nucleotide 230 to nucleotide 340; the nucleotide sequence of the full-length protein coding sequence of clone vq6\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vq6\_1 deposited with the ATCC under accession number 207114. In other preferred  
5  
10  
15  
embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 22 to amino acid 31 of SEQ ID NO:22.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:21.  
20

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
30 vq6\_1 deposited with the ATCC under accession number 207114;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:21 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:21, but  
20 excluding the poly(A) tail at the 3' end of SEQ ID NO:21. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 176 to nucleotide 340, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from nucleotide 176 to nucleotide 340, to a nucleotide  
25 sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 176 to nucleotide 340. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 230 to nucleotide 340, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from  
30 nucleotide 230 to nucleotide 340, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 230 to nucleotide 340.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins. Preferably such
- 10 protein comprises the amino acid sequence of SEQ ID NO:22. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a protein comprising a fragment of the amino acid sequence of SEQ
- 15 ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 22 to amino acid 31 of SEQ ID NO:22.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:23;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 29 to nucleotide 1111;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 167 to nucleotide 1111;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vr1\_1 deposited with the ATCC under accession number 207114;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number
- 30 207114;



- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vr1\_1 deposited with the ATCC under accession number 207114;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number 207114;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:23.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:23 from nucleotide 29 to nucleotide 1111; the nucleotide sequence of SEQ ID NO:23 from nucleotide 167 to nucleotide 1111; the nucleotide sequence of the full-length protein coding sequence of clone vr1\_1 deposited with the ATCC under accession number 207114; or the nucleotide sequence of a mature protein coding sequence of clone vr1\_1 deposited with the ATCC under accession number 207114. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number 207114.
- In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:24, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of

SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 175 to amino acid 184 of SEQ ID NO:24.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:23.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and

(ab) the nucleotide sequence of the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number 207114;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and

(bb) the nucleotide sequence of the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number 207114;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:23 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:23, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:23. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 29 to nucleotide 1111, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from nucleotide 29 to nucleotide 1111, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 29 to nucleotide 1111. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 167 to nucleotide 1111, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from  
15 nucleotide 167 to nucleotide 1111, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 167 to nucleotide 1111.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:24;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
vr1\_1 deposited with the ATCC under accession number 207114;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:24. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:24, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 175 to amino acid 184 of SEQ ID NO:24.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25 from nucleotide 13 to nucleotide 513;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc63\_1 deposited with the ATCC under accession number 207115;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc63\_1 deposited with the ATCC under  
15 accession number 207115;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:26;
- 20 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 25 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:25.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:25 from nucleotide 13 to nucleotide 513; the nucleotide sequence of the full-length protein coding sequence of clone vc63\_1 deposited with the ATCC under accession number 207115; or the nucleotide sequence of a mature protein coding sequence of clone  
5 vc63\_1 deposited with the ATCC under accession number 207115. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of  
10 SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:26, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:26.

15 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:25.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 20 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C. to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and
    - 25 (ab) the nucleotide sequence of the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115;
    - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
    - (iii) isolating the DNA polynucleotides detected with the  
30 probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and
- (bb) the nucleotide sequence of the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 10 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:25 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25 from nucleotide 13 to nucleotide 513, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:25 from nucleotide 13 to nucleotide 513, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:25 from nucleotide 13 to nucleotide 513.

15

20

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

25

- (a) the amino acid sequence of SEQ ID NO:26;
- (b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115;
- 30
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:26. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:26, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:26.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 79 to nucleotide 345;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 130 to nucleotide 345;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb25\_1 deposited with the ATCC under accession number PTA-362;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number  
20 PTA-362;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb25\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA  
25 insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:28;
- (i) a polynucleotide encoding a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:27.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:27 from nucleotide 79 to nucleotide 345; the nucleotide sequence of SEQ ID NO:27 from nucleotide 130 to nucleotide 345; the nucleotide sequence of the full-length protein coding sequence of clone vb25\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vb25\_1  
15 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28  
20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most-preferably thirty) contiguous amino acids of SEQ ID NO:28, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 39 to amino acid 48 of SEQ ID NO:28.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:27.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:



- (aa) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and
- (ab) the nucleotide sequence of the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and
- (bb) the nucleotide sequence of the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:27 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 79 to nucleotide 345, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 79 to nucleotide 345, to a nucleotide

sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 79 to nucleotide 345. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 130 to nucleotide 345, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 130 to nucleotide 345, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 130 to nucleotide 345.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:28;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:28. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 39 to amino acid 48 of SEQ ID NO:28.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 72 to nucleotide 236;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 150 to nucleotide 236;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:30;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:29.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:29 from nucleotide 72 to nucleotide 236; the nucleotide sequence of SEQ ID NO:29 from nucleotide 150 to nucleotide 236; the nucleotide sequence of the full-length protein coding sequence of clone vb27\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vb27\_1 deposited with the ATCC under accession number PTA-362. In other preferred

30

embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30  
5 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 22 to amino acid 31 of SEQ ID NO:30.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:29.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 15 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and
    - 20 (ab) the nucleotide sequence of the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize  
30 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and
- (bb) the nucleotide sequence of the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:29 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29. Also preferably the
- 15 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 72 to nucleotide 236, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 72 to nucleotide 236, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:29 from nucleotide
- 20 72 to nucleotide 236. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 150 to nucleotide 236, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 150 to nucleotide 236, to a nucleotide sequence corresponding to the 3' end of
- 25 said sequence of SEQ ID NO:29 from nucleotide 150 to nucleotide 236.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;
- 30 (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:30. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence  
10 from amino acid 22 to amino acid 31 of SEQ ID NO:30.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 884;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31 from nucleotide 183 to nucleotide 884;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb28\_1 deposited with the ATCC under  
20 accession number PTA-362;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362;
- 25 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb28\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-  
30 362;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:32;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:31.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 884; the nucleotide sequence of SEQ ID NO:31 from nucleotide 183 to nucleotide 884; the nucleotide sequence of the full-length protein coding sequence of clone vb28\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vb28\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:32, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising the amino acid sequence from amino acid 120 to amino acid 129 of SEQ ID NO:32.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:31.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (aa) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and

(ab) the nucleotide sequence of the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362;

10 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

15 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and

20 (bb) the nucleotide sequence of the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

25 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID  
30 NO:31 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence



corresponding to the cDNA sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 884, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 884, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 884. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31 from nucleotide 183 to nucleotide 884, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:31 from nucleotide 183 to nucleotide 884, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:31 from nucleotide 183 to nucleotide 884.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:32;
- (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:32. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:32, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising the amino acid sequence from amino acid 120 to amino acid 129 of SEQ ID NO:32.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33 from nucleotide 42 to nucleotide 206;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33 from nucleotide 111 to nucleotide 206;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:34;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:33.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:33 from nucleotide 42 to nucleotide 206; the nucleotide sequence of SEQ ID NO:33 from nucleotide 111 to nucleotide 206; the nucleotide sequence of the full-length protein coding sequence of clone vb29\_1 deposited with the ATCC under accession number PTA-

362; or the nucleotide sequence of a mature protein coding sequence of clone vb29\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-  
5 362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:34, or a  
10 polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 22 to amino acid 31 of SEQ ID NO:34.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:33.

Further embodiments of the invention provide isolated polynucleotides produced  
15 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (aa) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and

(ab) the nucleotide sequence of the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;

25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

30 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and
- (bb) the nucleotide sequence of the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- 10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33, and

15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:33 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33 from nucleotide

20 206, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:33 from nucleotide 42 to nucleotide 206, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:33 from nucleotide 42 to nucleotide 206. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

25 NO:33 from nucleotide 111 to nucleotide 206, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:33 from nucleotide 111 to nucleotide 206, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:33 from nucleotide 111 to nucleotide 206.

In other embodiments, the present invention provides a composition comprising

30 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;

- (b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:34. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:34, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 22 to amino acid 31 of SEQ ID NO:34.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 17 to nucleotide 253;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:35 from nucleotide 98 to nucleotide 253;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb30\_1 deposited with the ATCC under accession number PTA-362;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb30\_1 deposited with the ATCC under accession number PTA-362;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:35.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:35 from nucleotide 17 to nucleotide 253; the nucleotide sequence of SEQ ID NO:36 from nucleotide 98 to nucleotide 253; the nucleotide sequence of the full-length protein coding sequence of clone vb30\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vb30\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:36, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:36.
- Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:35.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and

(ab) the nucleotide sequence of the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and

(bb) the nucleotide sequence of the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID

NO:35 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35 from nucleotide 17 to nucleotide 5 253, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:35 from nucleotide 17 to nucleotide 253, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:35 from nucleotide 17 to nucleotide 253. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID 10 NO:35 from nucleotide 98 to nucleotide 253, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:35 from nucleotide 98 to nucleotide 253, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:35 from nucleotide 98 to nucleotide 253.

In other embodiments, the present invention provides a composition comprising 15 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:36;
- (b) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and
- 20 (c) the amino acid sequence encoded by the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:36. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino 25 acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:36, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:36.

30 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:



- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37 from nucleotide 68 to nucleotide 424;
- 5 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc67\_1 deposited with the ATCC under accession number PTA-362;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc67\_1 deposited with the ATCC under accession number
- 10 PTA-362;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc67\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA
- 15 insert of clone vc67\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:38;
- (h) a polynucleotide encoding a protein comprising a fragment of the
- 20 amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein
- 25 of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any
- 30 one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:37.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:37 from nucleotide 68 to nucleotide 424; the nucleotide sequence of the full-length

protein coding sequence of clone vc67\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vc67\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc67\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:38, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:38.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:37.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37; and

(ab) the nucleotide sequence of the cDNA insert of clone vc67\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37; and
- (bb) the nucleotide sequence of the cDNA insert of clone vc67\_1 deposited with the ATCC under accession number PTA-362;
- 10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37, and  
15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:37 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37 from nucleotide  
20 424, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:37 from nucleotide 68 to nucleotide 424, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:37 from nucleotide 68 to nucleotide 424.

In other embodiments, the present invention provides a composition comprising  
25 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:38;
- (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and
- 30 (c) the amino acid sequence encoded by the cDNA insert of clone vc67\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:38. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably  
5 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:38, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:38.

In one embodiment, the present invention provides a composition comprising an  
10 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39 from nucleotide 103 to nucleotide 261;
- 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39 from nucleotide 154 to nucleotide 261;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- 20 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vf4\_1 deposited with the ATCC under accession  
25 number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:40;
- 30 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:39.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:39 from nucleotide 103 to nucleotide 261; the nucleotide sequence of SEQ ID NO:39 from nucleotide 154 to nucleotide 261; the nucleotide sequence of the full-length protein coding sequence of clone vf4\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vf4\_1  
15 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40  
20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:40, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:40.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:39.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and
- (ab) the nucleotide sequence of the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- 5 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- 10 (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:39, but excluding the poly(A) tail at the 15 3' end of SEQ ID NO:39; and
- (bb) the nucleotide sequence of the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 20 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID

25 NO:39 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 103 to nucleotide 261, and extending contiguously from a nucleotide sequence corresponding to the 5' end

30 of said sequence of SEQ ID NO:39 from nucleotide 103 to nucleotide 261, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from nucleotide 103 to nucleotide 261. Also preferably the polynucleotide isolated according to the above

process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 154 to nucleotide 261, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:39 from nucleotide 154 to nucleotide 261, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from nucleotide 154 to nucleotide 261.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
  - 10 (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:40. In further preferred
- 15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:40, or a protein comprising a fragment of the amino acid sequence of SEQ
- 20 ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:40.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 25 NO:41;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 1575 to nucleotide 3038;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 1650 to nucleotide 3038;
- 30 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vg3\_1 deposited with the ATCC under accession number PTA-362;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:42;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:41.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:41 from nucleotide 1575 to nucleotide 3038; the nucleotide sequence of SEQ ID NO:41 from nucleotide 1650 to nucleotide 3038; the nucleotide sequence of the full-length protein coding sequence of clone vg3\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vg3\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by

30 the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42



having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid  
5 sequence from amino acid 239 to amino acid 248 of SEQ ID NO:42.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:41.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:41, but excluding the poly(A) tail at the  
15 3' end of SEQ ID NO:41; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:41, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:41; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vg3\_1 deposited with the ATCC under accession number PTA-362;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:41 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 1575 to  
10 nucleotide 3038, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 1575 to nucleotide 3038, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 1575 to nucleotide 3038. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the  
15 cDNA sequence of SEQ ID NO:41 from nucleotide 1650 to nucleotide 3038, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 1650 to nucleotide 3038, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 1650 to nucleotide 3038.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:42;
- (b) a fragment of the amino acid sequence of SEQ ID NO:42, the  
25 fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:42. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids

of SEQ ID NO:42, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid sequence from amino acid 239 to amino acid 248 of SEQ ID NO:42.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43 from nucleotide 2112 to nucleotide 2363;
- 10 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo2\_1 deposited with the ATCC under accession number PTA-362;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo2\_1 deposited with the ATCC under accession number  
15 PTA-362;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo2\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA  
20 insert of clone vo2\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:44;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment  
25 comprising eight contiguous amino acids of SEQ ID NO:44;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 30 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:43.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:43 from nucleotide 2112 to nucleotide 2363; the nucleotide sequence of the full-length protein coding sequence of clone vo2\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vo2\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo2\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:44.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:43.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and

(ab) the nucleotide sequence of the cDNA insert of clone vo2\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vo2\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:43 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:43, but  
20 excluding the poly(A) tail at the 3' end of SEQ ID NO:43. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43 from nucleotide 2112 to nucleotide 2363, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:43 from nucleotide 2112 to nucleotide 2363,  
25 to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:43 from nucleotide 2112 to nucleotide 2363.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

30 (a) the amino acid sequence of SEQ ID NO:44;

(b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo2\_1 deposited with the ATCC under accession number PTA-362; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:44. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising the amino acid sequence  
10 from amino acid 37 to amino acid 46 of SEQ ID NO:44.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45 from nucleotide 36 to nucleotide 707;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45 from nucleotide 393 to nucleotide 707;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo3\_1 deposited with the ATCC under  
20 accession number PTA-362;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;
- 25 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo3\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;
- 30 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:46;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:45.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:45 from nucleotide 36 to nucleotide 707; the nucleotide sequence of SEQ ID NO:45 from nucleotide 393 to nucleotide 707; the nucleotide sequence of the full-length protein coding sequence of clone vo3\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vo3\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:46, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence from amino acid 107 to amino acid 116 of SEQ ID NO:46.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:45.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (aa) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and

(ab) the nucleotide sequence of the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

10 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

15 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and

20 (bb) the nucleotide sequence of the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:45 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45. Also preferably the  
30 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45 from nucleotide 36 to nucleotide 707, and extending contiguously from a nucleotide sequence corresponding to the 5' end



of said sequence of SEQ ID NO:45 from nucleotide 36 to nucleotide 707, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:45 from nucleotide 36 to nucleotide 707. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
5 NO:45 from nucleotide 393 to nucleotide 707, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:45 from nucleotide 393 to nucleotide 707, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:45 from nucleotide 393 to nucleotide 707.

In other embodiments, the present invention provides a composition comprising  
10 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
- (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
- 15 (c) the amino acid sequence encoded by the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:46. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:46, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence from amino acid 107 to amino acid 116 of SEQ ID NO:46.

25 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:47 from nucleotide 74 to nucleotide 295;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47 from nucleotide 134 to nucleotide 295;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo5\_1 deposited with the ATCC under accession number PTA-362;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo5\_1 deposited with the ATCC under accession number PTA-362;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 20 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least
- 25 25% of the length of SEQ ID NO:47.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:47 from nucleotide 74 to nucleotide 295; the nucleotide sequence of SEQ ID NO:47 from nucleotide 134 to nucleotide 295; the nucleotide sequence of the full-length protein coding sequence of clone vo5\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vo5\_1 deposited with the ATCC under accession number PTA-362. In other preferred

30 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by

the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably  
5 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 32 to amino acid 41 of SEQ ID NO:48.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
10 ID NO:47.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
20 vo5\_1 deposited with the ATCC under accession number PTA-362;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- 25 and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 30 (ba) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and

- (bb) the nucleotide sequence of the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:47 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 74 to nucleotide 295, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 74 to nucleotide 295, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 74 to nucleotide 295. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 134 to nucleotide 295, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 134 to nucleotide 295, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 134 to nucleotide 295.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:48;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:48. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 32 to amino acid 41 of SEQ ID NO:48.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49 from nucleotide 45 to nucleotide 383;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49 from nucleotide 312 to nucleotide 383;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

5 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:49.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:49 from nucleotide 45 to nucleotide 383; the nucleotide sequence of SEQ ID NO:49  
10 from nucleotide 312 to nucleotide 383; the nucleotide sequence of the full-length protein coding sequence of clone vo6\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vo6\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by  
15 the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:50, or a  
20 polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising the amino acid sequence from amino acid 51 to amino acid 60 of SEQ ID NO:50.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:49.

25 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the  
30 group consisting of:

(aa) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and

- (ab) the nucleotide sequence of the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize
- 10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and
- (bb) the nucleotide sequence of the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- 15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:49 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49 from nucleotide 45 to nucleotide 383, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:49 from nucleotide 45 to nucleotide 383, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:49 from nucleotide
- 30 45 to nucleotide 383. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49 from nucleotide 312 to nucleotide 383, and extending contiguously from a

nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:49 from nucleotide 312 to nucleotide 383, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:49 from nucleotide 312 to nucleotide 383.

In other embodiments, the present invention provides a composition comprising  
5 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
- (b) a fragment of the amino acid sequence of SEQ ID NO:50, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
- 10 (c) the amino acid sequence encoded by the cDNA insert of clone  
vo6\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:50. In further preferred  
embodiments, the present invention provides a protein comprising a fragment of the amino  
15 acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably  
comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
of SEQ ID NO:50, or a protein comprising a fragment of the amino acid sequence of SEQ  
ID NO:50 having biological activity, the fragment comprising the amino acid sequence  
from amino acid 51 to amino acid 60 of SEQ ID NO:50.

20 In one embodiment, the present invention provides a composition comprising an  
isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:51;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
25 NO:51 from nucleotide 186 to nucleotide 1739;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:51 from nucleotide 288 to nucleotide 1739;
- (d) a polynucleotide comprising the nucleotide sequence of the full-  
length protein coding sequence of clone vo9\_1 deposited with the ATCC under  
30 accession number PTA-362;



- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:51.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:51 from nucleotide 186 to nucleotide 1739; the nucleotide sequence of SEQ ID NO:51 from nucleotide 288 to nucleotide 1739; the nucleotide sequence of the full-length protein coding sequence of clone vo9\_1 deposited with the ATCC under accession number PTA-362; or the nucleotide sequence of a mature protein coding sequence of clone vo9\_1 deposited with the ATCC under accession number PTA-362. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52

having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising the amino acid  
5 sequence from amino acid 254 to amino acid 263 of SEQ ID NO:52.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:51.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:51, but excluding the poly(A) tail at the  
15 3' end of SEQ ID NO:51; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vo9\_1 deposited with the ATCC under accession number PTA-362;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:51 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 186 to nucleotide  
10 1739, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 186 to nucleotide 1739, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide 186 to nucleotide 1739. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:51 from nucleotide 288 to nucleotide 1739, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 288 to nucleotide 1739, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide 288 to nucleotide 1739.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:52;
- (b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:52. In further preferred  
30 acid sequence of SEQ ID NO:52 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:52 having biological activity, the fragment comprising the amino acid sequence from amino acid 254 to amino acid 263 of SEQ ID NO:52.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5                   (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53 from nucleotide 440 to nucleotide 835;
- (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53 from nucleotide 632 to nucleotide 835;
- 10                  (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo11\_1 deposited with the ATCC under accession number PTA-366;
- (e)     a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo11\_1 deposited with the ATCC under accession number PTA-366;
- 15                  (f)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo11\_1 deposited with the ATCC under accession number PTA-366;
- 20                  (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo11\_1 deposited with the ATCC under accession number PTA-366;
- (h)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:54;
- 25                  (i)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;
- (j)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 30                  (k)     a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:53.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:53 from nucleotide 440 to nucleotide 835; the nucleotide sequence of SEQ ID NO:53 from nucleotide 632 to nucleotide 835; the nucleotide sequence of the full-length protein coding sequence of clone vo11\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo11\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo11\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:54.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:53.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and

- (ab) the nucleotide sequence of the cDNA insert of clone vo11\_1 deposited with the ATCC under accession number PTA-366;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize
- 10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 vo11\_1 deposited with the ATCC under accession number PTA-366;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 20 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:53 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:53, but

25 excluding the poly(A) tail at the 3' end of SEQ ID NO:53. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide 440 to nucleotide 835, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 440 to nucleotide 835, to a nucleotide

30 sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 440 to nucleotide 835. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

NO:53 from nucleotide 632 to nucleotide 835, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 632 to nucleotide 835, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 632 to nucleotide 835.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:54;
- (b) a fragment of the amino acid sequence of SEQ ID NO:54, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:54; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
vol1\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:54. In further preferred  
15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence  
20 from amino acid 61 to amino acid 70 of SEQ ID NO:54.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:55;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:55 from nucleotide 72 to nucleotide 329;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:55 from nucleotide 120 to nucleotide 329;
- 30 (d) a polynucleotide comprising the nucleotide sequence of the full-  
length protein coding sequence of clone vol12\_1 deposited with the ATCC under  
accession number PTA-366;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo12\_1 deposited with the ATCC under accession number PTA-366;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:56;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:55.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:55 from nucleotide 72 to nucleotide 329; the nucleotide sequence of SEQ ID NO:55 from nucleotide 120 to nucleotide 329; the nucleotide sequence of the full-length protein coding sequence of clone vo12\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo12\_1 deposited with the ATCC under accession number PTA-366. In other preferred

30 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide



encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of  
5 SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:56.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:55.

Further embodiments of the invention provide isolated polynucleotides produced  
10 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55; and

(ab) the nucleotide sequence of the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

25 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (ba) SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55; and

- (bb) the nucleotide sequence of the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID

10 NO:55 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 72 to nucleotide 329, and extending contiguously from a nucleotide sequence corresponding to the 5' end

15 of said sequence of SEQ ID NO:55 from nucleotide 72 to nucleotide 329, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide 72 to nucleotide 329. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 120 to nucleotide 329, and extending contiguously from a

20 nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:55 from nucleotide 120 to nucleotide 329, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide 120 to nucleotide 329.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

25 consisting of:

- (a) the amino acid sequence of SEQ ID NO:56;
- (b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and
- (c) the amino acid sequence encoded by the cDNA insert of clone
- 30 vo12\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:56. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:56.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57 from nucleotide 227 to nucleotide 439;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57 from nucleotide 287 to nucleotide 439;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:58;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;
- 30

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:57.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:57 from nucleotide 227 to nucleotide 439; the nucleotide sequence of SEQ ID NO:57 from nucleotide 287 to nucleotide 439; the nucleotide sequence of the full-length protein coding sequence of clone vo13\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo13\_1  
15 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58  
20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:58, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising the amino acid sequence from amino acid 30 to amino acid 39 of SEQ ID NO:58.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:57.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and

(ab) the nucleotide sequence of the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

10 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and

(bb) the nucleotide sequence of the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:57 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57 from nucleotide 227 to nucleotide 439, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:57 from nucleotide 227 to nucleotide 439, to a nucleotide

sequence corresponding to the 3' end of said sequence of SEQ ID NO:57 from nucleotide 227 to nucleotide 439. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57 from nucleotide 287 to nucleotide 439, and extending contiguously from a  
5 nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:57 from nucleotide 287 to nucleotide 439, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:57 from nucleotide 287 to nucleotide 439.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
10 consisting of:

- (a) the amino acid sequence of SEQ ID NO:58;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
15 vo13\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:58. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment preferably  
20 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:58, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising the amino acid sequence from amino acid 30 to amino acid 39 of SEQ ID NO:58.

In one embodiment, the present invention provides a composition comprising an  
25 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59 from nucleotide 96 to nucleotide 341;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:59 from nucleotide 174 to nucleotide 341;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any  
25 one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:59.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:59 from nucleotide 96 to nucleotide 341; the nucleotide sequence of SEQ ID NO:59 from nucleotide 174 to nucleotide 341; the nucleotide sequence of the full-length protein  
30 coding sequence of clone vo14\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo14\_1 deposited with the ATCC under accession number PTA-366. In other preferred

embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:60, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising the amino acid sequence from amino acid 36 to amino acid 45 of SEQ ID NO:60.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:59.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
    - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
      - (aa) SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59; and
      - (ab) the nucleotide sequence of the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
    - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
    - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
    - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:



- (ba) SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59; and
- (bb) the nucleotide sequence of the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- 5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:59 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59. Also preferably the
- 15 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59 from nucleotide 96 to nucleotide 341, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:59 from nucleotide 96 to nucleotide 341, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:59 from nucleotide
- 20 96 to nucleotide 341. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59 from nucleotide 174 to nucleotide 341, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:59 from nucleotide 174 to nucleotide 341, to a nucleotide sequence corresponding to the 3' end of
- 25 said sequence of SEQ ID NO:59 from nucleotide 174 to nucleotide 341.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;
- 30 (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:60. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:60, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising the amino acid sequence  
10 from amino acid 36 to amino acid 45 of SEQ ID NO:60.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61 from nucleotide 90 to nucleotide 599;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61 from nucleotide 165 to nucleotide 599;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo15\_1 deposited with the ATCC under  
20 accession number PTA-366;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366;
- 25 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo15\_1 deposited with the ATCC under accession number PTA-366;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-  
30 366;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:62;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;

5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

10 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:61.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:61 from nucleotide 90 to nucleotide 599; the nucleotide sequence of SEQ ID NO:61 from nucleotide 165 to nucleotide 599; the nucleotide sequence of the full-length protein coding sequence of clone vo15\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo15\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:62, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising the amino acid sequence from amino acid 80 to amino acid 89 of SEQ ID NO:62.

20 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:61.

30 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (aa) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and

(ab) the nucleotide sequence of the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366;

10 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

15 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and

20 (bb) the nucleotide sequence of the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

25 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID  
30 NO:61 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence

corresponding to the cDNA sequence of SEQ ID NO:61 from nucleotide 90 to nucleotide 599, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:61 from nucleotide 90 to nucleotide 599, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:61 from nucleotide 90 to nucleotide 599. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61 from nucleotide 165 to nucleotide 599, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:61 from nucleotide 165 to nucleotide 599, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:61 from nucleotide 165 to nucleotide 599.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:62;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:62. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:62, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising the amino acid sequence from amino acid 80 to amino acid 89 of SEQ ID NO:62.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 209 to nucleotide 451;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 398 to nucleotide 451;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo16\_1 deposited with the ATCC under accession number PTA-366;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo16\_1 deposited with the ATCC under accession number PTA-366;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:64;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:63.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:63 from nucleotide 209 to nucleotide 451; the nucleotide sequence of SEQ ID NO:63 from nucleotide 398 to nucleotide 451; the nucleotide sequence of the full-length protein coding sequence of clone vo16\_1 deposited with the ATCC under accession number PTA-

366; or the nucleotide sequence of a mature protein coding sequence of clone vo16\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-  
5 366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:64, or a  
10 polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:64.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:63.

Further embodiments of the invention provide isolated polynucleotides produced  
15 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (aa) SEQ ID NO:63, but excluding the poly(A) tail at the 3' end of SEQ ID NO:63; and

(ab) the nucleotide sequence of the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;

25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

30 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:63, but excluding the poly(A) tail at the 3' end of SEQ ID NO:63; and

(bb) the nucleotide sequence of the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;

10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63, and  
15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:63 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:63, but excluding the poly(A) tail at the 3' end of SEQ ID NO:63. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63 from nucleotide 209 to nucleotide  
20 451, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:63 from nucleotide 209 to nucleotide 451, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:63 from nucleotide 209 to nucleotide 451. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
25 NO:63 from nucleotide 398 to nucleotide 451, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:63 from nucleotide 398 to nucleotide 451, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:63 from nucleotide 398 to nucleotide 451.

In other embodiments, the present invention provides a composition comprising  
30 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:64;



- (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:64. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:64, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:64.

In one embodiment, the present invention provides a composition comprising an

15 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 31 to nucleotide 231;
- 20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 97 to nucleotide 231;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- 25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo18\_1 deposited with the ATCC under
- 30 accession number PTA-366;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:65.
- 15

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:65 from nucleotide 31 to nucleotide 231; the nucleotide sequence of SEQ ID NO:65 from nucleotide 97 to nucleotide 231; the nucleotide sequence of the full-length protein coding sequence of clone vo18\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo18\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:66, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising the amino acid sequence from amino acid 28 to amino acid 37 of SEQ ID NO:66.

20

25

30

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:65.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5                   (a)     a process comprising the steps of:
- (i)     preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa)    SEQ ID NO:65, but excluding the poly(A) tail at the
- 10                   3' end of SEQ ID NO:65; and
- (ab)    the nucleotide sequence of the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (ii)    hybridizing said probe(s) to human genomic DNA in
- 15                   conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii)   isolating the DNA polynucleotides detected with the probe(s);
- and
- (b)     a process comprising the steps of:
- 20                   (i)     preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba)    SEQ ID NO:65, but excluding the poly(A) tail at the
- 25                   3' end of SEQ ID NO:65; and
- (bb)    the nucleotide sequence of the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (ii)    hybridizing said primer(s) to human genomic DNA in
- conditions at least as stringent as 4X SSC at 50 degrees C;
- 30                   (iii)   amplifying human DNA sequences; and
- (iv)    isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:65 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:65, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:65. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 31 to nucleotide 231, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from nucleotide 31 to nucleotide 231, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:65 from nucleotide 31 to nucleotide 231. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 97 to nucleotide 231, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from  
15 nucleotide 97 to nucleotide 231, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:65 from nucleotide 97 to nucleotide 231.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:66;
- (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:66. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:66, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising the amino acid sequence from amino acid 28 to amino acid 37 of SEQ ID NO:66.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67 from nucleotide 23 to nucleotide 736;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67 from nucleotide 83 to nucleotide 736;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo19\_1 deposited with the ATCC under  
10 accession number PTA-366;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-366;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo19\_1 deposited with the ATCC under accession number PTA-366;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-  
20 366;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment  
25 comprising eight contiguous amino acids of SEQ ID NO:68;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:67.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:67 from nucleotide 23 to nucleotide 736; the nucleotide sequence of SEQ ID NO:67 from nucleotide 83 to nucleotide 736; the nucleotide sequence of the full-length protein coding sequence of clone vo19\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo19\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:68, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising the amino acid sequence from amino acid 114 to amino acid 123 of SEQ ID NO:68.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:67.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

5 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67; and

(bb) the nucleotide sequence of the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-366;

15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
20 nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67, and  
extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID  
NO:67 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:67, but  
excluding the poly(A) tail at the 3' end of SEQ ID NO:67. Also preferably the  
polynucleotide isolated according to the above process comprises a nucleotide sequence  
25 corresponding to the cDNA sequence of SEQ ID NO:67 from nucleotide 23 to nucleotide  
736, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
of said sequence of SEQ ID NO:67 from nucleotide 23 to nucleotide 736, to a nucleotide  
sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide  
23 to nucleotide 736. Also preferably the polynucleotide isolated according to the above  
30 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
NO:67 from nucleotide 83 to nucleotide 736, and extending contiguously from a  
nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:67 from

nucleotide 83 to nucleotide 736, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide 83 to nucleotide 736.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
10 vo19\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:68. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably  
15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:68, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising the amino acid sequence from amino acid 114 to amino acid 123 of SEQ ID NO:68.

In one embodiment, the present invention provides a composition comprising an  
20 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69 from nucleotide 104 to nucleotide 1399;
- 25 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69 from nucleotide 158 to nucleotide 1399;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo22\_1 deposited with the ATCC under accession number PTA-366;
- 30 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo22\_1 deposited with the ATCC under accession number PTA-366;



(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo22\_1 deposited with the ATCC under accession number PTA-366;

5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo22\_1 deposited with the ATCC under accession number PTA-366;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:70;

10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

15 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:69.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:69 from nucleotide 104 to nucleotide 1399; the nucleotide sequence of SEQ ID NO:69 from nucleotide 158 to nucleotide 1399; the nucleotide sequence of the full-length protein coding sequence of clone vo22\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo22\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo22\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a

polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising the amino acid sequence from amino acid 211 to amino acid 220 of SEQ ID NO:70.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
5 ID NO:69.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
15 vo22\_1 deposited with the ATCC under accession number PTA-366;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the  
20 probe(s);

and

- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vo22\_1 deposited with the ATCC under accession number PTA-366;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:69 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69 from nucleotide  
10 104 to nucleotide 1399, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:69 from nucleotide 104 to nucleotide 1399, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:69 from nucleotide 104 to nucleotide 1399. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:69 from nucleotide 158 to nucleotide 1399, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:69 from nucleotide 158 to nucleotide 1399, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:69 from nucleotide 158 to nucleotide 1399.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:70;
- (b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone vo22\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:70. In further preferred  
30 acid sequence of SEQ ID NO:70 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:70 having biological activity, the fragment comprising the amino acid sequence from amino acid 211 to amino acid 220 of SEQ ID NO:70.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5                   (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71 from nucleotide 174 to nucleotide 1595;
- (c)     a polynucleotide comprising the nucleotide sequence of the full-  
10     length protein coding sequence of clone vo23\_1 deposited with the ATCC under accession number PTA-366;
- (d)     a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-366;
- 15               (e)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo23\_1 deposited with the ATCC under accession number PTA-366;
- (f)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-  
20     366;
- (g)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:72;
- (h)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment  
25     comprising eight contiguous amino acids of SEQ ID NO:72;
- (i)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j)     a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 30               (k)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:71.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:71 from nucleotide 174 to nucleotide 1595; the nucleotide sequence of the full-length protein coding sequence of clone vo23\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo23\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:72, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 232 to amino acid 241 of SEQ ID NO:72.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:71.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and

(ab) the nucleotide sequence of the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

15 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:71 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71 from nucleotide 174 to nucleotide 1595, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:71 from nucleotide 174 to nucleotide 1595, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:71 from nucleotide 174 to nucleotide 1595.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

30 (a) the amino acid sequence of SEQ ID NO:72;

(b) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-366;

5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:72. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
10 of SEQ ID NO:72, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 232 to amino acid 241 of SEQ ID NO:72.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 129 to nucleotide 311;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
20 NO:73 from nucleotide 195 to nucleotide 311;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(e) a polynucleotide encoding the full-length protein encoded by the  
25 cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(g) a polynucleotide encoding a mature protein encoded by the cDNA  
30 insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:74;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:73.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:73 from nucleotide 129 to nucleotide 311; the nucleotide sequence of SEQ ID NO:73 from nucleotide 195 to nucleotide 311; the nucleotide sequence of the full-length protein coding sequence of clone vo24\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo24\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:74.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:73.



Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

(ab) the nucleotide sequence of the cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

(bb) the nucleotide sequence of the cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID

NO:73 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:73 , but excluding the poly(A) tail at the 3' end of SEQ ID NO:73. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 129 to nucleotide

5 311, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 129 to nucleotide 311, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 129 to nucleotide 311. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

10 NO:73 from nucleotide 195 to nucleotide 311, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 195 to nucleotide 311, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 195 to nucleotide 311.

In other embodiments, the present invention provides a composition comprising

15 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
- (b) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and
- 20 (c) the amino acid sequence encoded by the cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:74. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino

25 acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:74.

30 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 73 to nucleotide 798;
- 5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 142 to nucleotide 798;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- 20 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 25 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:75.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:75 from nucleotide 73 to nucleotide 798; the nucleotide sequence of SEQ ID NO:75 from nucleotide 142 to nucleotide 798; the nucleotide sequence of the full-length protein coding sequence of clone vo25\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo25\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 116 to amino acid 125 of SEQ ID NO:76.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:75.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:75, but excluding the poly(A) tail at the  
25 3' end of SEQ ID NO:75; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
  - (ii) hybridizing said probe(s) to human genomic DNA in  
30 conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and

(bb) the nucleotide sequence of the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

15 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:75 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75. Also preferably the  
20 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 73 to nucleotide 798, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 73 to nucleotide 798, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide  
25 73 to nucleotide 798. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 142 to nucleotide 798, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 142 to nucleotide 798, to a nucleotide sequence corresponding to the 3' end of  
30 said sequence of SEQ ID NO:75 from nucleotide 142 to nucleotide 798.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins. Preferably such
- 10 protein comprises the amino acid sequence of SEQ ID NO:76. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a protein comprising a fragment of the amino acid sequence of SEQ
- 15 ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 116 to amino acid 125 of SEQ ID NO:76.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:77;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 26 to nucleotide 307;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 101 to nucleotide 307;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo26\_1 deposited with the ATCC under accession number
- 30 PTA-366;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:77.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:77 from nucleotide 26 to nucleotide 307; the nucleotide sequence of SEQ ID NO:77 from nucleotide 101 to nucleotide 307; the nucleotide sequence of the full-length protein coding sequence of clone vo26\_1 deposited with the ATCC under accession number PTA-366; or the nucleotide sequence of a mature protein coding sequence of clone vo26\_1 deposited with the ATCC under accession number PTA-366. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo26\_1 deposited with the ATCC under accession number PTA-366. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:78, or a

polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 42 to amino acid 51 of SEQ ID NO:78.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
5 ID NO:77.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
15 vo26\_1 deposited with the ATCC under accession number PTA-366;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the  
20 probe(s);

and

- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vo26\_1 deposited with the ATCC under accession number PTA-366;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;



- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:77 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77 from nucleotide 26 to nucleotide  
10 307, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 26 to nucleotide 307, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from nucleotide 26 to nucleotide 307. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:77 from nucleotide 101 to nucleotide 307, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 101 to nucleotide 307, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from nucleotide 101 to nucleotide 307.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
- (b) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone vo26\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:78. In further preferred  
30 acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:78, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 42 to amino acid 51 of SEQ ID NO:78.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79 from nucleotide 43 to nucleotide 228;
- (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID  
10       NO:79 from nucleotide 94 to nucleotide 228;
- (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vp23\_1 deposited with the ATCC under accession number PTA-368;
- (e)     a polynucleotide encoding the full-length protein encoded by the  
15       cDNA insert of clone vp23\_1 deposited with the ATCC under accession number PTA-368;
- (f)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vp23\_1 deposited with the ATCC under accession number PTA-368;
- 20       (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vp23\_1 deposited with the ATCC under accession number PTA-368;
- (h)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
- 25       (i)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;
- (j)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 30       (k)     a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least  
5 25% of the length of SEQ ID NO:79.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:79 from nucleotide 43 to nucleotide 228; the nucleotide sequence of SEQ ID NO:79 from nucleotide 94 to nucleotide 228; the nucleotide sequence of the full-length protein coding sequence of clone vp23\_1 deposited with the ATCC under accession number PTA-  
10 368; or the nucleotide sequence of a mature protein coding sequence of clone vp23\_1 deposited with the ATCC under accession number PTA-368. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vp23\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide  
15 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid  
20 sequence from amino acid 26 to amino acid 35 of SEQ ID NO:80.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:79.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

25 (a) a process comprising the steps of:  
(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:  
(aa) SEQ ID NO:79, but excluding the poly(A) tail at the  
30 3' end of SEQ ID NO:79; and

- (ab) the nucleotide sequence of the cDNA insert of clone  
vp23\_1 deposited with the ATCC under accession number PTA-368;
- (ii) hybridizing said probe(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the  
probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the  
group consisting of:
- (ba) SEQ ID NO:79, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:79; and
- (bb) the nucleotide sequence of the cDNA insert of clone  
15 vp23\_1 deposited with the ATCC under accession number PTA-  
368;
- (ii) hybridizing said primer(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 20 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79, and  
extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID  
NO:79 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:79, but  
25 excluding the poly(A) tail at the 3' end of SEQ ID NO:79. Also preferably the  
polynucleotide isolated according to the above process comprises a nucleotide sequence  
corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 43 to nucleotide  
228, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
of said sequence of SEQ ID NO:79 from nucleotide 43 to nucleotide 228, to a nucleotide  
30 sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide  
43 to nucleotide 228. Also preferably the polynucleotide isolated according to the above  
process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

NO:79 from nucleotide 94 to nucleotide 228, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from nucleotide 94 to nucleotide 228, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide 94 to nucleotide 228.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:80;
- (b) a fragment of the amino acid sequence of SEQ ID NO:80, the
- 10 fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp23\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:80. In further preferred

15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence

20 from amino acid 26 to amino acid 35 of SEQ ID NO:80.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:81;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:81 from nucleotide 245 to nucleotide 427;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:81 from nucleotide 308 to nucleotide 427;
- (d) a polynucleotide comprising the nucleotide sequence of the full-
- 30 length protein coding sequence of clone vq7\_1 deposited with the ATCC under accession number PTA-368;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq7\_1 deposited with the ATCC under accession number PTA-368;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:82;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:82;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:81.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:81 from nucleotide 245 to nucleotide 427; the nucleotide sequence of SEQ ID NO:81  
25 from nucleotide 308 to nucleotide 427; the nucleotide sequence of the full-length protein coding sequence of clone vq7\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq7\_1 deposited with the ATCC under accession number PTA-368. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by  
30 the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82

having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:82, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment comprising the amino acid  
 5 sequence from amino acid 25 to amino acid 34 of SEQ ID NO:82.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:81.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:81, but excluding the poly(A) tail at the  
15 3' end of SEQ ID NO:81; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vq7\_1 deposited with the ATCC under accession number PTA-368;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:81 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81 from nucleotide 245 to nucleotide  
10 427, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:81 from nucleotide 245 to nucleotide 427, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:81 from nucleotide 245 to nucleotide 427. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:81 from nucleotide 308 to nucleotide 427, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:81 from nucleotide 308 to nucleotide 427, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:81 from nucleotide 308 to nucleotide 427.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of: \_\_\_\_\_

- (a) the amino acid sequence of SEQ ID NO:82;
- (b) a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:82. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino  
30 acid sequence of SEQ ID NO:82 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:82, or a protein comprising a fragment of the amino acid sequence of SEQ



ID NO:82 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:82.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5                   (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83 from nucleotide 119 to nucleotide 475;
- (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID  
10               NO:83 from nucleotide 185 to nucleotide 475;
- (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- (e)     a polynucleotide encoding the full-length protein encoded by the  
15               cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- (f)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- 20               (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- (h)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:84;
- (i)     a polynucleotide encoding a protein comprising a fragment of the  
25               amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:84;
- (j)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k)     a polynucleotide which encodes a species homologue of the protein  
30               of (h) or (i) above ;
- (l)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:83.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:83 from nucleotide 119 to nucleotide 475; the nucleotide sequence of SEQ ID NO:83 from nucleotide 185 to nucleotide 475; the nucleotide sequence of the full-length protein coding sequence of clone vq8\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq8\_1 deposited with the ATCC under accession number PTA-368. In other preferred  
10 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment preferably comprising eight (more preferably  
15 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:84, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:84.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:83.  
20

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:83, but excluding the poly(A) tail at the 3' end of SEQ ID NO:83; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
30 vq8\_1 deposited with the ATCC under accession number PTA-368;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:83, but excluding the poly(A) tail at the 3' end of SEQ ID NO:83; and

10 (bb) the nucleotide sequence of the cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:83 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:83, but  
20 excluding the poly(A) tail at the 3' end of SEQ ID NO:83. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83 from nucleotide 119 to nucleotide 475, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:83 from nucleotide 119 to nucleotide 475, to a nucleotide  
25 sequence corresponding to the 3' end of said sequence of SEQ ID NO:83 from nucleotide 119 to nucleotide 475. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83 from nucleotide 185 to nucleotide 475, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:83 from  
30 nucleotide 185 to nucleotide 475, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:83 from nucleotide 185 to nucleotide 475.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:84;
- 5 (b) a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins. Preferably such  
10 protein comprises the amino acid sequence of SEQ ID NO:84. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:84, or a protein comprising a fragment of the amino acid sequence of SEQ  
15 ID NO:84 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:84.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
20 NO:85;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:85 from nucleotide 90 to nucleotide 323;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:85 from nucleotide 141 to nucleotide 323;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq9\_1 deposited with the ATCC under accession number PTA-368;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number  
30 PTA-368;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq9\_1 deposited with the ATCC under accession number PTA-368;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number PTA-368;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:86;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:86;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:85.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:85 from nucleotide 90 to nucleotide 323; the nucleotide sequence of SEQ ID NO:85 from nucleotide 141 to nucleotide 323; the nucleotide sequence of the full-length protein coding sequence of clone vq9\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq9\_1 deposited with the ATCC under accession number PTA-368. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:86, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of

SEQ ID NO:86 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:86.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:85.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85; and

(ab) the nucleotide sequence of the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number PTA-368;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85; and

(bb) the nucleotide sequence of the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number PTA-368;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:85, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:85 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:85, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:85. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:85 from nucleotide 90 to nucleotide 323, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:85 from nucleotide 90 to nucleotide 323, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:85 from nucleotide 90 to nucleotide 323. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:85 from nucleotide 141 to nucleotide 323, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:85 from  
15 nucleotide 141 to nucleotide 323, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:85 from nucleotide 141 to nucleotide 323.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:86;  
(b) a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86; and  
(c) the amino acid sequence encoded by the cDNA insert of clone  
vq9\_1 deposited with the ATCC under accession number PTA-368;  
25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:86. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:86, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:86.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 18 to nucleotide 452;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 72 to nucleotide 452;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq10\_1 deposited with the ATCC under accession number PTA-368;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq10\_1 deposited with the ATCC under accession number PTA-368;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:88;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and



(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:87.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:87 from nucleotide 18 to nucleotide 452; the nucleotide sequence of SEQ ID NO:87 from nucleotide 72 to nucleotide 452; the nucleotide sequence of the full-length protein coding sequence of clone vq10\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq10\_1 deposited with the ATCC under accession number PTA-368. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 67 to amino acid 76 of SEQ ID NO:88.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:87.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

5 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87; and

(bb) the nucleotide sequence of the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368;

15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
20 nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:87 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
25 corresponding to the cDNA sequence of SEQ ID NO:87 from nucleotide 18 to nucleotide 452, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:87 from nucleotide 18 to nucleotide 452, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:87 from nucleotide 18 to nucleotide 452. Also preferably the polynucleotide isolated according to the above  
30 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87 from nucleotide 72 to nucleotide 452, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:87 from

nucleotide 72 to nucleotide 452, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:87 from nucleotide 72 to nucleotide 452.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:88;
- (b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and
- (c) the amino acid sequence encoded by the cDNA insert of clone

10 vq10\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:88. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably

15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 67 to amino acid 76 of SEQ ID NO:88.

In one embodiment, the present invention provides a composition comprising an

20 isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89 from nucleotide 196 to nucleotide 378;

25 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89 from nucleotide 262 to nucleotide 378;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

30 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq13\_1 deposited with the ATCC under accession number PTA-368;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq13\_1 deposited with the ATCC under accession number PTA-368;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:90;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:90;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:89.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:89 from nucleotide 196 to nucleotide 378; the nucleotide sequence of SEQ ID NO:89 from nucleotide 262 to nucleotide 378; the nucleotide sequence of the full-length protein coding sequence of clone vq13\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq13\_1 deposited with the ATCC under accession number PTA-368. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq13\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:90, or a

polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:90.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
5 ID NO:89.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
15 vq13\_1 deposited with the ATCC under accession number PTA-368;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the  
20 probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vq13\_1 deposited with the ATCC under accession number PTA-368;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:89 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89 from nucleotide 196 to nucleotide  
10 378, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:89 from nucleotide 196 to nucleotide 378, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:89 from nucleotide 196 to nucleotide 378. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:89 from nucleotide 262 to nucleotide 378, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:89 from nucleotide 262 to nucleotide 378, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:89 from nucleotide 262 to nucleotide 378.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:90;
- (b) a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:90. In further preferred  
30 acid sequence of SEQ ID NO:90 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:90, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:90 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:90.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5                   (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91 from nucleotide 35 to nucleotide 718;
- (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID  
10               NO:91 from nucleotide 173 to nucleotide 718;
- (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- (e)     a polynucleotide encoding the full-length protein encoded by the  
15               cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- (f)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- 20               (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- (h)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:92;
- 25               (i)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:92;
- (j)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 30               (k)     a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least  
5 25% of the length of SEQ ID NO:91.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:91 from nucleotide 35 to nucleotide 718; the nucleotide sequence of SEQ ID NO:91 from nucleotide 173 to nucleotide 718; the nucleotide sequence of the full-length protein coding sequence of clone vq16\_1 deposited with the ATCC under accession number PTA-  
10 368; or the nucleotide sequence of a mature protein coding sequence of clone vq16\_1 deposited with the ATCC under accession number PTA-368. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide  
15 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:92, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment comprising the amino acid  
20 sequence from amino acid 109 to amino acid 118 of SEQ ID NO:92.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:91.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 25 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:91, but excluding the poly(A) tail at the  
30 3' end of SEQ ID NO:91; and



- (ab) the nucleotide sequence of the cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
    - (i) preparing one or more polynucleotide primers that hybridize
    - 10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
      - (ba) SEQ ID NO:91, but excluding the poly(A) tail at the 3' end of SEQ ID NO:91; and
      - (bb) the nucleotide sequence of the cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
    - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
    - (iii) amplifying human DNA sequences; and
    - 20 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:91, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:91 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:91, but

25 excluding the poly(A) tail at the 3' end of SEQ ID NO:91. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:91 from nucleotide 35 to nucleotide 718, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:91 from nucleotide 35 to nucleotide 718, to a nucleotide

30 sequence corresponding to the 3' end of said sequence of SEQ ID NO:91 from nucleotide 35 to nucleotide 718. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

NO:91 from nucleotide 173 to nucleotide 718, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:91 from nucleotide 173 to nucleotide 718, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:91 from nucleotide 173 to nucleotide 718.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:92;
- (b) a fragment of the amino acid sequence of SEQ ID NO:92, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:92; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
vq16\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:92. In further preferred  
15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:92, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment comprising the amino acid sequence  
20 from amino acid 109 to amino acid 118 of SEQ ID NO:92.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:93;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:93 from nucleotide 1 to nucleotide 762;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:93 from nucleotide 70 to nucleotide 762;
- (d) a polynucleotide comprising the nucleotide sequence of the full-  
30 length protein coding sequence of clone vq19\_1 deposited with the ATCC under  
accession number PTA-368;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:94;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any  
20 one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:93.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID  
25 NO:93 from nucleotide 1 to nucleotide 762; the nucleotide sequence of SEQ ID NO:93 from nucleotide 70 to nucleotide 762; the nucleotide sequence of the full-length protein coding sequence of clone vq19\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq19\_1 deposited with the ATCC under accession number PTA-368. In other preferred  
30 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide

encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of  
5 SEQ ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 122 to amino acid 131 of SEQ ID NO:94.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:93.

Further embodiments of the invention provide isolated polynucleotides produced  
10 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and

(ab) the nucleotide sequence of the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

25 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (ba) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and

- (bb) the nucleotide sequence of the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID  
10 NO:93 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93 from nucleotide 1 to nucleotide 762, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
15 of said sequence of SEQ ID NO:93 from nucleotide 1 to nucleotide 762, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:93 from nucleotide 1 to nucleotide 762. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93 from nucleotide 70 to nucleotide 762, and extending contiguously from a  
20 nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:93 from nucleotide 70 to nucleotide 762, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:93 from nucleotide 70 to nucleotide 762.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
25 consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
30 vq19\_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:94. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 122 to amino acid 131 of SEQ ID NO:94.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95 from nucleotide 106 to nucleotide 792;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95 from nucleotide 172 to nucleotide 792;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq20\_1 deposited with the ATCC under accession number PTA-368;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq20\_1 deposited with the ATCC under accession number PTA-368;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:96;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:96;
- 30

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:95.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:95 from nucleotide 106 to nucleotide 792; the nucleotide sequence of SEQ ID NO:95 from nucleotide 172 to nucleotide 792; the nucleotide sequence of the full-length protein coding sequence of clone vq20\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq20\_1  
15 deposited with the ATCC under accession number PTA-368. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96  
20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:96, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising the amino acid sequence from amino acid 109 to amino acid 118 of SEQ ID NO:96.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:95.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and

(ab) the nucleotide sequence of the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

10 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and

(bb) the nucleotide sequence of the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:95 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95 from nucleotide 106 to nucleotide 792, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:95 from nucleotide 106 to nucleotide 792, to a nucleotide



sequence corresponding to the 3' end of said sequence of SEQ ID NO:95 from nucleotide 106 to nucleotide 792. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95 from nucleotide 172 to nucleotide 792, and extending contiguously from a  
5 nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:95 from nucleotide 172 to nucleotide 792, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:95 from nucleotide 172 to nucleotide 792.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
10 consisting of:

- (a) the amino acid sequence of SEQ ID NO:96;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
15 vq20\_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:96. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment preferably  
20 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:96, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising the amino acid sequence from amino acid 109 to amino acid 118 of SEQ ID NO:96.

In one embodiment, the present invention provides a composition comprising an  
25 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97 from nucleotide 40 to nucleotide 315;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:97 from nucleotide 124 to nucleotide 315;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq21\_1 deposited with the ATCC under accession number PTA-368;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq21\_1 deposited with the ATCC under accession number PTA-368;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:98;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:98;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:97.
- 25

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:97 from nucleotide 40 to nucleotide 315; the nucleotide sequence of SEQ ID NO:97 from nucleotide 124 to nucleotide 315; the nucleotide sequence of the full-length protein coding sequence of clone vq21\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vq21\_1 deposited with the ATCC under accession number PTA-368. In other preferred

30

embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98  
5 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:98, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising the amino acid sequence from amino acid 41 to amino acid 50 of SEQ ID NO:98.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:97.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - 15 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:97, but excluding the poly(A) tail at the 3' end of SEQ ID NO:97; and
    - 20 (ab) the nucleotide sequence of the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize  
30 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:97, but excluding the poly(A) tail at the 3' end of SEQ ID NO:97; and
- (bb) the nucleotide sequence of the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368;
- 5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:97 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:97, but excluding the poly(A) tail at the 3' end of SEQ ID NO:97. Also preferably the
- 15 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97 from nucleotide 40 to nucleotide 315, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:97 from nucleotide 40 to nucleotide 315, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:97 from nucleotide
- 20 40 to nucleotide 315. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97 from nucleotide 124 to nucleotide 315, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:97 from nucleotide 124 to nucleotide 315, to a nucleotide sequence corresponding to the 3' end of
- 25 said sequence of SEQ ID NO:97 from nucleotide 124 to nucleotide 315.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:98;
- 30 (b) a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98; and

(c) the amino acid sequence encoded by the cDNA insert of clone  
vq21\_1 deposited with the ATCC under accession number PTA-368;  
the protein being substantially free from other mammalian proteins. Preferably such  
protein comprises the amino acid sequence of SEQ ID NO:98. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino  
acid sequence of SEQ ID NO:98 having biological activity, the fragment preferably  
comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
of SEQ ID NO:98, or a protein comprising a fragment of the amino acid sequence of SEQ  
ID NO:98 having biological activity, the fragment comprising the amino acid sequence  
10 from amino acid 41 to amino acid 50 of SEQ ID NO:98.

In one embodiment, the present invention provides a composition comprising an  
isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:99;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:99 from nucleotide 70 to nucleotide 699;
- (c) a polynucleotide comprising the nucleotide sequence of the full-  
length protein coding sequence of clone vr2\_1 deposited with the ATCC under  
accession number PTA-368;
- 20 (d) a polynucleotide encoding the full-length protein encoded by the  
cDNA insert of clone vr2\_1 deposited with the ATCC under accession number  
PTA-368;
- (e) a polynucleotide comprising the nucleotide sequence of a mature  
protein coding sequence of clone vr2\_1 deposited with the ATCC under accession  
25 number PTA-368;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA  
insert of clone vr2\_1 deposited with the ATCC under accession number PTA-368;
- (g) a polynucleotide encoding a protein comprising the amino acid  
sequence of SEQ ID NO:100;
- 30 (h) a polynucleotide encoding a protein comprising a fragment of the  
amino acid sequence of SEQ ID NO:100 having biological activity, the fragment  
comprising eight contiguous amino acids of SEQ ID NO:100;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

5 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:99.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:99 from nucleotide 70 to nucleotide 699; the nucleotide sequence of the full-length protein coding sequence of clone vr2\_1 deposited with the ATCC under accession number PTA-368; or the nucleotide sequence of a mature protein coding sequence of clone vr2\_1 deposited with the ATCC under accession number PTA-368. In other preferred  
15 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vr2\_1 deposited with the ATCC under accession number PTA-368. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment preferably comprising eight (more preferably  
20 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:100, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising the amino acid sequence from amino acid 100 to amino acid 109 of SEQ ID NO:100.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
25 ID NO:99.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
30 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99; and
- (ab) the nucleotide sequence of the cDNA insert of clone vr2\_1 deposited with the ATCC under accession number PTA-368;
- 5 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- 10 (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:99, but excluding the poly(A) tail at the 15 3' end of SEQ ID NO:99; and
- (bb) the nucleotide sequence of the cDNA insert of clone vr2\_1 deposited with the ATCC under accession number PTA-368;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 20 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:99, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID

25 NO:99 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:99 from nucleotide 70 to nucleotide 699, and extending contiguously from a nucleotide sequence corresponding to the 5' end

30 of said sequence of SEQ ID NO:99 from nucleotide 70 to nucleotide 699, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:99 from nucleotide 70 to nucleotide 699.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
- 5 (b) a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vr2\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins. Preferably such  
10 protein comprises the amino acid sequence of SEQ ID NO:100. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:100, or a protein comprising a fragment of the amino acid sequence of SEQ  
15 ID NO:100 having biological activity, the fragment comprising the amino acid sequence from amino acid 100 to amino acid 109 of SEQ ID NO:100.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
20 NO:101;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101 from nucleotide 170 to nucleotide 394;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101 from nucleotide 227 to nucleotide 394;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a polynucleotide encoding the full-length protein encoded by the  
cDNA insert of clone vc69\_1 deposited with the ATCC under accession number  
30 PTA-1075;



- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;
- 5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- 10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:102;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein  
15 of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least  
20 25% of the length of SEQ ID NO:101.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:101 from nucleotide 170 to nucleotide 394; the nucleotide sequence of SEQ ID NO:101 from nucleotide 227 to nucleotide 394; the nucleotide sequence of the full-length protein coding sequence of clone vc69\_1 deposited with the ATCC under accession  
25 number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vc69\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc69\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a  
30 polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID

NO:102, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino acid 32 to amino acid 41 of SEQ ID NO:102.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
5 ID NO:101.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101; and
    - (ab) the nucleotide sequence of the cDNA insert of clone  
15 vc69\_1 deposited with the ATCC under accession number PTA-1075;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the  
20 probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vc69\_1 deposited with the ATCC under accession number PTA-1075;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:101 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101 from nucleotide 170 to  
10 nucleotide 394, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:101 from nucleotide 170 to nucleotide 394, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:101 from nucleotide 170 to nucleotide 394. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the  
15 cDNA sequence of SEQ ID NO:101 from nucleotide 227 to nucleotide 394, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:101 from nucleotide 227 to nucleotide 394, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:101 from nucleotide 227 to nucleotide 394.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;
- (b) a fragment of the amino acid sequence of SEQ ID NO:102, the  
25 fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:102. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids

of SEQ ID NO: 102, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO: 102 having biological activity, the fragment comprising the amino acid sequence from amino acid 32 to amino acid 41 of SEQ ID NO: 102.

In one embodiment, the present invention provides a composition comprising an  
5 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO: 103;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO: 103 from nucleotide 43 to nucleotide 198;
- 10 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO: 103 from nucleotide 85 to nucleotide 198;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- 15 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vc71\_1 deposited with the ATCC under  
20 accession number PTA-1075;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a polynucleotide encoding a protein comprising the amino acid  
25 sequence of SEQ ID NO: 104;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO: 104 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO: 104;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of  
30 (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least  
5 25% of the length of SEQ ID NO:103.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:103 from nucleotide 43 to nucleotide 198; the nucleotide sequence of SEQ ID NO:103 from nucleotide 85 to nucleotide 198; the nucleotide sequence of the full-length protein coding sequence of clone vc71\_1 deposited with the ATCC under accession number PTA-  
10 1075; or the nucleotide sequence of a mature protein coding sequence of clone vc71\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a polynucleotide  
15 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:104, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid  
20 sequence from amino acid 21 to amino acid 30 of SEQ ID NO:104.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:103.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

25 (a) a process comprising the steps of:  
(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:  
(aa) SEQ ID NO:103, but excluding the poly(A) tail at  
30 the 3' end of SEQ ID NO:103; and

- (ab) the nucleotide sequence of the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize
- 10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 20 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:103 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:103, but

25 excluding the poly(A) tail at the 3' end of SEQ ID NO:103. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103 from nucleotide 43 to nucleotide 198, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:103 from nucleotide 43 to nucleotide 198, to a nucleotide

30 sequence corresponding to the 3' end of said sequence of SEQ ID NO:103 from nucleotide 43 to nucleotide 198. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

NO:103 from nucleotide 85 to nucleotide 198, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:103 from nucleotide 85 to nucleotide 198, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:103 from nucleotide 85 to nucleotide 198.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
- (b) a fragment of the amino acid sequence of SEQ ID NO:104, the
- 10 fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:104. In further preferred

15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:104, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid sequence

20 from amino acid 21 to amino acid 30 of SEQ ID NO:104.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105 from nucleotide 260 to nucleotide 1552;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105 from nucleotide 335 to nucleotide 1552;
- (d) a polynucleotide comprising the nucleotide sequence of the full-
- 30 length protein coding sequence of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:106;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any  
20 one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:105.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID  
25 NO:105 from nucleotide 260 to nucleotide 1552; the nucleotide sequence of SEQ ID NO:105 from nucleotide 335 to nucleotide 1552; the nucleotide sequence of the full-length protein coding sequence of clone vo27\_1 deposited with the ATCC under accession number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vo27\_1 deposited with the ATCC under accession number PTA-1075. In other  
30 preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a



polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:106, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 210 to amino acid 219 of SEQ ID NO:106.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:105.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and

(ab) the nucleotide sequence of the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and

(bb) the nucleotide sequence of the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

5 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID  
10 NO:105 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105 from nucleotide 260 to nucleotide 1552, and extending contiguously from a nucleotide sequence corresponding  
15 to the 5' end of said sequence of SEQ ID NO:105 from nucleotide 260 to nucleotide 1552, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:105 from nucleotide 260 to nucleotide 1552. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105 from nucleotide 335 to nucleotide 1552, and  
20 extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:105 from nucleotide 335 to nucleotide 1552, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:105 from nucleotide 335 to nucleotide 1552.

In other embodiments, the present invention provides a composition comprising  
25 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:106;

(b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and

30 (c) the amino acid sequence encoded by the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:106. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably  
5 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:106, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 210 to amino acid 219 of SEQ ID NO:106.

In one embodiment, the present invention provides a composition comprising an  
10 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107 from nucleotide 15 to nucleotide 320;
- 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107 from nucleotide 72 to nucleotide 320;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;
- 20 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo31\_1 deposited with the ATCC under  
25 accession number PTA-1075;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a polynucleotide encoding a protein comprising the amino acid  
30 sequence of SEQ ID NO:108;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:108;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:107.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:107 from nucleotide 15 to nucleotide 320; the nucleotide sequence of SEQ ID NO:107 from nucleotide 72 to nucleotide 320; the nucleotide sequence of the full-length protein coding sequence of clone vo31\_1 deposited with the ATCC under accession number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vo31\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:108, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:108.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:107.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (aa) SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107; and

(ab) the nucleotide sequence of the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;

10 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

15 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107; and

20 (bb) the nucleotide sequence of the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

25 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:107, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID  
30 NO:107 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence

corresponding to the cDNA sequence of SEQ ID NO:107 from nucleotide 15 to nucleotide 320, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:107 from nucleotide 15 to nucleotide 320, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:107 from nucleotide 15 to nucleotide 320. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:107 from nucleotide 72 to nucleotide 320, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:107 from nucleotide 72 to nucleotide 320, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:107 from nucleotide 72 to nucleotide 320.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:108;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:108. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:108, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:108.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109 from nucleotide 38 to nucleotide 1255;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109 from nucleotide 86 to nucleotide 1255;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:110;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:110;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:109.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:109 from nucleotide 38 to nucleotide 1255; the nucleotide sequence of SEQ ID NO:109 from nucleotide 86 to nucleotide 1255; the nucleotide sequence of the full-length protein coding sequence of clone vo32\_1 deposited with the ATCC under accession

number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vo32\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo32\_1 deposited with the ATCC under accession  
5 number PTA-1075. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:110, or a polynucleotide encoding a protein comprising a fragment of the amino acid  
10 sequence of SEQ ID NO:110 having biological activity, the fragment comprising the amino acid sequence from amino acid 198 to amino acid 207 of SEQ ID NO:110.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:109.

Further embodiments of the invention provide isolated polynucleotides produced  
15 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (aa) SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109; and

(ab) the nucleotide sequence of the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;

25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

30 (b) a process comprising the steps of:



(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109; and
- (bb) the nucleotide sequence of the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- 10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:109, and

15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:109 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:109 from nucleotide

20 1255, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:109 from nucleotide 38 to nucleotide 1255, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:109 from nucleotide 38 to nucleotide 1255. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

25 NO:109 from nucleotide 86 to nucleotide 1255, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:109 from nucleotide 86 to nucleotide 1255, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:109 from nucleotide 86 to nucleotide 1255.

In other embodiments, the present invention provides a composition comprising

30 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:110;

- (b) a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:110. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:110, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising the amino acid sequence from amino acid 198 to amino acid 207 of SEQ ID NO:110.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111 from nucleotide 80 to nucleotide 1276;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:111 from nucleotide 131 to nucleotide 1276;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;

- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:112;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:111.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:111 from nucleotide 80 to nucleotide 1276; the nucleotide sequence of SEQ ID NO:111 from nucleotide 131 to nucleotide 1276; the nucleotide sequence of the full-length protein coding sequence of clone vo33\_1 deposited with the ATCC under accession number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vo33\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:112, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising the amino acid sequence from amino acid 194 to amino acid 203 of SEQ ID NO:112.
- Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:111.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111; and

(ab) the nucleotide sequence of the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111; and

(bb) the nucleotide sequence of the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID

NO:111 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111 from nucleotide 80 to nucleotide 1276, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 80 to nucleotide 1276, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 80 to nucleotide 1276. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111 from nucleotide 131 to nucleotide 1276, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 131 to nucleotide 1276, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 131 to nucleotide 1276.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:112;
- (b) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:112. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:112, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising the amino acid sequence from amino acid 194 to amino acid 203 of SEQ ID NO:112.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113 from nucleotide 202 to nucleotide 429;
- 5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113 from nucleotide 292 to nucleotide 429;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:114;
- 20 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:114;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 25 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:113.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:113 from nucleotide 202 to nucleotide 429; the nucleotide sequence of SEQ ID NO:113 from nucleotide 292 to nucleotide 429; the nucleotide sequence of the full-length protein coding sequence of clone vq23\_1 deposited with the ATCC under accession number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vq23\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:114, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising the amino acid sequence from amino acid 33 to amino acid 42 of SEQ ID NO:114.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:113.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 25 (aa) SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113; and
    - (ab) the nucleotide sequence of the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 30 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113; and

(bb) the nucleotide sequence of the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

15 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:113 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113. Also preferably the  
20 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113 from nucleotide 202 to nucleotide 429, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:113 from nucleotide 202 to nucleotide 429, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:113  
25 from nucleotide 202 to nucleotide 429. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113 from nucleotide 292 to nucleotide 429, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:113 from nucleotide 292 to nucleotide 429, to a nucleotide sequence  
30 corresponding to the 3' end of said sequence of SEQ ID NO:113 from nucleotide 292 to nucleotide 429.



In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:114;
  - 5 (b) a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins. Preferably such
- 10 protein comprises the amino acid sequence of SEQ ID NO:114. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:114, or a protein comprising a fragment of the amino acid sequence of SEQ
- 15 ID NO:114 having biological activity, the fragment comprising the amino acid sequence from amino acid 33 to amino acid 42 of SEQ ID NO:114.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:115;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115 from nucleotide 37 to nucleotide 1113;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115 from nucleotide 88 to nucleotide 1113;
- 25 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vq24\_1 deposited with the ATCC under accession number
- 30 PTA-1075;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:116;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:116;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:115.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:115 from nucleotide 37 to nucleotide 1113; the nucleotide sequence of SEQ ID NO:115 from nucleotide 88 to nucleotide 1113; the nucleotide sequence of the full-length protein coding sequence of clone vq24\_1 deposited with the ATCC under accession number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vq24\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq24\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID

NO:116, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising the amino acid sequence from amino acid 174 to amino acid 183 of SEQ ID NO:116.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
5 ID NO:115.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
    - (i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
      - (aa) SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115; and
      - (ab) the nucleotide sequence of the cDNA insert of clone  
15 vq24\_1 deposited with the ATCC under accession number PTA-1075;
    - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
    - (iii) isolating the DNA polynucleotides detected with the  
20 probe(s);
- and
- (b) a process comprising the steps of:
    - (i) preparing one or more polynucleotide primers that hybridize  
25 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
      - (ba) SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115; and
      - (bb) the nucleotide sequence of the cDNA insert of clone  
30 vq24\_1 deposited with the ATCC under accession number PTA-1075;
    - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:115, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:115 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:115 from nucleotide 37 to nucleotide  
10 1113, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:115 from nucleotide 37 to nucleotide 1113, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:115 from nucleotide 37 to nucleotide 1113. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:115 from nucleotide 88 to nucleotide 1113, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:115 from nucleotide 88 to nucleotide 1113, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:115 from nucleotide 88 to nucleotide 1113.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:116;
- (b) a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:116. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino  
30 acid sequence of SEQ ID NO:116 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:116, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:116 having biological activity, the fragment comprising the amino acid sequence from amino acid 174 to amino acid 183 of SEQ ID NO:116.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5                   (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:117;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:117 from nucleotide 40 to nucleotide 207;
- (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID  
10               NO:117 from nucleotide 103 to nucleotide 207;
- (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (e)     a polynucleotide encoding the full-length protein encoded by the  
15               cDNA insert of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (f)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- 20               (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (h)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:118;
- 25               (i)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:118;
- (j)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 30               (k)     a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:117.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:117 from nucleotide 40 to nucleotide 207; the nucleotide sequence of SEQ ID NO:117 from nucleotide 103 to nucleotide 207; the nucleotide sequence of the full-length protein coding sequence of clone vq26\_1 deposited with the ATCC under accession number PTA-1075; or the nucleotide sequence of a mature protein coding sequence of clone vq26\_1 deposited with the ATCC under accession number PTA-1075. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone vq26\_1 deposited with the ATCC under accession number PTA-1075. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:118, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising the amino acid sequence from amino acid 23 to amino acid 32 of SEQ ID NO:118.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:117.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:117, but excluding the poly(A) tail at the 3' end of SEQ ID NO:117; and

- (ab) the nucleotide sequence of the cDNA insert of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize
- 10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:117, but excluding the poly(A) tail at the 3' end of SEQ ID NO:117; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 20 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:117, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:117 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:117, but

25 excluding the poly(A) tail at the 3' end of SEQ ID NO:117. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:117 from nucleotide 40 to nucleotide 207, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:117 from nucleotide 40 to nucleotide 207, to a nucleotide

30 sequence corresponding to the 3' end of said sequence of SEQ ID NO:117 from nucleotide 40 to nucleotide 207. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

NO:117 from nucleotide 103 to nucleotide 207, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:117 from nucleotide 103 to nucleotide 207, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:117 from nucleotide 103 to nucleotide 207.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:118;
- (b) a fragment of the amino acid sequence of SEQ ID NO:118, the  
10 fragment comprising eight contiguous amino acids of SEQ ID NO:118; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
vq26\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:118. In further preferred  
15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:118, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising the amino acid sequence  
20 from amino acid 23 to amino acid 32 of SEQ ID NO:118.

In certain preferred embodiments, the polynucleotide is operably linked to an expression control sequence. The invention also provides a host cell, including bacterial, yeast, insect and mammalian cells, transformed with such polynucleotide compositions.  
25 Also provided by the present invention are organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein.

Processes are also provided for producing a protein, which comprise:

- (a) growing a culture of the host cell transformed with such  
30 polynucleotide compositions in a suitable culture medium; and
- (b) purifying the protein from the culture.

The protein produced according to such methods is also provided by the present invention.



Protein compositions of the present invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody which specifically reacts with such protein are also provided by the present invention.

Methods are also provided for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition comprising a protein of the present invention and a pharmaceutically acceptable carrier.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B are schematic representations of the pED6 and pNOTs vectors, respectively, used for deposit of clones disclosed herein.

#### DETAILED DESCRIPTION

##### ISOLATED PROTEINS AND POLYNUCLEOTIDES

Nucleotide and amino acid sequences, as presently determined, are reported below for each clone and protein disclosed in the present application. The nucleotide sequence of each clone can readily be determined by sequencing of the deposited clone in accordance with known methods. The predicted amino acid sequence (both full-length and mature forms) can then be determined from such nucleotide sequence. The amino acid sequence of the protein encoded by a particular clone can also be determined by expression of the clone in a suitable host cell, collecting the protein and determining its sequence. For each disclosed protein applicants have identified what they have determined to be the reading frame best identifiable with sequence information available at the time of filing.

As used herein a "secreted" protein is one which, when expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported across the membrane of the endoplasmic reticulum.

Clone "vc62\_1"

A polynucleotide of the present invention has been identified as clone "vc62\_1". vc62\_1 was isolated from a human fetal brain cDNA library and was identified as  
5 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc62\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc62\_1 protein").

The nucleotide sequence of vc62\_1 as presently determined is reported in SEQ ID NO:1, and includes a poly(A) tail. What applicants presently believe to be the proper  
10 reading frame and the predicted amino acid sequence of the vc62\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:2. Amino acids 3 to 15 of SEQ ID NO:2 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted  
15 leader/signal sequence not be separated from the remainder of the vc62\_1 protein. If the 'G' residue at position 254 of SEQ ID NO:1 were deleted, another potential vc62\_1 reading frame and predicted amino acid sequence that would then be encoded by nucleotides 27 to 365 of SEQ ID NO:1 is reported in SEQ ID NO:169.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20 vc62\_1 should be approximately 4221 bp.

The nucleotide sequence disclosed herein for vc62\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc62\_1 demonstrated at least some similarity with sequences identified as AA580489 (nn22a10.s1 NCI\_CGAP\_Co12 Homo sapiens cDNA clone  
25 IMAGE 1084602, mRNA sequence), AF047042 (Homo sapiens citrate synthase mRNA, complete cds), and T04200 (Sugar beet citrate synthase cDNA; standard; cDNA to mRNA). The predicted amino acid sequence disclosed herein for vc62\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc62\_1 protein demonstrated at least some similarity to  
30 sequences identified as AF047042 (citrate synthase [Homo sapiens]) and R82839 (Sugar beet citrate synthase). Based upon sequence similarity, vc62\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vp10\_1"

A polynucleotide of the present invention has been identified as clone "vp10\_1". vp10\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp10\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp10\_1 protein").

The nucleotide sequence of vp10\_1 as presently determined is reported in SEQ ID NO:3, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp10\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:4. Amino acids 19 to 31 of SEQ ID NO:4 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 32. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp10\_1 protein. If another 'G' residue were inserted in SEQ ID NO:3 after the 'G' residue at position 868, another potential vp10\_1 reading frame and predicted amino acid sequence that would be encoded by what would then be nucleotides 6 to 968 of SEQ ID NO:3 is reported in SEQ ID NO:170.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp10\_1 should be approximately 1401 bp.

The nucleotide sequence disclosed herein for vp10\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp10\_1 demonstrated at least some similarity with sequences identified as AA733074 (zg79d07.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 399565 3' similar to WP:C15H9.5 CE06834; mRNA sequence). The predicted amino acid sequence disclosed herein for vp10\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vp10\_1 protein demonstrated at least some similarity to the sequence identified as U56965 (unknown protein [Caenorhabditis elegans]). Based upon sequence similarity, vp10\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vp10\_1 protein sequence centered around amino acid 270 of SEQ ID NO:4.

Clone "vp11\_1"

A polynucleotide of the present invention has been identified as clone "vp11\_1". vp11\_1 was isolated from a human adult prostate cDNA library and was identified as  
5 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp11\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp11\_1 protein").

The nucleotide sequence of vp11\_1 as presently determined is reported in SEQ ID NO:5, and includes a poly(A) tail. What applicants presently believe to be the proper  
10 reading frame and the predicted amino acid sequence of the vp11\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:6. Amino acids 5 to 17 of SEQ ID NO:6 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted  
15 leader/signal sequence not be separated from the remainder of the vp11\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp11\_1 should be approximately 1329 bp.

The nucleotide sequence disclosed herein for vp11\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
20 FASTA search protocols. No hits were found in the database.

Clone "vp13\_1"

A polynucleotide of the present invention has been identified as clone "vp13\_1". vp13\_1 was isolated from a human adult prostate cDNA library and was identified as  
25 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp13\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp13\_1 protein").

The nucleotide sequence of vp13\_1 as presently determined is reported in SEQ ID NO:7, and includes a poly(A) tail. What applicants presently believe to be the proper  
30 reading frame and the predicted amino acid sequence of the vp13\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:8. Amino acids 13 to 25 of SEQ ID NO:8 are a predicted leader/signal sequence, with the predicted mature amino

acid sequence beginning at amino acid 26. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp13\_1 protein.

Other potential vp13\_1 reading frames and predicted amino acid sequences are  
5 encoded by nucleotides 151 to 267 of SEQ ID NO:7, with the encoded amino acid  
sequence reported in SEQ ID NO:171, and by nucleotides 209 to 787 of SEQ ID NO:7,  
with the encoded amino acid sequence reported in SEQ ID NO:172. Amino acids 1 to 13  
of SEQ ID NO:172 are a predicted leader/signal sequence, with the predicted mature  
amino acid sequence beginning at amino acid 14. Due to the hydrophobic nature of this  
10 predicted leader/signal sequence, it is likely to act as a transmembrane domain should it  
not be separated from the remainder of the protein of SEQ ID NO:172. The protein of  
SEQ ID NO:172 also demonstrates significant homology to the human Notch protein,  
Delta proteins from various species, and other EGF-repeat-containing transmembrane  
proteins. A deletion or insertion causing a frame-shift in the nucleotide sequence of SEQ  
15 ID NO:7 in the region approximately between nucleotides 208 and 267 of SEQ ID NO:7  
could join the reading frames of SEQ ID NO:171 and SEQ ID NO:172 into a single  
reading frame encoding an EGF-repeat-containing protein. Further, the region  
approximately between nucleotides 605 and 850 may be an alternatively spliced exon.

If the 'A' residue at position 423 of SEQ ID NO:7 were deleted, another potential  
20 vp13\_1 reading frame and predicted amino acid sequence that would be encoded by what  
would then be nucleotides 288 to 503 of SEQ ID NO:7 is reported in SEQ ID NO:173.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
vp13\_1 should be approximately 1048 bp.

The nucleotide sequence disclosed herein for vp13\_1 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
FASTA search protocols. vp13\_1 demonstrated at least some similarity with sequences  
identified as AA190865 (zp85b02.s1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA  
clone 626955 3' similar to TR G1336628 G1336628 EGF REPEAT TRANSMEMBRANE  
PROTEIN; mRNA sequence), and U57368 (Mus musculus EGF repeat transmembrane  
30 protein mRNA, complete cds). The predicted amino acid sequence disclosed herein for  
vp13\_1 was searched against the GenPept and GeneSeq amino acid sequence databases  
using the BLASTX search protocol. The predicted vp13\_1 protein demonstrated at least

some similarity to sequences identified as AC004663 (Notch 3 [Homo sapiens]), R28960 (Delta D11), and U57368 (EGF repeat transmembrane protein [Mus musculus]). Based upon sequence similarity, vp13\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential  
5 transmembrane domain within the vp13\_1 protein sequence centered around amino acid 56 of SEQ ID NO:8.

Clone "vp16\_1"

A polynucleotide of the present invention has been identified as clone "vp16\_1".  
10 vp16\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp16\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp16\_1 protein").

The nucleotide sequence of vp16\_1 as presently determined is reported in SEQ ID  
15 NO:9, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp16\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:10. Amino acids 34 to 46 of SEQ ID NO:10 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 47. Due to the hydrophobic nature of the predicted  
20 leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp16\_1 protein. Another potential vp16\_1 reading frame and predicted amino acid sequence is encoded by basepairs 1621 to 1839 of SEQ ID NO:9 and is reported in SEQ ID NO:174.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
25 vp16\_1 should be approximately 2105 bp.

The nucleotide sequence disclosed herein for vp16\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp16\_1 demonstrated at least some similarity with sequences identified as AA523851 (ng31e01.s1 NCI\_CGAP\_Co3 Homo sapiens cDNA clone  
30 IMAGE:936408, mRNA sequence). Based upon sequence similarity, vp16\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the vp16\_1 protein

sequence, one centered around amino acid 36 and another around amino acid 69 of SEQ ID NO:10. The nucleotide sequence of vp16\_1 indicates that it may contain an Alu repetitive element.

5        Clone "vp21\_1"

A polynucleotide of the present invention has been identified as clone "vp21\_1". vp21\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp21\_1 is a full-length clone, including the  
10    entire coding sequence of a secreted protein (also referred to herein as "vp21\_1 protein").

The nucleotide sequence of vp21\_1 as presently determined is reported in SEQ ID NO:11, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp21\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:12. Amino acids 62 to 74  
15    of SEQ ID NO:12 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 75. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp21\_1 protein. Another potential vp21\_1 reading frame and predicted amino acid sequence encoded by  
20    basepairs 598 to 831 of SEQ ID NO:11 is reported in SEQ ID NO:175. Amino acids 1 to 6 of SEQ ID NO:175 and amino acids 41 to 43 of SEQ ID NO:175 are predicted leader/signal sequences, with the predicted mature amino acid sequences beginning at amino acid 7 or at amino acid 44, respectively.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
25    vp21\_1 should be approximately 1538 bp.

The nucleotide sequence disclosed herein for vp21\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp21\_1 demonstrated at least some similarity with sequences identified as AC004076 (Homo sapiens chromosome 19, cosmid R30217, complete  
30    sequence). The predicted amino acid sequence disclosed herein for vp21\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vp21\_1 protein demonstrated at least some similarity to

sequences identified as AC003682 (Zinc finger protein F18547\_1 [Homo sapiens]) and W19106 (Tat pheromone receptor VN5). Based upon sequence similarity, vp21\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts potential transmembrane domains within the predicted vp21\_1 protein sequences, one centered around amino acid 70 of SEQ ID NO:12, and one centered around amino acid 17 of SEQ ID NO:175.

#### Clone "vp22\_1"

A polynucleotide of the present invention has been identified as clone "vp22\_1". vp22\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp22\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vp22\_1 protein").

The nucleotide sequence of vp22\_1 as presently determined is reported in SEQ ID NO:13, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp22\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:14. Amino acids 13 to 25 of SEQ ID NO:14 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 26. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp22\_1 protein.

Another potential vp22\_1 reading frame and predicted amino acid sequence encoded by basepairs 408 to 1154 of SEQ ID NO:13 is reported in SEQ ID NO:176. Amino acids 40 to 52 of SEQ ID NO:176 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 53. Due to the hydrophobic nature of this predicted leader/signal sequence, it is likely to act as a transmembrane domain should it not be separated from the remainder of the protein of SEQ ID NO:176. A frameshift within the nucleotide sequence of SEQ ID NO:13 approximately between nucleotides 163 and 477 could join the openreading frames of SEQ ID NO:14 and SEQ ID NO:176.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp22\_1 should be approximately 1718 bp.



The nucleotide sequence disclosed herein for vp22\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp22\_1 demonstrated at least some similarity with sequences identified as AA526186 (ni94h03.s1 NCI\_CGAP\_Pr21 Homo sapiens cDNA clone  
5 IMAGE:984533, mRNA sequence), AA570505 (nk64h01.s1 NCI\_CGAP\_Sch1 Homo sapiens cDNA clone IMAGE 1018321, mRNA sequence), AB006085 (Danio rerio mRNA for MINDIN2, complete cds), and T78360 (Human neuronal attachment factor-1 DNA; standard; DNA). The predicted amino acid sequence disclosed herein for vp22\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the  
10 BLASTX search protocol. The predicted vp22\_1 protein demonstrated at least some similarity to sequences identified as AB006085 (MINDIN2 [Danio rerio]) and W23663 (Human neuronal attachment factor-1). Based upon sequence similarity, vp22\_1 proteins and each similar protein or peptide may share at least some activity.

15        Clone "vq2\_1"

A polynucleotide of the present invention has been identified as clone "vq2\_1". vq2\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq2\_1 is a full-length clone, including the entire coding  
20 sequence of a secreted protein (also referred to herein as "vq2\_1 protein").

The nucleotide sequence of vq2\_1 as presently determined is reported in SEQ ID NO:15, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq2\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:16. Amino acids 4 to 16  
25 of SEQ ID NO:16 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq2\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
30 vq2\_1 should be approximately 896 bp.

The nucleotide sequence disclosed herein for vq2\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and

FASTA search protocols. vq2\_1 demonstrated at least some similarity with sequences identified as AI203981 (qe76h05.x1 Soares\_fetal\_lung\_NbHL19W Homo sapiens cDNA clone IMAGE:1744953 3', mRNA sequence) and T97082 (Human haematopoietic-specific protein (HSP) DNA; standard; DNA). The predicted amino acid sequence disclosed herein  
5 for vq2\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vq2\_1 protein demonstrated at least some similarity to the sequence identified as W35904 (Human haematopoietic-specific protein (HSP)). Based upon sequence similarity, vq2\_1 proteins and each similar protein or peptide may share at least some activity.

10

#### Clone "vq3\_1"

A polynucleotide of the present invention has been identified as clone "vq3\_1". vq3\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid  
15 sequence of the encoded protein. vq3\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq3\_1 protein").

The nucleotide sequence of vq3\_1 as presently determined is reported in SEQ ID NO:17, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq3\_1 protein corresponding  
20 to the foregoing nucleotide sequence is reported in SEQ ID NO:18. Amino acids 11 to 23 of SEQ ID NO:18 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq3\_1 protein.

25 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq3\_1 should be approximately 1490 bp.

The nucleotide sequence disclosed herein for vq3\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No significant hits were found in the database. The nucleotide  
30 sequence of vq3\_1 indicates that it may contain an Alu repetitive element.

Clone "vq5\_1"

A polynucleotide of the present invention has been identified as clone "vq5\_1". vq5\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid  
5 sequence of the encoded protein. vq5\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq5\_1 protein").

The nucleotide sequence of vq5\_1 as presently determined is reported in SEQ ID NO:19, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq5\_1 protein corresponding  
10 to the foregoing nucleotide sequence is reported in SEQ ID NO:20. Amino acids 9 to 21 of SEQ ID NO:20 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq5\_1 protein.

15 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq5\_1 should be approximately 2207 bp.

The nucleotide sequence disclosed herein for vq5\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq5\_1 demonstrated at least some similarity with sequences  
20 identified as AQ036276 (CIT-HSP-2331M15.TF CIT-HSP Homo sapiens genomic clone 2331M15, genomic survey sequence) and T24918 (Human gene signature HUMGS07027; standard; cDNA to mRNAP). Based upon sequence similarity, vq5\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts that the signal sequence at residue 22 of SEQ ID NO:20 is also a  
25 potential transmembrane domain.

Clone "vq6\_1"

A polynucleotide of the present invention has been identified as clone "vq6\_1". vq6\_1 was isolated from a human adult lung cDNA library and was identified as encoding  
30 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq6\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq6\_1 protein").

The nucleotide sequence of vq6\_1 as presently determined is reported in SEQ ID NO:21, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq6\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:22. Amino acids 6 to 18 of SEQ ID NO:22 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 19. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq6\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq6\_1 should be approximately 1875 bp.

The nucleotide sequence disclosed herein for vq6\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq6\_1 demonstrated at least some similarity with sequences identified as AA729043 (nw22d09.s1 NCI\_CGAP\_GCB0 Homo sapiens cDNA clone IMAGE:1241201 similar to contains Alu repetitive element; mRNA sequence). Based upon sequence similarity, vq6\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the vq6\_1 protein sequence centered around amino acid 37 of SEQ ID NO:22. The nucleotide sequence of vq6\_1 indicates that it may contain an Alu repetitive element.

#### Clone "vr1\_1"

A polynucleotide of the present invention has been identified as clone "vr1\_1". vr1\_1 was isolated from a human adult muscle cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vr1\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vr1\_1 protein").

The nucleotide sequence of vr1\_1 as presently determined is reported in SEQ ID NO:23, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vr1\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:24. Amino acids 34 to 46 of SEQ ID NO:24 are a predicted leader/signal sequence, with the predicted mature amino

acid sequence beginning at amino acid 47. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vr1\_1 protein. The region of SEQ ID NO:23 approximately between nucleotides 1931 and 1977 of SEQ ID NO:23 may be an alternatively spliced exon.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vr1\_1 should be approximately 1512 bp.

The nucleotide sequence disclosed herein for vr1\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vr1\_1 demonstrated at least some similarity with sequences identified as AL031602 (Human DNA sequence \*\*\* SEQUENCING IN PROGRESS \*\*\* from clone 1174N9; HTGS phase 1), I64695 (Sequence 1 from patent US 5665588), and T35233 (Natural killer lytic associated protein cDNA; standard; cDNA). The predicted amino acid sequence disclosed herein for vr1\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vr1\_1 protein demonstrated at least some similarity to sequences identified as R99256 (Natural killer lytic associated protein), and X71642 (GEG-154 gene product [Mus musculus]). Based upon sequence similarity, vr1\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the vr1\_1 protein sequence centered around amino acid 150 of SEQ ID NO:24.

#### Clone "vc63\_1"

A polynucleotide of the present invention has been identified as clone "vc63\_1". vc63\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc63\_1 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "vc63\_1 protein").

The nucleotide sequence of vc63\_1 as presently determined is reported in SEQ ID NO:25, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc63\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:26. Another potential

vc63\_1 reading frame and predicted amino acid sequence encoded by basepairs 528 to 1100 of SEQ ID NO:25 is reported in SEQ ID NO:177. Amino acids 140 to 152 of SEQ ID NO:177 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 153. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:177.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc63\_1 should be approximately 2397 bp.

10 The nucleotide sequence disclosed herein for vc63\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc63\_1 demonstrated at least some similarity with sequences identified as N66555 (yy69b07.s1 Homo sapiens cDNA clone 278773 3') and T21367 (Human gene signature HUMGS02731; standard; cDNA to mRNA). The predicted amino acid sequence disclosed herein for vc63\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc63\_1 protein demonstrated at least some similarity to the sequence identified as Z36948 (D2089.2 [Caenorhabditis elegans]). Based upon sequence similarity, vc63\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the protein sequence of SEQ ID NO:177, centered around amino acid 153 of SEQ ID NO:177.

#### Clone "vb25\_1"

A polynucleotide of the present invention has been identified as clone "vb25\_1". vb25\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb25\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb25\_1 protein").

The nucleotide sequence of vb25\_1 as presently determined is reported in SEQ ID NO:27, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb25\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:28. Amino acids 5 to 17

of SEQ ID NO:28 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb25\_1 protein.

5           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb25\_1 should be approximately 1677 bp.

          The nucleotide sequence disclosed herein for vb25\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb25\_1 demonstrated at least some similarity with sequences  
10 identified as Z73429 (Human DNA sequence from cosmid cN32F9 on chromosome 22q11.2-qter Contains CpG island). Based upon sequence similarity, vb25\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vb25\_1 indicates that it may contain one or more of the following repetitive elements: AC simple repeat, AG simple repeat, ALU, MIR.

15

Clone "vb27\_1"

          A polynucleotide of the present invention has been identified as clone "vb27\_1". vb27\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the  
20 amino acid sequence of the encoded protein. vb27\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb27\_1 protein").

          The nucleotide sequence of vb27\_1 as presently determined is reported in SEQ ID NO:29, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb27\_1 protein corresponding  
25 to the foregoing nucleotide sequence is reported in SEQ ID NO:30. Amino acids 14 to 26 of SEQ ID NO:30 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb27\_1 protein.

30           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb27\_1 should be approximately 3456 bp.

The nucleotide sequence disclosed herein for vb27\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb27\_1 demonstrated at least some similarity with sequences identified as AC005035 (Homo sapiens BAC clone NH0353P23 from 2, complete  
5 sequence) and H73579 (yu29f09.r1 Homo sapiens cDNA clone 235241 5'). Based upon sequence similarity, vb27\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vb27\_1 indicates that it may contain one or more of the following repetitive elements: ALU, Mer3.

10        Clone "vb28\_1"

A polynucleotide of the present invention has been identified as clone "vb28\_1". vb28\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb28\_1 is a full-length clone, including the  
15 entire coding sequence of a secreted protein (also referred to herein as "vb28\_1 protein").

The nucleotide sequence of vb28\_1 as presently determined is reported in SEQ ID NO:31, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb28\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:32. Amino acids 4 to 16  
20 of SEQ ID NO:32 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb28\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
25 vb28\_1 should be approximately 3008 bp.

The nucleotide sequence disclosed herein for vb28\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb28\_1 demonstrated at least some similarity with sequences identified as AA046671 (zf12d09.r1 Soares\_fetal\_heart\_NbHH19W Homo sapiens cDNA  
30 clone IMAGE:376721 5' similar to PIR:A38745 A38745 cell adhesion molecule CD44 precursor - rat; mRNA sequence) and V22687 (DNA encoding a CD44-like protein). The predicted amino acid sequence disclosed herein for vb28\_1 was searched against the



GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vb28\_1 protein demonstrated at least some similarity to sequences identified as W56249 (Amino acid sequence of a CD44-like protein) and X66081 (CD44 [Mus musculus]). Based upon sequence similarity, vb28\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "vb29\_1"

A polynucleotide of the present invention has been identified as clone "vb29\_1". vb29\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb29\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb29\_1 protein").

The nucleotide sequence of vb29\_1 as presently determined is reported in SEQ ID NO:33, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb29\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:34. Amino acids 11 to 23 of SEQ ID NO:34 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb29\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb29\_1 should be approximately 2970 bp.

The nucleotide sequence disclosed herein for vb29\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb29\_1 demonstrated at least some similarity with sequences identified as AA084068 (zn16d12.r1 Stratagene neuroepithelium NT2RAMI 937234 Homo sapiens cDNA clone 547607 5', mRNA sequence) and AQ418918 (RPCI-11-185K12.TV RPCI-11 Homo sapiens genomic clone RPCI-11-185K12, genomic survey sequence). Based upon sequence similarity, vb29\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vb29\_1 protein sequence centered

around amino acid 41 of SEQ ID NO:34. The nucleotide sequence of vb29\_1 indicates that it may contain an Alu repetitive element.

Clone "vb30\_1"

5           A polynucleotide of the present invention has been identified as clone "vb30\_1". vb30\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb30\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb30\_1 protein").

10           The nucleotide sequence of vb30\_1 as presently determined is reported in SEQ ID NO:35, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb30\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:36. Amino acids 15 to 27 of SEQ ID NO:36 are a predicted leader/signal sequence, with the predicted mature amino  
15           acid sequence beginning at amino acid 28. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb30\_1 protein.

          The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb30\_1 should be approximately 3325 bp.

20           The nucleotide sequence disclosed herein for vb30\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No significant hits were found in the databases. The nucleotide sequence of vb30\_1 indicates that it may contain an Alu repetitive element.

25           Clone "vc67\_1"

          A polynucleotide of the present invention has been identified as clone "vc67\_1". vc67\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc67\_1 is a full-length clone, including the  
30           entire coding sequence of a secreted protein (also referred to herein as "vc67\_1 protein").

          The nucleotide sequence of vc67\_1 as presently determined is reported in SEQ ID NO:37, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the vc67\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:38. Another potential vc67\_1 reading frame and predicted amino acid sequence encoded by basepairs 3 to 242 of SEQ ID NO:37 is reported in SEQ ID NO:178.

- 5           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc67\_1 should be approximately 2305 bp.

          The nucleotide sequence disclosed herein for vc67\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc67\_1 demonstrated at least some similarity with sequences  
10 identified as T23222 (Human gene signature HUMGS05018), W87297 (zh67h03.s1 Soares\_fetal\_liver\_spleen\_1NFLS\_S1 Homo sapiens cDNA clone IMAGE 417173 3', mRNA sequence), and Z97201 (Human DNA sequence \*\*\* SEQUENCING IN PROGRESS \*\*\* from clone 94M16, WORKING DRAFT SEQUENCE). The predicted amino acid sequence disclosed herein for vc67\_1 was searched against the GenPept and  
15 GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc67\_1 protein demonstrated at least some similarity to sequences identified as W69427 (Human secreted protein bk291\_3) and Z68751 (Similarity to Yeast hypothetical protein YKK0 (SW YKK0\_YEAST); cDNA EST EMBL C12578 comes from this gene; cDNA EST yk329g12.5 comes from this gene; cDNA EST yk415). Based upon sequence  
20 similarity, vc67\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the vc67\_1 protein sequence of SEQ ID NO:38, one centered around amino acid 58 and another around amino acid 85 of SEQ ID NO:38.

25           Clone "vf4\_1"

          A polynucleotide of the present invention has been identified as clone "vf4\_1". vf4\_1 was isolated from a human adult heart cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vf4\_1 is a full-length clone, including the entire coding  
30 sequence of a secreted protein (also referred to herein as "vf4\_1 protein").

          The nucleotide sequence of vf4\_1 as presently determined is reported in SEQ ID NO:39, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the vf4\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:40. Amino acids 5 to 17 of SEQ ID NO:40 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vf4\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vf4\_1 should be approximately 972 bp.

The nucleotide sequence disclosed herein for vf4\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vf4\_1 demonstrated at least some similarity with sequences identified as AA813690 (ai71a09.s1 Soares\_testis\_NHT Homo sapiens cDNA clone 1376248 3', mRNA sequence) and V86544 (EST clone AZ285). Based upon sequence similarity, vf4\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "vg3\_1"

A polynucleotide of the present invention has been identified as clone "vg3\_1". vg3\_1 was isolated from a human adult brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vg3\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vg3\_1 protein").

The nucleotide sequence of vg3\_1 as presently determined is reported in SEQ ID NO:41, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vg3\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:42. Amino acids 13 to 25 of SEQ ID NO:42 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 26. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vg3\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vg3\_1 should be approximately 3667 bp.

The nucleotide sequence disclosed herein for vg3\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vg3\_1 demonstrated at least some similarity with sequences identified as AI283122 (qm51h10.x1 Soares\_placenta\_8to9weeks\_2NbHP8to9W Homo sapiens cDNA clone IMAGE 1892323 3', mRNA sequence). The predicted amino acid sequence disclosed herein for vg3\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vg3\_1 protein demonstrated at least some similarity to sequences identified as U53155 (ZC513.5 [Caenorhabditis elegans]). Based upon sequence similarity, vg3\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts the following transmembrane domains within the vg3\_1 protein sequence: four certain transmembrane domains centered around amino acids 78, 133, 156, and 298 of SEQ ID NO:42, respectively; four strongly putative transmembrane domains centered around amino acids 105, 189, 221, and 354 of SEQ ID NO:42, respectively; and six possible transmembrane domains centered around amino acids 262, 272, 322, 367, 432, and 460 of SEQ ID NO:42, respectively. Motifs analysis detected a Crystallins beta and gamma 'Greek key' motif signature around amino acid 52 of SEQ ID NO:42. The nucleotide sequence of vg3\_1 indicates that it may contain an Alu repetitive element.

#### 20        Clone "vo2\_1"

A polynucleotide of the present invention has been identified as clone "vo2\_1". vo2\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo2\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo2\_1 protein").

The nucleotide sequence of vo2\_1 as presently determined is reported in SEQ ID NO:43, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo2\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:44.

30        Another potential vo2\_1 reading frame and predicted amino acid sequence encoded by basepairs 95 to 280 of SEQ ID NO:43 is reported in SEQ ID NO:179. Amino acids 9 to 21 of SEQ ID NO:179 are a predicted leader/signal sequence, with the predicted mature

amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:179.

5           Another potential vo2\_1 reading frame and predicted amino acid sequence encoded by basepairs 76 to 258 of SEQ ID NO:43 is reported in SEQ ID NO:180. Amino acids 18 to 30 of SEQ ID NO:180 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 31. Due to the hydrophobic nature of this predicted leader/signal sequence, it is likely to act as a transmembrane domain should it  
10   not be separated from the remainder of the protein of SEQ ID NO:180.

          Another potential vo2\_1 reading frame and predicted amino acid sequence encoded by basepairs 2131 to 2310 of SEQ ID NO:43 is reported in SEQ ID NO:181. Amino acids 38 to 50 of SEQ ID NO:181 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 11; amino acids 19 to 31 of SEQ ID  
15   NO:181 are also a possible leader/signal sequence, with the predicted mature amino acid sequence in this case beginning at amino acid 32. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the protein of SEQ ID NO:181.

          The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20   vo2\_1 should be approximately 2903 bp.

          The nucleotide sequence disclosed herein for vo2\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo2\_1 demonstrated at least some similarity with sequences identified as AI094627 (oy61b07.s1 NCI\_CGAP\_Brn23 Homo sapiens cDNA clone  
25   IMAGE 1670293 3', mRNA sequence). Based upon sequence similarity, vo2\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "vo3\_1"

          A polynucleotide of the present invention has been identified as clone "vo3\_1".  
30   vo3\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the

amino acid sequence of the encoded protein. vo3\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo3\_1 protein").

The nucleotide sequence of vo3\_1 as presently determined is reported in SEQ ID NO:45, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the vo3\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:46. Amino acids 107 to 119 of SEQ ID NO:46 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 120. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the  
10 predicted leader/signal sequence not be separated from the remainder of the vo3\_1 protein.

If a "C" residue were to be deleted from the nucleotide sequence of SEQ ID NO:45 at either position 917 or position 918, another potential vo3\_1 reading frame and predicted amino acid sequence encoded by what would then be basepairs 697 to 1377 of SEQ ID NO:45 is reported in SEQ ID NO:182. Amino acids 62 to 74 of SEQ ID NO:182 are a  
15 predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 75. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:182.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20 vo3\_1 should be approximately 1592 bp.

The nucleotide sequence disclosed herein for vo3\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo3\_1 demonstrated at least some similarity with sequences identified as AA530997 (nj07a06.s1 NCI\_CGAP\_Pr22 Homo sapiens cDNA clone  
25 IMAGE:985618 3', mRNA sequence), AA683481 (zl55b03.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone IMAGE:505805 3', mRNA sequence), D88158 (Pig mRNA for cytochrome b561, complete cds), and V84516 (Human secreted protein gene 106 clone HTOEY16). The predicted amino acid sequence disclosed herein for vo3\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the  
30 BLASTX search protocol. The predicted vo3\_1 protein demonstrated at least some similarity to sequences identified as U06715 (HCYTO B561 [Homo sapiens]) and W89024 (Polypeptide fragment encoded by gene 156). Based upon sequence similarity, vo3\_1

proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts five potential transmembrane domains within the vo3\_1 protein sequence, centered around amino acids 35, 75, 113, 146, and 191 of SEQ ID NO:46, respectively.

5

Clone "vo5\_1"

A polynucleotide of the present invention has been identified as clone "vo5\_1". vo5\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo5\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo5\_1 protein").

The nucleotide sequence of vo5\_1 as presently determined is reported in SEQ ID NO:47, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo5\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:48. Amino acids 8 to 20 of SEQ ID NO:48 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vo5\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo5\_1 should be approximately 2487 bp.

The nucleotide sequence disclosed herein for vo5\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo5\_1 demonstrated at least some similarity with sequences identified as AA868551 (ak43f09.s1 Soares testis NHT Homo sapiens cDNA clone IMAGE:1408745 3', mRNA sequence) and AC005500 (complete sequence [Homo sapiens Chromosome 22q11 PAC Clone p52f6 In DGCR Region]). Based upon sequence similarity, vo5\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vo5\_1 indicates that it may contain an Alu repetitive element.



Clone "vo6\_1"

A polynucleotide of the present invention has been identified as clone "vo6\_1". vo6\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo6\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo6\_1 protein").

The nucleotide sequence of vo6\_1 as presently determined is reported in SEQ ID NO:49, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo6\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:50. Amino acids 77 to 89 of SEQ ID NO:50 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 90. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vo6\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo6\_1 should be approximately 1272 bp.

The nucleotide sequence disclosed herein for vo6\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo6\_1 demonstrated at least some similarity with sequences identified as AL020989 (Human DNA sequence \*\*\* SEQUENCING IN PROGRESS \*\*\* from clone 192P9; HTGS phase 1), T34592 (NTII-11 nerve protein coding sequence), and U13617 (Rattus norvegicus Sprague-Dawley plasmolipin mRNA, complete cds). The predicted amino acid sequence disclosed herein for vo6\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo6\_1 protein demonstrated at least some similarity to sequences identified as R99799 (NTII-11 nerve protein, facilitates regeneration of nerve cells) and U13617 (plasmolipin [Rattus norvegicus]). Plasmolipin is an 18-kDa proteolipid protein found in kidney and brain, where it is restricted to the apical surface of tubular epithelial cells and to mammalian myelinated tracts, respectively; addition of plasmolipin to lipid bilayers induces the formation of ion channels, which are voltage-dependent and K(+)-selective. (See Fischer and Sapirstein, 1994, *J. Biol. Chem.* 269(40): 24912-24919, which is incorporated by reference herein). Based upon sequence similarity, vo6\_1 proteins and

each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three potential transmembrane domains within the vo6\_1 protein sequence, centered around amino acids 14, 42, and 90 of SEQ ID NO:50, respectively.

5           Clone "vo9\_1"

A polynucleotide of the present invention has been identified as clone "vo9\_1". vo9\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo9\_1 is a full-length clone, including the  
10   entire coding sequence of a secreted protein (also referred to herein as "vo9\_1 protein").

The nucleotide sequence of vo9\_1 as presently determined is reported in SEQ ID NO:51, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo9\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:52. Amino acids 22 to 34  
15   of SEQ ID NO: are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 35. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vo9\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20   vo9\_1 should be approximately 3331 bp.

The nucleotide sequence disclosed herein for vo9\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo9\_1 demonstrated at least some similarity with sequences identified as AA936961 (oo65f04.s1 NCI\_CGAP\_GC4 Homo sapiens cDNA clone  
25   IMAGE 1571071 3', mRNA sequence), AF010496 9Rhodobacter capsulatus strain SB1003, partial genome), AL035661 (Human DNA sequence \*\*\* SEQUENCING IN PROGRESS \*\*\* from clone 568C11, WORKING DRAFT SEQUENCE), and Q24673 (facA gene). The predicted amino acid sequence disclosed herein for vo9\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX  
30   search protocol. The predicted vo9\_1 protein demonstrated at least some similarity to sequences identified as R23968 (facA gene product) and Y15417 (acetate--CoA ligase

[Coprinus cinereus]). Based upon sequence similarity, vo9\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vo11\_1"

5           A polynucleotide of the present invention has been identified as clone "vo11\_1". vo11\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo11\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo11\_1 protein").

10           The nucleotide sequence of vo11\_1 as presently determined is reported in SEQ ID NO:53, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo11\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:54. Amino acids 52 to 64 of SEQ ID NO:54 are a predicted leader/signal sequence, with the predicted mature amino  
15   acid sequence beginning at amino acid 65.

          Another potential vo11\_1 reading frame and predicted amino acid sequence, encoded by basepairs 18 to 308 of SEQ ID NO:53, is reported in SEQ ID NO:183. Amino acids 10 to 22 of SEQ ID NO:183 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature  
20   of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:183.

          The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo11\_1 should be approximately 1509 bp.

25           The nucleotide sequence disclosed herein for vo11\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo11\_1 demonstrated at least some similarity with sequences identified as D83866 (similar to none, mRNA sequence). Based upon sequence similarity, vo11\_1 proteins and each similar protein or peptide may share at least some activity.

30

Clone "vo12\_1"

A polynucleotide of the present invention has been identified as clone "vo12\_1". vo12\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo12\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo12\_1 protein").

The nucleotide sequence of vo12\_1 as presently determined is reported in SEQ ID NO:55, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo12\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:56. Amino acids 4 to 16 of SEQ ID NO:56 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17.

Another potential vo12\_1 reading frame and predicted amino acid sequence, encoded by basepairs 107 to 310 of SEQ ID NO:55, is reported in SEQ ID NO:184. Amino acids 14 to 26 and amino acids 18 to 30 of SEQ ID NO:184 are predicted leader/signal sequences, with the predicted mature amino acid sequence beginning at amino acid 27 or at amino acid 31, respectively. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:184.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo12\_1 should be approximately 986 bp.

The nucleotide sequence disclosed herein for vo12\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo12\_1 demonstrated at least some similarity with sequences identified as AA444152 (zv51g06.r1 Soares testis NHT Homo sapiens cDNA clone 757210 5', mRNA sequence). Based upon sequence similarity, vo12\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vo12\_1 protein sequence centered around amino acid 51 of SEQ ID NO:56.

Clone "vo13\_1"

A polynucleotide of the present invention has been identified as clone "vo13\_1". vo13\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. The vo13\_1 clone includes coding sequence of a secreted protein (also referred to herein as "vo13\_1 protein").

The nucleotide sequence of vo13\_1 as presently determined is reported in SEQ ID NO:57, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo13\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:58. Amino acids 8 to 20 of SEQ ID NO:58 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo13\_1 should be approximately 1073 bp.

The nucleotide sequence disclosed herein for vo13\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo13\_1 demonstrated at least some similarity with sequences identified as AA988298 (os32a02.s1 NCI\_CGAP\_Br2 Homo sapiens cDNA clone IMAGE:1607018 3', mRNA sequence) and V69614 (Human secreted protein gene 4 clone HE8ND56). The predicted amino acid sequence disclosed herein for vo13\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo13\_1 protein demonstrated at least some similarity to sequences identified as W83934 (Human secreted protein from gene 4 clone HE8ND56). Based upon sequence similarity, vo13\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vo13\_1 protein sequence centered around amino acid 50 of SEQ ID NO:58.

Clone "vo14\_1"

A polynucleotide of the present invention has been identified as clone "vo14\_1". vo14\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the

amino acid sequence of the encoded protein. vo14\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo14\_1 protein").

The nucleotide sequence of vo14\_1 as presently determined is reported in SEQ ID NO:59, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the vo14\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:60. Amino acids 14 to 26 of SEQ ID NO:60 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
10 vo14\_1 should be approximately 1605 bp.

The nucleotide sequence disclosed herein for vo14\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No significant hits were found in the databases. Based upon sequence similarity, vo14\_1 proteins and each similar protein or peptide may share at least  
15 some activity. The nucleotide sequence of vo14\_1 indicates that it may contain one or more of the following repetitive elements: Alu, TAAAA repeat.

#### Clone "vo15\_1"

A polynucleotide of the present invention has been identified as clone "vo15\_1".  
20 vo15\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo15\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo15\_1 protein").

The nucleotide sequence of vo15\_1 as presently determined is reported in SEQ ID  
25 NO:61, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo15\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:62. Amino acids 13 to 25 of SEQ ID NO:62 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 26.

30 If a nucleotide were deleted between nucleotide 458 and nucleotide 460 of SEQ ID NO:61, another potential vo15\_1 reading frame and predicted amino acid sequence, encoded by what would then be basepairs 90 to 515 of SEQ ID NO:61, is reported in SEQ

ID NO:185. Amino acids 16 to 28 and amino acids 13 to 25 of SEQ ID NO:185 are predicted leader/signal sequences, with the predicted mature amino acid sequence beginning at amino acid 29 or at amino acid 26, respectively. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:185.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo15\_1 should be approximately 2842 bp.

The nucleotide sequence disclosed herein for vo15\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo15\_1 demonstrated at least some similarity with sequences identified as AI096756 (qb46e10.x1 NCI\_CGAP\_Brn23 Homo sapiens cDNA clone IMAGE 1703178 3', mRNA sequence). Based upon sequence similarity, vo15\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vo15\_1 protein sequence centered around amino acid 126 of SEQ ID NO:62. The nucleotide sequence of vo15\_1 indicates that it may contain one or more repeat sequences.

#### Clone "vo16\_1"

A polynucleotide of the present invention has been identified as clone "vo16\_1". vo16\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo16\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo16\_1 protein").

The nucleotide sequence of vo16\_1 as presently determined is reported in SEQ ID NO:63, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo16\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:64. Amino acids 51 to 63 of SEQ ID NO:64 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 64. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vo16\_1 protein.

If an "A" or "G" nucleotide were inserted between nucleotides 102 and 103 of SEQ ID NO:63 and an additional "A" residue inserted between nucleotides 271 and 273 of SEQ ID NO:63, another potential vo16\_1 reading frame and predicted amino acid sequence, encoded by what would then be basepairs 6 to 338 of SEQ ID NO:63, is reported in SEQ ID NO:186. Amino acids 5 to 17 and amino acids 4 to 16 of SEQ ID NO:186 are predicted leader/signal sequences, with the predicted mature amino acid sequence beginning at amino acid 18 or at amino acid 17, respectively. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:186.

Another potential vo16\_1 reading frame and predicted amino acid sequence, encoded by basepairs 846 to 1061 of SEQ ID NO:63, is reported in SEQ ID NO:187. Amino acids 12 to 24 and amino acids 11 to 23 of SEQ ID NO:187 are predicted leader/signal sequences, with the predicted mature amino acid sequence beginning at amino acid 25 or at amino acid 24, respectively. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:187.

Nucleotides 1 to 133 of SEQ ID NO:63 are nearly identical to nucleotides 862 to 994 of SEQ ID NO:63, resulting in amino acids 1 to 33 of SEQ ID NO:186 being identical to amino acids 8 to 40 of SEQ ID NO:187.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo16\_1 should be approximately 2113 bp.

The nucleotide sequence disclosed herein for vo16\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo16\_1 demonstrated at least some similarity with sequences identified as R79825 (y189a06.s1 Soares placenta Nb2HP Homo sapiens cDNA clone IMAGE:146386 3', mRNA sequence). Based upon sequence similarity, vo16\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vo16\_1 protein sequence centered around amino acid 64 of SEQ ID NO:64. The nucleotide sequence of vo16\_1 indicates that it may contain an Alu repeat region.



Clone "vo18\_1"

A polynucleotide of the present invention has been identified as clone "vo18\_1". vo18\_1 was isolated from a human adult pancreas cDNA library and was identified as  
5 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo18\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo18\_1 protein").

The nucleotide sequence of vo18\_1 as presently determined is reported in SEQ ID NO:65, and includes a poly(A) tail. What applicants presently believe to be the proper  
10 reading frame and the predicted amino acid sequence of the vo18\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:66. Amino acids 10 to 22 of SEQ ID NO:66 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
15 vo18\_1 should be approximately 624 bp.

The nucleotide sequence disclosed herein for vo18\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo18\_1 demonstrated at least some similarity with sequences identified as AI198956 (qf66h01.x1 Soares\_testis\_NHT Homo sapiens cDNA clone  
20 IMAGE 1755025 3', mRNA sequence). Based upon sequence similarity, vo18\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vo19\_1"

A polynucleotide of the present invention has been identified as clone "vo19\_1".  
25 vo19\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo19\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo19\_1 protein").

The nucleotide sequence of vo19\_1 as presently determined is reported in SEQ ID  
30 NO:67, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo19\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:68. Amino acids 8 to 20

of SEQ ID NO:68 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo19\_1 should be approximately 1957 bp.

5       The nucleotide sequence disclosed herein for vo19\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo19\_1 demonstrated at least some similarity with sequences identified as AI524085 (th01e09.x1 NCI\_CGAP\_CLL1 Homo sapiens cDNA clone IMAGE:2117032 3', mRNA sequence) and V42646 (DNA encoding a human  
10 pathogenesis-related protein designated HPRP). The predicted amino acid sequence disclosed herein for vo19\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo19\_1 protein demonstrated at least some similarity to sequences identified as U16307 (glioma pathogenesis-related protein [Homo sapiens] and W63115 (A human pathogenesis-related  
15 protein designated HPRP). Based upon sequence similarity, vo19\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "vo22\_1"

A polynucleotide of the present invention has been identified as clone "vo22\_1".  
20 vo22\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo22\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo22\_1 protein").

The nucleotide sequence of vo22\_1 as presently determined is reported in SEQ ID  
25 NO:69, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo22\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:70. Amino acids 6 to 18 of SEQ ID NO:70 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 19.

30       If one of the "G" nucleotides at positions 385 and 386 of SEQ ID NO:69 were deleted, and the "G" residue at position 312 of SEQ ID NO:69 changed to a "T", another potential vo22\_1 reading frame and predicted amino acid sequence, encoded by what

would then be basepairs 104 to 430 of SEQ ID NO:69, is reported in SEQ ID NO:188. Amino acids 8 to 20, amino acids 7 to 19, amino acids 6 to 18, and amino acids 9 to 21 of SEQ ID NO:188 are predicted leader/signal sequences, with the predicted mature amino acid sequence beginning at amino acid 21, or at amino acid 20, or at amino acid 19, or at amino acid 22, respectively. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:188.

Another potential vo22\_1 reading frame and predicted amino acid sequence, encoded by basepairs 1150 to 1357 of SEQ ID NO:69, is reported in SEQ ID NO:189. Amino acids 3 to 15 of SEQ ID NO:189 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:189.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo22\_1 should be approximately 2091 bp.

The nucleotide sequence disclosed herein for vo22\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo22\_1 demonstrated at least some similarity with sequences identified as AA706247 (ah28c11.s1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 1240148 3', mRNA sequence) and V34194 (Human secreted protein gene 41 clone HNTME13). The predicted amino acid sequence disclosed herein for vo22\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo22\_1 protein demonstrated at least some similarity to sequences identified as AF01644 (No definition line found [Caenorhabditis elegans]) and W75155 (Human secreted protein encoded by gene 41 clone HNTME13). Based upon sequence similarity, vo22\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts 9 potential transmembrane domains within the vo22\_1 protein sequence, centered around amino acids 50, 120, 165, 250, 275, 309, 356, 374, and 392 of SEQ ID NO:70, respectively.

Clone "vo23\_1"

A polynucleotide of the present invention has been identified as clone "vo23\_1". vo23\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo23\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo23\_1 protein").

The nucleotide sequence of vo23\_1 as presently determined is reported in SEQ ID NO:71, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo23\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:72.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo23\_1 should be approximately 2598 bp.

The nucleotide sequence disclosed herein for vo23\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo23\_1 demonstrated at least some similarity with sequences identified as T23658 (Human gene signature HUMGS05523), W81246 (zd85b01.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 347401 5', mRNA sequence), and Z84488 (Human DNA sequence from PAC 93H18 on chromosome 6 contains ESTs heterochromatin protein HP1Hs-gamma pseudogene, STS and CpG island). Based upon sequence similarity, vo23\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the vo23\_1 protein sequence, one centered around amino acid 428 and another around amino acid 472 of SEQ ID NO:72.

Clone "vo24\_1"

A polynucleotide of the present invention has been identified as clone "vo24\_1". vo24\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo24\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo24\_1 protein").

The nucleotide sequence of vo24\_1 as presently determined is reported in SEQ ID NO:73, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the vo24\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:74. Amino acids 10 to 22 of SEQ ID NO:74 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23.

5           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo24\_1 should be approximately 3484 bp.

          The nucleotide sequence disclosed herein for vo24\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo24\_1 demonstrated at least some similarity with sequences  
10 identified as AC003117 (\*\*\*) SEQUENCING IN PROGRESS (\*\*\*) Human chromosome 1 BAC 308G1 genomic sequence; HTGS phase 1, 3 unordered pieces), V10696 (Human 3.5 kB DNA fragment predicted to contain CH1-9a11-2 gene), and Z94054 (Human DNA sequence from PAC 125H23 on chromosome 1q24-1q25). The predicted amino acid sequence disclosed herein for vo24\_1 was searched against the GenPept and GeneSeq  
15 amino acid sequence databases using the BLASTX search protocol. The predicted vo24\_1 protein demonstrated at least some similarity to sequences identified as W58774 (Human breast cancer gene CH1-9a11-2 protein fragment #1). Based upon sequence similarity, vo24\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vo24\_1 indicates that it may contain one or more of the following  
20 repetitive elements: Alu, Mer33.

#### Clone "vo25\_1"

          A polynucleotide of the present invention has been identified as clone "vo25\_1". vo25\_1 was isolated from a human adult pancreas cDNA library and was identified as  
25 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo25\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo25\_1 protein").

          The nucleotide sequence of vo25\_1 as presently determined is reported in SEQ ID NO:75, and includes a poly(A) tail. What applicants presently believe to be the proper  
30 reading frame and the predicted amino acid sequence of the vo25\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:76. Amino acids 11 to 23

of SEQ ID NO:76 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo25\_1 should be approximately 1200 bp.

5        The nucleotide sequence disclosed herein for vo25\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo25\_1 demonstrated at least some similarity with sequences identified as AI300566 (qn56a09.x1 NCI\_CGAP\_Kid5 Homo sapiens cDNA clone IMAGE 1902232 3' similar to WP C35D10.1 CE01190 ;, mRNA sequence), V34218  
10 (Human secreted protein gene 65 clone HSREG44), and Z55702 (H.sapiens CpG island DNA genomic MseI fragment, clone 58e10, forward read cpg58e10.ft1a). The predicted amino acid sequence disclosed herein for vo25\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo25\_1 protein demonstrated at least some similarity to sequences identified as  
15 U21324 (similar to S. cerevisiae hypothetical protein YKL166 [Caenorhabditis elegans]) and W57893 (Protein of clone AT340\_1). Based upon sequence similarity, vo25\_1 proteins and each similar protein or peptide may share at least some activity. Motifs analysis detected an ATP/GTP-binding site motif A (P-loop) centered around residue 229 of SEQ ID NO:76. The TopPredII computer program predicts a potential transmembrane  
20 domain within the vo25\_1 protein sequence centered around amino acid 170 of SEQ ID NO:76.

#### Clone "vo26\_1"

A polynucleotide of the present invention has been identified as clone "vo26\_1".  
25 vo26\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo26\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo26\_1 protein").

The nucleotide sequence of vo26\_1 as presently determined is reported in SEQ ID  
30 NO:77, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo26\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:78. Amino acids 13 to 25

of SEQ ID NO:78 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 26.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo26\_1 should be approximately 2503 bp.

5       The nucleotide sequence disclosed herein for vo26\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo26\_1 demonstrated at least some similarity with sequences identified as AC004707 (Homo sapiens chromosome 17, clone hRPC.117\_B\_12, complete sequence), AI160442 (qc08g02.x1 Soares\_fetal\_heart\_NbHH19W Homo sapiens cDNA  
10   clone IMAGE 1709042 3' similar to SW RM02\_YEAST P12687 MITOCHONDRIAL 60S RIBOSOMAL PROTEIN L2 PRECURSOR; mRNA sequence), and T23473 (Human gene signature HUMGS05312). The predicted amino acid sequence disclosed herein for vo26\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo26\_1 protein demonstrated at least  
15   some similarity to sequences identified as L37877 (ribosomal protein L27 [Filobasidiella neoformans]). Based upon sequence similarity, vo26\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vo26\_1 indicates that it may contain a Mir repeat.

20       Clone "vp23\_1"

      A polynucleotide of the present invention has been identified as clone "vp23\_1". vp23\_1 was isolated from a human adult prostate cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vp23\_1 is a full-length clone, including the  
25   entire coding sequence of a secreted protein (also referred to herein as "vp23\_1 protein").

      The nucleotide sequence of vp23\_1 as presently determined is reported in SEQ ID NO:79, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vp23\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:80. Amino acids 5 to 17  
30   of SEQ ID NO:80 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted

leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vp23\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vp23\_1 should be approximately 1220 bp.

5        The nucleotide sequence disclosed herein for vp23\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vp23\_1 demonstrated at least some similarity with sequences identified as AL021578 (Human DNA sequence from clone 453C12 on chromosome 20q12-13.12 Contains SDC4 (syndecan 4 (amphiglycan, ryudocan)), predicts a gene like  
10    the mouse transcription factor RBP-L). Based upon sequence similarity, vp23\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vp23\_1 indicates that it may contain an Alu repetitive element.

#### Clone "vq7\_1"

15        A polynucleotide of the present invention has been identified as clone "vq7\_1". vq7\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq7\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq7\_1 protein").

20        The nucleotide sequence of vq7\_1 as presently determined is reported in SEQ ID NO:81, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq7\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:82. Amino acids 9 to 21 of SEQ ID NO:82 are a predicted leader/signal sequence, with the predicted mature amino  
25    acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq7\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq7\_1 should be approximately 1326 bp.

30        The nucleotide sequence disclosed herein for vq7\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq7\_1 demonstrated at least some similarity with sequences



identified as AA036918 (zk32e03.r1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 484540 5', mRNA sequence). The predicted amino acid sequence disclosed herein for vq7\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vq7\_1 protein demonstrated at least  
5 some similarity to sequences identified as AF142780 (butyrophilin-like protein [Mus musculus]). Butyrophilin is a glycoprotein of the immunoglobulin superfamily that is secreted in association with the milk-fat-globule membrane from mammary epithelial cells (Ogg *et al.*, 1996, *Mamm. Genome* 7 (12): 900-905, which is incorporated by reference herein). Based upon sequence similarity, vq7\_1 proteins and each similar protein or  
10 peptide may share at least some activity. The nucleotide sequence of vq7\_1 indicates that it may contain a repetitive element.

#### Clone "vq8\_1"

A polynucleotide of the present invention has been identified as clone "vq8\_1".  
15 vq8\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq8\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq8\_1 protein").

The nucleotide sequence of vq8\_1 as presently determined is reported in SEQ ID  
20 NO:83, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq8\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:84. Amino acids 10 to 22 of SEQ ID NO:84 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted  
25 leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq8\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq8\_1 should be approximately 695 bp.

The nucleotide sequence disclosed herein for vq8\_1 was searched against the  
30 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq8\_1 demonstrated at least some similarity with sequences identified as AA433968 (zw23f07.r1 Soares ovary tumor NbHOT Homo sapiens cDNA

clone 770149 5', mRNA sequence) and V69618 (Human secreted protein gene 8 clone HLHCM89). The predicted amino acid sequence disclosed herein for vq8\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vq8\_1 protein demonstrated at least some similarity to sequences identified as W83953 (Polypeptide encoded by gene 7 clone HJPDJ64). Based upon sequence similarity, vq8\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vq9\_1"

10 A polynucleotide of the present invention has been identified as clone "vq9\_1". vq9\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq9\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq9\_1 protein").

15 The nucleotide sequence of vq9\_1 as presently determined is reported in SEQ ID NO:85, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq9\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:86. Amino acids 5 to 17 of SEQ ID NO:86 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq9\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq9\_1 should be approximately 1218 bp.

25 The nucleotide sequence disclosed herein for vq9\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq9\_1 demonstrated at least some similarity with sequences identified as AA769310 (nz39f03.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:1290173, mRNA sequence). The predicted amino acid sequence disclosed herein for vq9\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vq9\_1 protein demonstrated at least some similarity to sequences identified as U79260 (unknown [Homo sapiens]) and

W48351 (Human breast cancer related protein BCRB2). Based upon sequence similarity, vq9\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vq10\_1"

5           A polynucleotide of the present invention has been identified as clone "vq10\_1". vq10\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq10\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq10\_1 protein").

10           The nucleotide sequence of vq10\_1 as presently determined is reported in SEQ ID NO:87, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq10\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:88. Amino acids 6 to 18 of SEQ ID NO:88 are a predicted leader/signal sequence, with the predicted mature amino

15           acid sequence beginning at amino acid 19. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq10\_1 protein.

          Another potential reading frame, encoded by nucleotides 331 to 834 of SEQ ID NO:87, is reported as the amino acid sequence of SEQ ID NO:190. Amino acids 29 to 41

20           of SEQ ID NO:190 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 42. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:190.

25           If one nucleotide was deleted from the group of nucleotides at positions 330 and 331 of SEQ ID NO:87, another potential reading frame would be created from what would then be nucleotides 18 to 836, with a predicted amino acid sequence reported as SEQ ID NO:191. Amino acids 6 to 18 of SEQ ID NO:191 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 19. Due to the

30           hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:191.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq10\_1 should be approximately 1516 bp.

The nucleotide sequence disclosed herein for vq10\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq10\_1 demonstrated at least some similarity with sequences identified as AA359702 (EST68843 Fetal lung II Homo sapiens cDNA 5' end similar to similar to pulmonary surfactant protein B, mRNA sequence), I08571 (Sequence 14 from Patent WO 8706588), and Q79287 (Human pulmonary surfactant protein B (SPB)). The predicted amino acid sequence disclosed herein for vq10\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vq10\_1 protein demonstrated at least some similarity to sequences identified as J02761 (pulmonary surfactant-associated protein SP-B [Homo sapiens]) and P70664 (6kd pulmonary surfactant protein). Pulmonary surfactant associated proteins such as SP-B promote alveolar stability by lowering the surface tension at the air-liquid interface in the peripheral air spaces. Based upon sequence similarity, vq10\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of vq10\_1 indicates that it may contain an Alu repetitive element.

#### Clone "vq13\_1"

A polynucleotide of the present invention has been identified as clone "vq13\_1". vq13\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq13\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq13\_1 protein").

The nucleotide sequence of vq13\_1 as presently determined is reported in SEQ ID NO:89, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq13\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:90. Amino acids 10 to 22 of SEQ ID NO:90 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq13\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq13\_1 should be approximately 2284 bp.

The nucleotide sequence disclosed herein for vq13\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
5 FASTA search protocols. vq13\_1 demonstrated at least some similarity with sequences identified as AA928678 (on48e07.s1 NCI\_CGAP\_Co8 Homo sapiens cDNA clone IMAGE 1559940 3', mRNA sequence), AB023187 (Homo sapiens mRNA for KIAA0970 protein, complete cds), and T19039 (Human gene signature HUMGS00046). Based upon  
10 sequence similarity, vq13\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "vq16\_1"

A polynucleotide of the present invention has been identified as clone "vq16\_1". vq16\_1 was isolated from a human adult lung cDNA library and was identified as  
15 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq16\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq16\_1 protein").

The nucleotide sequence of vq16\_1 as presently determined is reported in SEQ ID NO:91, and includes a poly(A) tail. What applicants presently believe to be the proper  
20 reading frame and the predicted amino acid sequence of the vq16\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:92. Amino acids 34 to 46 of SEQ ID NO:92 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 47. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted  
25 leader/signal sequence not be separated from the remainder of the vq16\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq16\_1 should be approximately 1087 bp.

The nucleotide sequence disclosed herein for vq16\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
30 FASTA search protocols. vq16\_1 demonstrated at least some similarity with sequences identified as AA400700 (zu70g11.r1 Soares\_testis\_NHT Homo sapiens cDNA clone IMAGE:743396 5' similar to WP:R05D3.2 CE00281; mRNA sequence). The predicted

amino acid sequence disclosed herein for vq16\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vq16\_1 protein demonstrated at least some similarity to sequences identified as AF05611 (unknown [Fugu rubripes]). Based upon sequence similarity, vq16\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three additional potential transmembrane domains within the vq16\_1 protein sequence, centered around amino acids 90, 134, and 174 of SEQ ID NO:92, respectively.

10        Clone "vq19\_1"

A polynucleotide of the present invention has been identified as clone "vq19\_1". vq19\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq19\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq19\_1 protein").

The nucleotide sequence of vq19\_1 as presently determined is reported in SEQ ID NO:93, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq19\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:94. Amino acids 11 to 23 of SEQ ID NO:94 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq19\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq19\_1 should be approximately 1833 bp.

The nucleotide sequence disclosed herein for vq19\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq19\_1 demonstrated at least some similarity with sequences identified as AA577696 (nn22h03.s1 NCI\_CGAP\_Co12 Homo sapiens cDNA clone IMAGE:1084661 3' similar to contains Alu repetitive element; mRNA sequence. Based upon sequence similarity, vq19\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential

transmembrane domains within the vq19\_1 protein sequence centered around amino acid 214 of SEQ ID NO:94. The nucleotide sequence of vq19\_1 indicates that it may contain an Alu repetitive element.

5           Clone "vq20\_1"

A polynucleotide of the present invention has been identified as clone "vq20\_1". vq20\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq20\_1 is a full-length clone, including the  
10 entire coding sequence of a secreted protein (also referred to herein as "vq20\_1 protein").

The nucleotide sequence of vq20\_1 as presently determined is reported in SEQ ID NO:95, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq20\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:96. Amino acids 10 to 22  
15 of SEQ ID NO:96 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq20\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20 vq20\_1 should be approximately 1275 bp.

The nucleotide sequence disclosed herein for vq20\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq20\_1 demonstrated at least some similarity with sequences identified as AA826249 (of11c04.s1 NCI\_CGAP\_Co12 Homo sapiens cDNA clone  
25 IMAGE 1420806 3' similar to TR Q13445 Q13445 PUTATIVE T1/ST2 RECEPTOR BINDING PROTEIN PRECURSOR; mRNA sequence), AI129838 (qc49h11.x1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone IMAGE:1712997 3' similar to TR:Q13445 Q13445 PUTATIVE T1/ST2 RECEPTOR BINDING PROTEIN PRECURSOR; mRNA sequence), U41805 (Mus musculus putative T1/ST2 receptor  
30 binding protein precursor mRNA, partial cds), and V17729 (Human T1 receptor-like ligand II cDNA). The predicted amino acid sequence disclosed herein for vq20\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the

BLASTX search protocol. The predicted vq20\_1 protein demonstrated at least some similarity to sequences identified as U41804 (putative T1/ST2 receptor binding protein precursor [Homo sapiens]) and W48335 (Human T1 receptor-like ligand II). T1/ST2 is a receptor-like molecule homologous to the type I interleukin-1 receptor (Gayle *et al.*, 1996, *J. Biol. Chem.* **271** (10): 5784-5789, which is incorporated by reference herein). Based upon sequence similarity, vq20\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the vq20\_1 protein sequence centered around amino acid 208 of SEQ ID NO:96.

10

#### Clone "vq21\_1"

A polynucleotide of the present invention has been identified as clone "vq21\_1". vq21\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq21\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq21\_1 protein").

The nucleotide sequence of vq21\_1 as presently determined is reported in SEQ ID NO:97, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq21\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:98. Amino acids 16 to 28 of SEQ ID NO:98 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 29. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq21\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq21\_1 should be approximately 1230 bp.

The nucleotide sequence disclosed herein for vq21\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq21\_1 demonstrated at least some similarity with sequences identified as AA149768 (zo01g05.s1 Stratagene colon (#937204) Homo sapiens cDNA clone IMAGE 566456 3' similar to contains Alu repetitive element; mRNA sequence), AC005282 (Homo sapiens clone DJ0826E18, WORKING DRAFT SEQUENCE, 4



unordered pieces), T25413 (Human gene signature HUMGS07579). The predicted amino acid sequence disclosed herein for vq21\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vq21\_1 protein demonstrated at least some similarity to sequences identified as U67577 (cell division protein FtsJ [Methanococcus jannaschii]). Based upon sequence similarity, vq21\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "vr2\_1"

A polynucleotide of the present invention has been identified as clone "vr2\_1". vr2\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vr2\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vr2\_1 protein").

The nucleotide sequence of vr2\_1 as presently determined is reported in SEQ ID NO:99. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vr2\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:100.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vr2\_1 should be approximately 1382 bp.

The nucleotide sequence disclosed herein for vr2\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No significant similarities were identified in the databases. The TopPredII computer program predicts a potential transmembrane domain within the vr2\_1 protein sequence centered around amino acid 85 of SEQ ID NO:100. The nucleotide sequence of vr2\_1 indicates that it may contain one or more of the following repetitive elements: Alu, MER2, MER4B.

#### Clone "vc69\_1"

A polynucleotide of the present invention has been identified as clone "vc69\_1". vc69\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the

amino acid sequence of the encoded protein. vc69\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc69\_1 protein").

The nucleotide sequence of vc69\_1 as presently determined is reported in SEQ ID NO:101, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the vc69\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:102. Amino acids 7 to 19 of SEQ ID NO:102 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the  
10 predicted leader/signal sequence not be separated from the remainder of the vc69\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc69\_1 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for vc69\_1 was searched against the  
15 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc69\_1 demonstrated at least some similarity with sequences identified as AB023138 (Homo sapiens mRNA for KIAA0921 protein, partial cds), and AI421941 (tf45c01.x1 NCI\_CGAP\_Brn23 Homo sapiens cDNA clone IMAGE 2099136 3' similar to TR Q63376 Q63376 NEUREXIN II-BETA-A PRECURSOR; mRNA  
20 sequence). The predicted amino acid sequence disclosed herein for vc69\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vc69\_1 protein demonstrated at least some similarity to sequences identified as AB02313 (KIAA0921 protein [Homo sapiens]), and various isoforms of *Rattus norvegicus* neurexin II protein. Based upon sequence similarity,  
25 vc69\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "vc71\_1"

A polynucleotide of the present invention has been identified as clone "vc71\_1". vc71\_1 was isolated from a human fetal brain cDNA library and was identified as  
30 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vc71\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vc71\_1 protein").

The nucleotide sequence of vc71\_1 as presently determined is reported in SEQ ID NO:103, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vc71\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:104. Amino acids 2 to 14 of SEQ ID NO:104 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vc71\_1 protein.

10           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vc71\_1 should be approximately 760 bp.

          The nucleotide sequence disclosed herein for vc71\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vc71\_1 demonstrated at least some similarity with sequences identified as AI393859 (tg65f04.x1 Soares\_NhHMPu\_S1 Homo sapiens cDNA clone IMAGE 2113663 3', mRNA sequence) and AL050018 (Homo sapiens mRNA; cDNA DKFZp564B116 (from clone DKFZp564B116)). Based upon sequence similarity, vc71\_1 proteins and each similar protein or peptide may share at least some activity.

20           Clone "vo27\_1"

          A polynucleotide of the present invention has been identified as clone "vo27\_1". vo27\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo27\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo27\_1 protein").

          The nucleotide sequence of vo27\_1 as presently determined is reported in SEQ ID NO:105, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo27\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:106. Amino acids 13 to 25 of SEQ ID NO:106 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 26. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the

predicted leader/signal sequence not be separated from the remainder of the vo27\_1 protein.

Another potential reading frame, encoded by nucleotides 1665 to 1844 of SEQ ID NO:105, is reported as the amino acid sequence of SEQ ID NO:192. Amino acids 4 to 16 of SEQ ID NO:192 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17; amino acids 28 to 40 of SEQ ID NO:192 are also a possible leader/signal sequence, with the predicted mature amino acid sequence beginning in that case at amino acid 41. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the protein of SEQ ID NO:192.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo27\_1 should be approximately 2433 bp.

The nucleotide sequence disclosed herein for vo27\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo27\_1 demonstrated at least some similarity with sequences identified as AC007621 (Homo sapiens clone RPCI11-757G14, WORKING DRAFT SEQUENCE, 142 unordered pieces), AI207832 (ao89g11.x1 Schiller meningioma Homo sapiens cDNA clone IMAGE 1953092 3' similar to contains Alu repetitive element; mRNA sequence), and X80059 (Human PRO361 nucleotide sequence). Based upon sequence similarity, vo27\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vo27\_1 protein sequence centered around amino acid 400 of SEQ ID NO:106. The nucleotide sequence of vo27\_1 indicates that it may contain an Alu repetitive element.

25

#### Clone "vo31\_1"

A polynucleotide of the present invention has been identified as clone "vo31\_1". vo31\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo31\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo31\_1 protein").

30

The nucleotide sequence of vo31\_1 as presently determined is reported in SEQ ID NO:107, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo31\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:108. Amino acids 7 to 19 of SEQ ID NO:108 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vo31\_1 protein.

10 Another potential reading frame, encoded by nucleotides 1937 to 3007 of SEQ ID NO:107, is reported as the amino acid sequence of SEQ ID NO:193.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo31\_1 should be approximately 3222 bp.

The nucleotide sequence disclosed herein for vo31\_1 was searched against the  
15 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo31\_1 demonstrated at least some similarity with sequences identified as AF022147 (*Rattus norvegicus* uterus-ovary specific putative transmembrane protein (uo) mRNA, complete cds), AI417638 (tg80e01.x1 Soares\_NhHMPu\_S1 Homo sapiens cDNA clone IMAGE 2115096 3' similar to TR O35360 O35360 UTERUS-  
20 OVARY SPECIFIC PUTATIVE TRANSMEMBRANE PROTEIN; mRNA sequence), and X52248 (Protein PRO257 cDNA clone DNA35841-1173). The predicted amino acid sequence disclosed herein for vo31\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo31\_1 protein demonstrated at least some similarity to sequences identified as AF02214  
25 (uterus-ovary specific putative transmembrane protein [*Rattus norvegicus*]) and Y13377 (Amino acid sequence of protein PRO257). Based upon sequence similarity, vo31\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the protein sequence of SEQ ID NO:193, centered around amino acid 328 of SEQ ID NO:193.  
30 Hidden markov model analysis indicates the presence of Zona-pellucida-like domains at amino acids 26-115 and 146-273 of SEQ ID NO:193. The nucleotide sequence of vo31\_1 indicates that it may contain a Mer5a repetitive element.

Clone "vo32\_1"

A polynucleotide of the present invention has been identified as clone "vo32\_1". vo32\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo32\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo32\_1 protein").

The nucleotide sequence of vo32\_1 as presently determined is reported in SEQ ID NO:109, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo32\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:110. Amino acids 4 to 16 of SEQ ID NO:110 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vo32\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vo32\_1 should be approximately 1868 bp.

The nucleotide sequence disclosed herein for vo32\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vo32\_1 demonstrated at least some similarity with sequences identified as AF028740 (Mus musculus olfactomedin mRNA, complete cds), AI078144 (oz30b06.x1 Soares\_total\_fetus\_Nb2HF8\_9w Homo sapiens cDNA clone IMAGE 1676819 3' similar to TR Q99784 Q99784 NEURONAL OLFACTOMEDIN-RELATED ER LOCALIZED PROTEIN; mRNA sequence), AI869993 (wl63e09.x1 NCI\_CGAP Brn25 Homo sapiens cDNA clone IMAGE:2429608 3' similar to SW:NOMR\_HUMAN Q99784 NEURONAL OLFACTOMEDIN-RELATED ER LOCALIZED PROTEIN; mRNA sequence), and V34217 (Human secreted protein gene 64 clone HSLDJ95). The predicted amino acid sequence disclosed herein for vo32\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vo32\_1 protein demonstrated at least some similarity to sequences identified as U03416 (neuronal olfactomedin-related ER localized protein [Rattus norvegicus]) and W75120 (Human secreted protein encoded by gene 64 clone HSLDJ95). Based upon

sequence similarity, vo32\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "vo33\_1"

5           A polynucleotide of the present invention has been identified as clone "vo33\_1". vo33\_1 was isolated from a human adult pancreas cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vo33\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vo33\_1 protein").

10           The nucleotide sequence of vo33\_1 as presently determined is reported in SEQ ID NO:111, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vo33\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:112. Amino acids 5 to 17 of SEQ ID NO:112 are a predicted leader/signal sequence, with the predicted mature  
15 amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vo33\_1 protein.

          The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20 vo33\_1 should be approximately 2879 bp.

The nucleotide sequence disclosed herein for vo33\_1 was searched against the  
GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
FASTA search protocols. vo33\_1 demonstrated at least some similarity with sequences  
identified as AI225613 (uj13e01.y1 Sugano mouse kidney mkia Mus musculus cDNA  
25 clone IMAGE:1907928 5' similar to TR:Q14624 Q14624 INTER-ALPHA-TRYPSIN  
INHIBITOR FAMILY HEAVY CHAIN-RELATED PROTEIN; mRNA sequence) and  
X80054 (Human PRO354 nucleotide sequence). The predicted amino acid sequence  
disclosed herein for vo33\_1 was searched against the GenPept and GeneSeq amino acid  
sequence databases using the BLASTX search protocol. The predicted vo33\_1 protein  
30 demonstrated at least some similarity to sequences identified as D38535 (PK-120 precursor  
[Homo sapiens]), Y11545 (inter-alpha-inhibitor heavy-chain H2 [Sus scrofa]), and the H2  
proteins of several species, including *Homo sapiens*. Based upon sequence similarity,

vo33\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the vo33\_1 protein sequence centered around amino acid 386 of SEQ ID NO:112. The nucleotide sequence of vo33\_1 indicates that it may contain an Alu repetitive element.

5

Clone "vq23\_1"

A polynucleotide of the present invention has been identified as clone "vq23\_1". vq23\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq23\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq23\_1 protein").

The nucleotide sequence of vq23\_1 as presently determined is reported in SEQ ID NO:113, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq23\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:114. Amino acids 18 to 30 of SEQ ID NO:114 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 31. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq23\_1 protein.

Another potential reading frame, encoded by nucleotides 1012 to 1518 of SEQ ID NO:113, is reported as the amino acid sequence of SEQ ID NO:194. Amino acids 83 to 94 of SEQ ID NO:194 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 95. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of protein of SEQ ID NO:194.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq23\_1 should be approximately 1793 bp.

The nucleotide sequence disclosed herein for vq23\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq23\_1 demonstrated at least some similarity with sequences



identified as AA625521 (af72f02.r1 Soares\_NhHMPu\_S1 Homo sapiens cDNA clone IMAGE 1047579 5', mRNA sequence) and AC002364 (Homo sapiens Xp22 Cosmids U15E4, U115H5, U132E12, U115B9 (Lawrence Livermore human cosmid library) complete sequence). Based upon sequence similarity, vq23\_1 proteins and each similar  
5 protein or peptide may share at least some activity. The nucleotide sequence of vq23\_1 indicates that it may contain an Alu repetitive element.

Clone "vq24\_1"

A polynucleotide of the present invention has been identified as clone "vq24\_1".  
10 vq24\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq24\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq24\_1 protein").

The nucleotide sequence of vq24\_1 as presently determined is reported in SEQ ID  
15 NO:115, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq24\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:116. Amino acids 5 to 17 of SEQ ID NO:116 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the  
20 predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq24\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq24\_1 should be approximately 2168 bp.

25 The nucleotide sequence disclosed herein for vq24\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq24\_1 demonstrated at least some similarity with sequences identified as N29315 (yx43d06.r1 Soares melanocyte 2NbHM Homo sapiens cDNA clone IMAGE:264491 5' similar to SP:SW:FCG1\_HUMAN P12315 HIGH AFFINITY  
30 IMMUNOGLOBULIN GAMMA FC RECEPTOR I 'B FORM' PRECURSOR; mRNA sequence). The predicted amino acid sequence disclosed herein for vq24\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX

search protocol. The predicted vq24\_1 protein demonstrated at least some similarity to sequences identified as AF14317 (high affinity immunoglobulin gamma Fc receptor I [Mus musculus]) and R12428 (Hybrid Fc(gamma)RII/I receptor). Based upon sequence similarity, vq24\_1 proteins and each similar protein or peptide may share at least some activity. Hidden markov model analysis detects immunoglobulin superfamily signatures in the vq24\_1 protein sequence from amino acid 92 to amino acid 145, and from amino acid 185 to amino acid 242, of SEQ ID NO:116. The nucleotide sequence of vq24\_1 indicates that it may contain one or more of the following repetitive elements: Mer, MLT1a.

10

#### Clone "vq26\_1"

A polynucleotide of the present invention has been identified as clone "vq26\_1". vq26\_1 was isolated from a human adult lung cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vq26\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vq26\_1 protein").

The nucleotide sequence of vq26\_1 as presently determined is reported in SEQ ID NO:117, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vq26\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:118. Amino acids 9 to 21 of SEQ ID NO:118 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vq26\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vq26\_1 should be approximately 1419 bp.

The nucleotide sequence disclosed herein for vq26\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vq26\_1 demonstrated at least some similarity with sequences identified as AA191552 (zp82g04.s1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone IMAGE 626742 3', mRNA sequence) and AA573741 (nk07a05.s1 NCI\_CGAP\_Co2

Homo sapiens cDNA clone IMAGE:1012784 3', mRNA sequence). Based upon sequence similarity, vq26\_1 proteins and each similar protein or peptide may share at least some activity.

5           Deposit of Clones

Clones vc62\_1, vp10\_1, vp11\_1, vp13\_1, vp16\_1, vp21\_1, vp22\_1, vq2\_1, vq3\_1, vq5\_1, vq6\_1, and vr1\_1 were deposited on February 17, 1999 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession  
10   number ATCC 207114, from which each clone comprising a particular polynucleotide is obtainable.

Clone vc63\_1 was deposited on February 17, 1999 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number  
15   ATCC 207115, from which the vc63\_1 clone comprising a particular polynucleotide is obtainable.

Clones vb25\_1, vb27\_1, vb28\_1, vb29\_1, vb30\_1, vc67\_1, vf4\_1, vg3\_1, vo2\_1, vo3\_1, vo5\_1, vo6\_1, and vo9\_1 were deposited on July 15, 1999 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia  
20   20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number PTA-362, from which each clone comprising a particular polynucleotide is obtainable.

Clones vo11\_1, vo12\_1, vo13\_1, vo14\_1, vo15\_1, vo16\_1, vo18\_1, vo19\_1, vo22\_1, vo23\_1, vo24\_1, vo25\_1, and vo26\_1 were deposited on July 15, 1999 with the  
25   ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number PTA-366, from which each clone comprising a particular polynucleotide is obtainable.

Clones vp23\_1, vq7\_1, vq8\_1, vq9\_1, vq10\_1, vq13\_1, vq16\_1, vq19\_1, vq20\_1, vq21\_1, and vr2\_1 were deposited on July 15, 1999 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.)  
30

as an original deposit under the Budapest Treaty and were given the accession number PTA-368, from which each clone comprising a particular polynucleotide is obtainable.

Clones vc69\_1, vc71\_1, vo27\_1, vo31\_1, vo32\_1, vo33\_1, vq23\_1, vq24\_1, and vq26\_1 were deposited on December 21, 1999 with the ATCC (American Type Culture  
5 Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number PTA-1075, from which each clone comprising a particular polynucleotide is obtainable.

All restrictions on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent, except for the requirements specified  
10 in 37 C.F.R. § 1.808(b), and the term of the deposit will comply with 37 C.F.R. § 1.806.

Each clone has been transfected into separate bacterial cells (*E. coli*) in these composite deposits. Each clone can be removed from the vector in which it was deposited by performing an EcoRI/NotI digestion (5' site, EcoRI; 3' site, NotI) to produce the appropriate fragment for such clone. Each clone was deposited in either the pED6 or  
15 pNOTs vector depicted in Figures 1A and 1B, respectively. The pED6dpc2 vector ("pED6") was derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning (Kaufman *et al.*, 1991, *Nucleic Acids Res.* 19: 4485-4490); the pNOTs vector was derived from pMT2 (Kaufman *et al.*, 1989, *Mol. Cell. Biol.* 9: 946-958) by deletion of the DHFR sequences, insertion of a new polylinker, and insertion of the M13  
20 origin of replication in the ClaI site. In some instances, the deposited clone can become "flipped" (i.e., in the reverse orientation) in the deposited isolate. In such instances, the cDNA insert can still be isolated by digestion with EcoRI and NotI. However, NotI will then produce the 5' site and EcoRI will produce the 3' site for placement of the cDNA in proper orientation for expression in a suitable vector. The cDNA may also be expressed  
25 from the vectors in which they were deposited.

Bacterial cells containing a particular clone can be obtained from the composite deposit as follows:

An oligonucleotide probe or probes should be designed to the sequence that is known for that particular clone. This sequence can be derived from the sequences  
30 provided herein, or from a combination of those sequences. The sequence of an oligonucleotide probe that was used to isolate or to sequence each full-length clone is identified below, and should be most reliable in isolating the clone of interest.

<u>Clone</u>	<u>Probe Sequence</u>
vc62_1	SEQ ID NO:119
vp10_1	SEQ ID NO:120
vp11_1	SEQ ID NO:121
5 vp13_1	SEQ ID NO:122
vp16_1	SEQ ID NO:123
vp21_1	SEQ ID NO:124
vp22_1	SEQ ID NO:125
vq2_1	SEQ ID NO:126
10 vq3_1	SEQ ID NO:127
vq5_1	SEQ ID NO:128
vq6_1	SEQ ID NO:129
vr1_1	SEQ ID NO:130
vc63_1	SEQ ID NO:131
15 vb25_1	SEQ ID NO:132
vb27_1	SEQ ID NO:133
vb28_1	SEQ ID NO:134
vb29_1	SEQ ID NO:135
vb30_1	SEQ ID NO:136
20 vc67_1	SEQ ID NO:137
vf4_1	SEQ ID NO:138
vg3_1	SEQ ID NO:139
vo2_1	SEQ ID NO:140
vo3_1	SEQ ID NO:141
25 vo5_1	SEQ ID NO:142
vo6_1	SEQ ID NO:143
vo9_1	SEQ ID NO:144
vo11_1	SEQ ID NO:145
vo12_1	SEQ ID NO:146
30 vo13_1	SEQ ID NO:147
vo14_1	SEQ ID NO:148
vo15_1	SEQ ID NO:149

	vo16_1	SEQ ID NO:150
	vo18_1	SEQ ID NO:151
	vo19_1	SEQ ID NO:152
	vo22_1	SEQ ID NO:153
5	vo23_1	SEQ ID NO:154
	vo24_1	SEQ ID NO:155
	vo25_1	SEQ ID NO:156
	vo26_1	SEQ ID NO:157
	vp23_1	SEQ ID NO:158
10	vq7_1	SEQ ID NO:159
	vq8_1	SEQ ID NO:160
	vq9_1	SEQ ID NO:161
	vq10_1	SEQ ID NO:162
	vq13_1	SEQ ID NO:163
15	vq16_1	SEQ ID NO:164
	vq19_1	SEQ ID NO:165
	vq20_1	SEQ ID NO:166
	vq21_1	SEQ ID NO:167
	vr2_1	SEQ ID NO:168

20

In the sequences listed above which include an N at position 2, that position is occupied in preferred probes/primers by a biotinylated phosphoramidite residue rather than a nucleotide (such as, for example, that produced by use of biotin phosphoramidite (1-dimethoxytrityloxy-2-(N-biotinyl-4-aminobutyl)-propyl-3-O-(2-cyanoethyl)-(N,N-diisopropyl)-phosphoramidite) (Glen Research, cat. no. 10-1953)).

25

The design of the oligonucleotide probe should preferably follow these parameters:

- (a) It should be designed to an area of the sequence which has the fewest ambiguous bases ("N's"), if any;
- (b) It should be designed to have a  $T_m$  of approx. 80 ° C (assuming 2° for each A or T and 4 degrees for each G or C).

30

The oligonucleotide should preferably be labeled with  $\gamma$ -<sup>32</sup>P ATP (specific activity 6000 Ci/mmol) and T4 polynucleotide kinase using commonly employed techniques for

labeling oligonucleotides. Other labeling techniques can also be used. Unincorporated label should preferably be removed by gel filtration chromatography or other established methods. The amount of radioactivity incorporated into the probe should be quantitated by measurement in a scintillation counter. Preferably, specific activity of the resulting  
5 probe should be approximately  $4 \times 10^6$  dpm/pmole.

The bacterial culture containing the pool of full-length clones should preferably be thawed and 100  $\mu$ l of the stock used to inoculate a sterile culture flask containing 25 ml of sterile L-broth containing ampicillin at 100  $\mu$ g/ml. The culture should preferably be grown to saturation at 37°C, and the saturated culture should preferably be diluted in fresh  
10 L-broth. Aliquots of these dilutions should preferably be plated to determine the dilution and volume which will yield approximately 5000 distinct and well-separated colonies on solid bacteriological media containing L-broth containing ampicillin at 100  $\mu$ g/ml and agar at 1.5% in a 150 mm petri dish when grown overnight at 37°C. Other known methods of obtaining distinct, well-separated colonies can also be employed.

15 Standard colony hybridization procedures should then be used to transfer the colonies to nitrocellulose filters and lyse, denature and bake them.

The filter is then preferably incubated at 65°C for 1 hour with gentle agitation in 6X SSC (20X stock is 175.3 g NaCl/liter, 88.2 g Na citrate/liter, adjusted to pH 7.0 with NaOH) containing 0.5% SDS, 100  $\mu$ g/ml of yeast RNA, and 10 mM EDTA  
20 (approximately 10 mL per 150 mm filter). Preferably, the probe is then added to the hybridization mix at a concentration greater than or equal to  $1 \times 10^6$  dpm/mL. The filter is then preferably incubated at 65°C with gentle agitation overnight. The filter is then preferably washed in 500 mL of 2X SSC/0.5% SDS at room temperature without agitation, preferably followed by 500 mL of 2X SSC/0.1% SDS at room temperature with gentle  
25 shaking for 15 minutes. A third wash with 0.1X SSC/0.5% SDS at 65°C for 30 minutes to 1 hour is optional. The filter is then preferably dried and subjected to autoradiography for sufficient time to visualize the positives on the X-ray film. Other known hybridization methods can also be employed.

The positive colonies are picked, grown in culture, and plasmid DNA isolated  
30 using standard procedures. The clones can then be verified by restriction analysis, hybridization analysis, or DNA sequencing.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H.U. Saragovi, *et al.*, *Bio/Technology* 10, 773-778 (1992) and in R.S. McDowell, *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein, such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to generate such fusions. For example, a protein - IgM fusion would generate a decavalent form of the protein of the invention.

The present invention also provides both full-length and mature forms of the disclosed proteins. The full-length form of the such proteins is identified in the sequence listing by translation of the nucleotide sequence of each disclosed clone. The mature form(s) of such protein may be obtained by expression of the disclosed full-length polynucleotide (preferably those deposited with the ATCC) in a suitable mammalian cell or other host cell. The sequence(s) of the mature form(s) of the protein may also be determinable from the amino acid sequence of the full-length form.

The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that are transcribed to produce the mRNAs from which cDNA polynucleotide sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited to coding sequences, 5' and 3' untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene" is a gene that has been separated from the adjacent coding sequences, if any, present in the genome of the organism from which the gene was isolated.



The chromosomal location corresponding to the polynucleotide sequences disclosed herein may also be determined, for example by hybridizing appropriately labeled polynucleotides of the present invention to chromosomes *in situ*. It may also be possible to determine the corresponding chromosomal location for a disclosed polynucleotide by identifying significantly similar nucleotide sequences in public databases, such as expressed sequence tags (ESTs), that have already been mapped to particular chromosomal locations. For at least some of the polynucleotide sequences disclosed herein, public database sequences having at least some similarity to the polynucleotide of the present invention have been listed by database accession number. Searches using the GenBank accession numbers of these public database sequences can then be performed at an Internet site provided by the National Center for Biotechnology Information having the address <http://www.ncbi.nlm.nih.gov/UniGene/>, in order to identify "UniGene clusters" of overlapping sequences. Many of the "UniGene clusters" so identified will already have been mapped to particular chromosomal sites.

Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided. The desired change in gene expression can be achieved through the use of antisense polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, *Trends Pharmacol. Sci.* **15**(7): 250-254; Lavarosky *et al.*, 1997, *Biochem. Mol. Med.* **62**(1): 11-22; and Hampel, 1998, *Prog. Nucleic Acid Res. Mol. Biol.* **58**: 1-39; all of which are incorporated by reference herein). The desired change in gene expression can also be achieved through the use of double-stranded ribonucleotide molecules having some complementarity to the mRNA transcribed from the gene, and which interfere with the transcription, stability, or expression of the mRNA ("RNA interference" or "RNAi"; Fire *et al.*, 1998, *Nature* **391** (6669): 806-811; Montgomery *et al.*, 1998, *Proc. Natl. Acad. Sci. USA* **95** (26): 15502-15507; and Sharp, 1999, *Genes Dev.* **13** (2): 139-141; all of which are incorporated by reference herein). Transgenic animals that have multiple copies of the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are also provided (see European

Patent No. 0 649 464 B1, incorporated by reference herein). In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially or completely inactivated, through insertion of extraneous sequences into the corresponding gene(s) or through deletion of all or part of the  
5 corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of transposable elements (Plasterk, 1992, *Bioessays* 14(9): 629-633; Zwaal *et al.*, 1993, *Proc. Natl. Acad. Sci. USA* 90(16): 7431-7435; Clark *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91(2): 719-722; all of which are incorporated by reference herein), or through homologous recombination, preferably  
10 detected by positive/negative genetic selection strategies (Mansour *et al.*, 1988, *Nature* 336: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms are useful for the development of non-human models for the  
15 study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms, part  
20 or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known techniques for determination of such domains from sequence information. For example, the TopPredII computer program can be used to predict the location of  
25 transmembrane domains in an amino acid sequence, domains which are described by the location of the center of the transmembrane domain, with at least ten transmembrane amino acids on each side of the reported central residue(s).

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most  
30 preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid

sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

In particular, sequence identity may be determined using WU-BLAST (Washington University BLAST) version 2.0 software, which builds upon WU-BLAST version 1.4, which in turn is based on the public domain NCBI-BLAST version 1.4 (Altschul and Gish, 1996, Local alignment statistics, Doolittle *ed.*, *Methods in Enzymology* 266: 460-480; Altschul *et al.*, 1990, Basic local alignment search tool, *Journal of Molecular Biology* 215: 403-410; Gish and States, 1993, Identification of protein coding regions by database similarity search, *Nature Genetics* 3: 266-272; Karlin and Altschul, 1993, Applications and statistics for multiple high-scoring segments in molecular sequences, *Proc. Natl. Acad. Sci. USA* 90: 5873-5877; all of which are incorporated by reference herein). WU-BLAST version 2.0 executable programs for several UNIX platforms can be downloaded from <ftp://blast.wustl.edu/blast/executables>. The complete suite of search programs (BLASTP, BLASTN, BLASTX, TBLASTN, and TBLASTX) is provided at that site, in addition to several support programs. WU-BLAST 2.0 is copyrighted and may not be sold or redistributed in any form or manner without the express written consent of the author; but the posted executables may otherwise be freely used for commercial, nonprofit, or academic purposes. In all search programs in the suite -- BLASTP, BLASTN, BLASTX, TBLASTN and TBLASTX -- the gapped alignment routines are integral to the database search itself, and thus yield much better sensitivity and selectivity while producing the more easily interpreted output. Gapping can optionally be turned off in all of these programs, if desired. The default penalty (Q) for a gap of length one is Q=9 for proteins and BLASTP, and Q=10 for BLASTN, but may be changed to any integer value including zero, one through eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. The default per-residue penalty for extending a gap (R) is R=2 for proteins and BLASTP, and R=10 for BLASTN, but may be changed to any integer value including zero, one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. Any

combination of values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps. The default amino acid comparison matrix is BLOSUM62, but other amino acid comparison matrices such as PAM can be utilized.

5           Species homologues of the disclosed polynucleotides and proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide. Preferably, polynucleotide species homologues have at least 60% sequence  
10 identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, and protein species homologues have at least 30% sequence identity (more preferably, at least 45% identity; most preferably at least 60% identity) with the given protein, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides or the amino acid sequences of the proteins when aligned so as to maximize  
15 overlap and identity while minimizing sequence gaps. Species homologues may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species. Preferably, species homologues are those isolated from mammalian species. Most preferably, species homologues are those isolated from certain mammalian species such as, for example, *Pan troglodytes*, *Gorilla*  
20 *gorilla*, *Pongo pygmaeus*, *Hylobates concolor*, *Macaca mulatta*, *Papio papio*, *Papio hamadryas*, *Cercopithecus aethiops*, *Cebus capucinus*, *Aotus trivirgatus*, *Sanguinus oedipus*, *Microcebus murinus*, *Mus musculus*, *Rattus norvegicus*, *Cricetulus griseus*, *Felis*  
*catus*, *Mustela vison*, *Canis familiaris*, *Oryctolagus cuniculus*, *Bos taurus*, *Ovis aries*, *Sus scrofa*, and *Equus caballus*, for which genetic maps have been created allowing the  
25 identification of syntenic relationships between the genomic organization of genes in one species and the genomic organization of the related genes in another species (O'Brien and Seuáñez, 1988, *Ann. Rev. Genet.* 22: 323-351; O'Brien *et al.*, 1993, *Nature Genetics* 3:103-112; Johansson *et al.*, 1995, *Genomics* 25: 682-690; Lyons *et al.*, 1997, *Nature Genetics* 15: 47-56; O'Brien *et al.*, 1997, *Trends in Genetics* 13(10): 393-399; Carver and  
30 Stubbs, 1997, *Genome Research* 7:1123-1137; all of which are incorporated by reference herein).

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotides which also encode proteins which are identical or have significantly similar sequences to those encoded by the disclosed polynucleotides. Preferably, allelic variants have at least  
5 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps. Allelic variants may be isolated and identified by making suitable probes or primers from the sequences provided  
10 herein and screening a suitable nucleic acid source from individuals of the appropriate species.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The present invention also includes polynucleotides that hybridize under reduced  
15 stringency conditions, more preferably stringent conditions, and most preferably highly stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

	Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) <sup>†</sup>	Hybridization Temperature and Buffer <sup>†</sup>	Wash Temperature and Buffer <sup>†</sup>
5	A	DNA:DNA	≥ 50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
	B	DNA:DNA	<50	T <sub>B</sub> *; 1xSSC	T <sub>B</sub> *; 1xSSC
	C	DNA:RNA	≥ 50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	67°C; 0.3xSSC
	D	DNA:RNA	<50	T <sub>D</sub> *; 1xSSC	T <sub>D</sub> *; 1xSSC
	E	RNA:RNA	≥ 50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	70°C; 0.3xSSC
	F	RNA:RNA	<50	T <sub>F</sub> *; 1xSSC	T <sub>F</sub> *; 1xSSC
10	G	DNA:DNA	≥ 50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	65°C; 1xSSC
	H	DNA:DNA	<50	T <sub>H</sub> *; 4xSSC	T <sub>H</sub> *; 4xSSC
	I	DNA:RNA	≥ 50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	67°C; 1xSSC
	J	DNA:RNA	<50	T <sub>J</sub> *; 4xSSC	T <sub>J</sub> *; 4xSSC
	K	RNA:RNA	≥ 50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	67°C; 1xSSC
	L	RNA:RNA	<50	T <sub>L</sub> *; 2xSSC	T <sub>L</sub> *; 2xSSC
15	M	DNA:DNA	≥ 50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	50°C; 2xSSC
	N	DNA:DNA	<50	T <sub>N</sub> *; 6xSSC	T <sub>N</sub> *; 6xSSC
	O	DNA:RNA	≥ 50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	55°C; 2xSSC
	P	DNA:RNA	<50	T <sub>P</sub> *; 6xSSC	T <sub>P</sub> *; 6xSSC
	Q	RNA:RNA	≥ 50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	60°C; 2xSSC
	R	RNA:RNA	<50	T <sub>R</sub> *; 4xSSC	T <sub>R</sub> *; 4xSSC

\*: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

<sup>†</sup>: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub>, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

\*T<sub>B</sub> - T<sub>R</sub>: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T<sub>m</sub>) of the hybrid, where T<sub>m</sub> is determined according to the following equations. For hybrids less than 18 base pairs in length, T<sub>m</sub>(°C) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T<sub>m</sub>(°C) = 81.5 + 16.6(log<sub>10</sub>[Na<sup>+</sup>]) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na<sup>+</sup>] is the concentration of sodium ions in the hybridization buffer ([Na<sup>+</sup>] for 1xSSC = 0.165 M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

The isolated polynucleotide encoding the protein of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable

bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (i.e., from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLabs (Beverly, MA), Pharmacia (Piscataway, NJ) and Invitrogen Corporation (Carlsbad, CA), respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope.



One such epitope ("Flag") is commercially available from the Eastman Kodak Company (New Haven, CT).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are known to those skilled in the art. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Patent No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

## **USES AND BIOLOGICAL ACTIVITY**

The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

### **Research Uses and Utilities**

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, those described in Gyuris *et al.*, 1993, *Cell*

75: 791-803 and in Rossi *et al.*, 1997, *Proc. Natl. Acad. Sci. USA* **94**: 8405-8410, all of which are incorporated by reference herein) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to  
5 determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue  
10 differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or  
15 small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning:  
20 A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

#### Nutritional Uses

25 Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid  
30 preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors  
 5 discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G,  
 10 M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek,  
 15 D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., J. Immunol. 149:3778-3783, 1992;  
 20 Bowman et al., J. Immunol. 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and  
 25 Measurement of mouse and human Interferon  $\gamma$ , Schreiber, R.D. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine  
 30 Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al.,

- Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6 - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of
- 5 human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.
- 10 Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience
- 15 (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

20

#### Immune Stimulating or Suppressing Activity

- A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies
- 25 and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral,
- 30 bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course,

in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, *i.e.*, in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems.

10 Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to regulate immune responses in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

15  
20

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), *e.g.*, preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble,

25  
30

monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow *et al.*, Science 257:789-792 (1992) and Turka *et al.*, Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the

disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosus in MRL/lpr/lpr mice or NZB hybrid mice, murine  
5 autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy.  
10 Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B  
15 lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro*  
20 activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory  
25 signal to, and thereby activate, T cells *in vivo*.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (*e.g.*, sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present  
30 invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an



expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to  
5 target a tumor cell for transfection *in vivo*.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II  
10 molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (*e.g.*, a cytoplasmic-domain truncated portion) of an MHC class I  $\alpha$  chain protein and  $\beta_2$  microglobulin protein or an MHC class II  $\alpha$  chain protein and an MHC class II  $\beta$  chain protein to thereby express MHC class I or MHC class II proteins on the cell surface.  
15 Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (*e.g.*, B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the  
20 activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured  
25 by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte  
30 Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol.

137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 5 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: 10 Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: *In vitro* antibody production, Mond, J.J. and Brunswick, M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, 15 those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, *Immunologic studies in Humans*); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. 20 Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., *Journal of Experimental Medicine* 173:549-559, 1991; Macatonia et al., *Journal of Immunology* 25 154:5071-5079, 1995; Porgador et al., *Journal of Experimental Medicine* 182:255-260, 1995; Nair et al., *Journal of Virology* 67:4062-4069, 1993; Huang et al., *Science* 264:961-965, 1994; Macatonia et al., *Journal of Experimental Medicine* 169:1255-1264, 1989; Bhardwaj et al., *Journal of Clinical Investigation* 94:797-807, 1994; and Inaba et al., *Journal of Experimental Medicine* 172:631-640, 1990.

30 Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz

et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

- 5           Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

10           Hematopoiesis Regulating Activity

- A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and
- 15           proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for
- 20           example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic
- 25           stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction
- 30           with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

- 5        Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.
- 10       Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic
- 15       colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term
- 20       bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

25

#### Tissue Growth Activity

- A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions
- 30       and ulcers.

A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone

fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma  
5 induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of  
10 progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the  
15 protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing  
20 protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also  
25 useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The  
30 compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue.

5 More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include

10 mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of

15 non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac)

20 and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or

25 regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

30 The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium ).

- 5        Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

10        Activin/Inhibin Activity

- A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin  $\alpha$  family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- $\beta$  group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

- 25        The activity of a protein of the invention may, among other means, be measured by the following methods:

- Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al., Nature 321:776-779, 1986; Mason et al., Nature 318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA 83:3091-3095, 1986.

Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

10 A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

#### Hemostatic and Thrombolytic Activity

30 A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation



and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels  
5 (e.g., stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis  
10 Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

#### Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as receptors,  
15 receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs  
20 involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

25 The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and  
30 Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med.

169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

#### Anti-Inflammatory Activity

5 Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or  
10 suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin  
15 lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

#### 20 Cadherin/Tumor Invasion Suppressor Activity

Cadherins are calcium-dependent adhesion molecules that appear to play major roles during development, particularly in defining specific cell types. Loss or alteration of normal cadherin expression can lead to changes in cell adhesion properties linked to tumor growth and metastasis. Cadherin malfunction is also implicated in other human diseases, such as pemphigus  
25 vulgaris and pemphigus foliaceus (auto-immune blistering skin diseases), Crohn's disease, and some developmental abnormalities.

The cadherin superfamily includes well over forty members, each with a distinct pattern of expression. All members of the superfamily have in common conserved extracellular repeats (cadherin domains), but structural differences are found in other parts of the molecule. The  
30 cadherin domains bind calcium to form their tertiary structure and thus calcium is required to mediate their adhesion. Only a few amino acids in the first cadherin domain provide the basis for homophilic adhesion; modification of this recognition site can change the specificity of a cadherin

so that instead of recognizing only itself, the mutant molecule can now also bind to a different cadherin. In addition, some cadherins engage in heterophilic adhesion with other cadherins.

E-cadherin, one member of the cadherin superfamily, is expressed in epithelial cell types. Pathologically, if E-cadherin expression is lost in a tumor, the malignant cells become invasive and the cancer metastasizes. Transfection of cancer cell lines with polynucleotides expressing E-cadherin has reversed cancer-associated changes by returning altered cell shapes to normal, restoring cells' adhesiveness to each other and to their substrate, decreasing the cell growth rate, and drastically reducing anchorage-independent cell growth. Thus, reintroducing E-cadherin expression reverts carcinomas to a less advanced stage. It is likely that other cadherins have the same invasion suppressor role in carcinomas derived from other tissue types. Therefore, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to treat cancer. Introducing such proteins or polynucleotides into cancer cells can reduce or eliminate the cancerous changes observed in these cells by providing normal cadherin expression.

Cancer cells have also been shown to express cadherins of a different tissue type than their origin, thus allowing these cells to invade and metastasize in a different tissue in the body. Proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be substituted in these cells for the inappropriately expressed cadherins, restoring normal cell adhesive properties and reducing or eliminating the tendency of the cells to metastasize.

Additionally, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to generate antibodies recognizing and binding to cadherins. Such antibodies can be used to block the adhesion of inappropriately expressed tumor-cell cadherins, preventing the cells from forming a tumor elsewhere. Such an anti-cadherin antibody can also be used as a marker for the grade, pathological type, and prognosis of a cancer, i.e. the more progressed the cancer, the less cadherin expression there will be, and this decrease in cadherin expression can be detected by the use of a cadherin-binding antibody.

Fragments of proteins of the present invention with cadherin activity, preferably a polypeptide comprising a decapeptide of the cadherin recognition site, and polynucleotides of the present invention encoding such protein fragments, can also be used to block cadherin function by binding to cadherins and preventing them from binding in ways that produce undesirable effects. Additionally, fragments of proteins of the present invention with cadherin activity, preferably truncated soluble cadherin fragments which have been found to be stable in the

circulation of cancer patients, and polynucleotides encoding such protein fragments, can be used to disturb proper cell-cell adhesion.

Assays for cadherin adhesive and invasive suppressor activity include, without limitation, those described in: Hortsch et al. J Biol Chem 270 (32): 18809-18817, 1995; Miyaki et al.

- 5 Oncogene 11: 2547-2552, 1995; Ozawa et al. Cell 63: 1033-1038, 1990.

#### Tumor Inhibition Activity

In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities.

- 10 A protein may inhibit tumor growth directly or indirectly (such as, for example, via antibody-dependent cell-mediated cytotoxicity (ADCC)). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit
- 15 tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth.

#### Other Activities

- A protein of the invention may also exhibit one or more of the following additional
- 20 activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution,
- 25 change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress,
- 30 cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic

lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen  
5 in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

### **ADMINISTRATION AND DOSING**

A protein of the present invention (from whatever source derived, including  
10 without limitation from recombinant and non-recombinant sources) may be used in a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the  
15 effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF,  
20 thrombopoietin, stem cell factor, and erythropoietin. The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or complement its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may  
25 be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

A protein of the present invention may be active in multimers (e.g., heterodimers  
30 or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in combination  
5 with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the  
10 attending physician will decide on the appropriate sequence of administering protein of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of protein of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a  
15 variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of protein of the present invention is administered orally, protein of the present invention will be in the form of a tablet, capsule,  
20 powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal  
25 or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein  
30 of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu\text{g}$  to about 100 mg (preferably about 0.1ng to about 10 mg, more preferably about 0.1  $\mu\text{g}$  to about 1 mg) of protein of the present invention per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the protein of the present invention will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. As used herein, the



term "antibody" includes without limitation a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a single-chain antibody, a CDR-grafted antibody, a humanized antibody, or fragments thereof which bind to the indicated protein. Such term also includes any other species derived from an antibody or antibody sequence which is capable  
5 of binding the indicated protein.

Antibodies to a particular protein can be produced by methods well known to those skilled in the art. For example, monoclonal antibodies can be produced by generation of antibody-producing hybridomas in accordance with known methods (see for example, Goding, 1983, *Monoclonal antibodies: principles and practice*, Academic Press Inc., New  
10 York; and Yokoyama, 1992, "Production of Monoclonal Antibodies" in *Current Protocols in Immunology*, Unit 2.5, Greene Publishing Assoc. and John Wiley & Sons). Polyclonal sera and antibodies can be produced by inoculation of a mammalian subject with the relevant protein or fragments thereof in accordance with known methods. Fragments of antibodies, receptors, or other reactive peptides can be produced from the corresponding  
15 antibodies by cleavage of and collection of the desired fragments in accordance with known methods (see for example, Goding, *supra*; and Andrew et al., 1992, "Fragmentation of Immunoglobulins" in *Current Protocols in Immunology*, Unit 2.8, Greene Publishing Assoc. and John Wiley & Sons). Chimeric antibodies and single chain antibodies can also be produced in accordance with known recombinant methods (see for example, 5,169,939,  
20 5,194,594, and 5,576,184). Humanized antibodies can also be made from corresponding murine antibodies in accordance with well known methods (see for example, U.S. Patent Nos. 5,530,101, 5,585,089, and 5,693,762). Additionally, human antibodies may be produced in non-human animals such as mice that have been genetically altered to express human antibody molecules (see for example Fishwild *et al.*, 1996, *Nature Biotechnology*  
25 14: 845-851; Mendez *et al.*, 1997, *Nature Genetics* 15: 146-156 (erratum *Nature Genetics* 16: 410); and U.S. Patents 5,877,397 and 5,625,126). Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for  
30 synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, J. Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, *et al.*, FEBS Lett. 211, 10 (1987).

Monoclonal antibodies binding to the protein of the invention may be useful diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where  
5 abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein.

For compositions of the present invention which are useful for bone, cartilage,  
10 tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or  
15 tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition  
20 would include a matrix capable of delivering the protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability,  
25 mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined,  
30 such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics.

Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

5           Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions from disassociating from the matrix.

10           A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic  
15 acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt% based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are  
20 prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells.

In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth factor  
25 (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins of the present invention.

30           The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be

formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (e.g., bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA).

Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

Patent and literature references cited herein are incorporated by reference as if fully set forth.

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:1;
- (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 27 to nucleotide 260;
- (c) the nucleotide sequence of SEQ ID NO:1 from nucleotide 72 to nucleotide 260;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc62\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc62\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:1.

2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.
3. A host cell transformed with the polynucleotide of claim 2.
4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:
  - (a) growing a culture of a host cell in a suitable culture medium, wherein the host cell has been transformed with the polynucleotide of claim 2; and
  - (b) purifying said protein from the culture.
6. A protein produced according to the process of claim 5.
7. An isolated polynucleotide encoding the protein of claim 6.
8. The polynucleotide of claim 7, wherein the polynucleotide comprises the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114.
9. A protein comprising an amino acid sequence selected from the group consisting of:
  - (a) the amino acid sequence of SEQ ID NO:2;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc62\_1 deposited with the ATCC under accession number 207114;the protein being substantially free from other mammalian proteins.
10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:2.

11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.

12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:3;
- (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 6 to nucleotide 1325;
- (c) the nucleotide sequence of SEQ ID NO:3 from nucleotide 99 to nucleotide 1325;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp10\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp10\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:3.

13. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp10\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

14. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:5;
- (b) the nucleotide sequence of SEQ ID NO:5 from nucleotide 149 to nucleotide 322;
- (c) the nucleotide sequence of SEQ ID NO:5 from nucleotide 200 to nucleotide 322;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp11\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp11\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;



(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:5.

15. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:6;

(b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp11\_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

16. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:7;

(b) the nucleotide sequence of SEQ ID NO:7 from nucleotide 288 to nucleotide 629;

(c) the nucleotide sequence of SEQ ID NO:7 from nucleotide 363 to nucleotide 629;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vp13\_1 deposited with the ATCC under accession number 207114;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114;

(f) the nucleotide sequence of a mature protein coding sequence of clone vp13\_1 deposited with the ATCC under accession number 207114;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:8;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:7.

17. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:8;

(b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp13\_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

18. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:9;

(b) the nucleotide sequence of SEQ ID NO:9 from nucleotide 11 to nucleotide 298;

(c) the nucleotide sequence of SEQ ID NO:9 from nucleotide 149 to nucleotide 298;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp16\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp16\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:9.

19. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp16\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

20. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:11;
- (b) the nucleotide sequence of SEQ ID NO:11 from nucleotide 257 to nucleotide 607;
- (c) the nucleotide sequence of SEQ ID NO:11 from nucleotide 479 to nucleotide 607;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp21\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp21\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:11.

21. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp21\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

22. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:13;
- (b) the nucleotide sequence of SEQ ID NO:13 from nucleotide 163 to nucleotide 477;
- (c) the nucleotide sequence of SEQ ID NO:13 from nucleotide 238 to nucleotide 477;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp22\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp22\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:14;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:13.

23. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:14;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp22\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

24. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:15;
- (b) the nucleotide sequence of SEQ ID NO:15 from nucleotide 58 to nucleotide 624;
- (c) the nucleotide sequence of SEQ ID NO:15 from nucleotide 106 to nucleotide 624;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq2\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq2\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:16;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:15.

25. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:16;

(b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq2\_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

26. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:17;

(b) the nucleotide sequence of SEQ ID NO:17 from nucleotide 773 to nucleotide 1090;

(c) the nucleotide sequence of SEQ ID NO:17 from nucleotide 842 to nucleotide 1090;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq3\_1 deposited with the ATCC under accession number 207114;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq3\_1 deposited with the ATCC under accession number 207114;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:18;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:17.

27. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:18;

(b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq3\_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

28. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:



- (a) the nucleotide sequence of SEQ ID NO:19;
- (b) the nucleotide sequence of SEQ ID NO:19 from nucleotide 96 to nucleotide 275;
- (c) the nucleotide sequence of SEQ ID NO:19 from nucleotide 159 to nucleotide 275;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq5\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq5\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:19.

29. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;

- (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq5\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

30. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:21;
- (b) the nucleotide sequence of SEQ ID NO:21 from nucleotide 176 to nucleotide 340;
- (c) the nucleotide sequence of SEQ ID NO:21 from nucleotide 230 to nucleotide 340;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq6\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq6\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:21.

31. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq6\_1 deposited with the ATCC under accession number 207114;
- the protein being substantially free from other mammalian proteins.

32. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:23;
- (b) the nucleotide sequence of SEQ ID NO:23 from nucleotide 29 to nucleotide 1111;
- (c) the nucleotide sequence of SEQ ID NO:23 from nucleotide 167 to nucleotide 1111;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vr1\_1 deposited with the ATCC under accession number 207114;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number 207114;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vr1\_1 deposited with the ATCC under accession number 207114;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number 207114;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:24;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:23.

33. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:24;

(b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and

(c) the amino acid sequence encoded by the cDNA insert of clone vr1\_1 deposited with the ATCC under accession number 207114;

the protein being substantially free from other mammalian proteins.

34. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:25;

(b) the nucleotide sequence of SEQ ID NO:25 from nucleotide 13 to nucleotide 513;

(c) the nucleotide sequence of the full-length protein coding sequence of clone vc63\_1 deposited with the ATCC under accession number 207115;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:26;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:25.

35. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:26;

(b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc63\_1 deposited with the ATCC under accession number 207115;

the protein being substantially free from other mammalian proteins.

36. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:27;

(b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 79 to nucleotide 345;

(c) the nucleotide sequence of SEQ ID NO:27 from nucleotide 130 to nucleotide 345;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vb25\_1 deposited with the ATCC under accession number PTA-362;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362;

(f) the nucleotide sequence of a mature protein coding sequence of clone vb25\_1 deposited with the ATCC under accession number PTA-362;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:28;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:27.

37. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:28;

(b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb25\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

38. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:29;
- (b) the nucleotide sequence of SEQ ID NO:29 from nucleotide 72 to nucleotide 236;
- (c) the nucleotide sequence of SEQ ID NO:29 from nucleotide 150 to nucleotide 236;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:30;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:29.

39. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;

- (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb27\_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

40. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:31;
- (b) the nucleotide sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 884;
- (c) the nucleotide sequence of SEQ ID NO:31 from nucleotide 183 to nucleotide 884;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb28\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb28\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:32;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and



(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:31.

41. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:32;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb28\_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

42. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:33;
- (b) the nucleotide sequence of SEQ ID NO:33 from nucleotide 42 to nucleotide 206;
- (c) the nucleotide sequence of SEQ ID NO:33 from nucleotide 111 to nucleotide 206;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb29\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:34;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:33.

43. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:34;

(b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb29\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

44. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 17 to nucleotide 253;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 98 to nucleotide 253;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:36;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:35.

45. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:36;

(b) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb30\_1 deposited with the ATCC under accession number PTA-362; the protein being substantially free from other mammalian proteins.

46. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:37;
- (b) the nucleotide sequence of SEQ ID NO:37 from nucleotide 68 to nucleotide 424;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vc67\_1 deposited with the ATCC under accession number PTA-362;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc67\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:38;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:37.

47. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:38;
- (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc67\_1 deposited with the ATCC under accession number PTA-362; the protein being substantially free from other mammalian proteins.

48. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:39;
- (b) the nucleotide sequence of SEQ ID NO:39 from nucleotide 103 to nucleotide 261;
- (c) the nucleotide sequence of SEQ ID NO:39 from nucleotide 154 to nucleotide 261;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:40;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:39.

49. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vf4\_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

50. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:41;
- (b) the nucleotide sequence of SEQ ID NO:41 from nucleotide 1575 to nucleotide 3038;
- (c) the nucleotide sequence of SEQ ID NO:41 from nucleotide 1650 to nucleotide 3038;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:42;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:41.

51. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:42;

(b) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and

(c) the amino acid sequence encoded by the cDNA insert of clone vg3\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

52. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:43;

(b) the nucleotide sequence of SEQ ID NO:43 from nucleotide 2112 to nucleotide 2363;

(c) the nucleotide sequence of the full-length protein coding sequence of clone vo2\_1 deposited with the ATCC under accession number PTA-362;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo2\_1 deposited with the ATCC under accession number PTA-362;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:44;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:43.

53. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:44;

(b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo2\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

54. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:45;

(b) the nucleotide sequence of SEQ ID NO:45 from nucleotide 36 to nucleotide 707;

(c) the nucleotide sequence of SEQ ID NO:45 from nucleotide 393 to nucleotide 707;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo3\_1 deposited with the ATCC under accession number PTA-362;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;



- (f) the nucleotide sequence of a mature protein coding sequence of clone vo3\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:46;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:45.

55. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
- (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo3\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

56. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:47;
- (b) the nucleotide sequence of SEQ ID NO:47 from nucleotide 74 to nucleotide 295;

(c) the nucleotide sequence of SEQ ID NO:47 from nucleotide 134 to nucleotide 295;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo5\_1 deposited with the ATCC under accession number PTA-362;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo5\_1 deposited with the ATCC under accession number PTA-362;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:48;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:47.

57. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:48;

(b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo5\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

58. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:49;
- (b) the nucleotide sequence of SEQ ID NO:49 from nucleotide 45 to nucleotide 383;
- (c) the nucleotide sequence of SEQ ID NO:49 from nucleotide 312 to nucleotide 383;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:49.

59. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo6\_1 deposited with the ATCC under accession number PTA-362;
- the protein being substantially free from other mammalian proteins.

60. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:51;
- (b) the nucleotide sequence of SEQ ID NO:51 from nucleotide 186 to nucleotide 1739;
- (c) the nucleotide sequence of SEQ ID NO:51 from nucleotide 288 to nucleotide 1739;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:51.

61. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:52;

(b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo9\_1 deposited with the ATCC under accession number PTA-362;

the protein being substantially free from other mammalian proteins.

62. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:53;

(b) the nucleotide sequence of SEQ ID NO:53 from nucleotide 440 to nucleotide 835;

(c) the nucleotide sequence of SEQ ID NO:53 from nucleotide 632 to nucleotide 835;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo11\_1 deposited with the ATCC under accession number PTA-366;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo11\_1 deposited with the ATCC under accession number PTA-366;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo11\_1 deposited with the ATCC under accession number PTA-366;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo11\_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:54;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:53.

63. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:54;

(b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo11\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

64. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:55;

(b) the nucleotide sequence of SEQ ID NO:55 from nucleotide 72 to nucleotide 329;

(c) the nucleotide sequence of SEQ ID NO:55 from nucleotide 120 to nucleotide 329;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo12\_1 deposited with the ATCC under accession number PTA-366;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo12\_1 deposited with the ATCC under accession number PTA-366;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:56;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:55.

65. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:56;

(b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo12\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

66. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:57;
- (b) the nucleotide sequence of SEQ ID NO:57 from nucleotide 227 to nucleotide 439;
- (c) the nucleotide sequence of SEQ ID NO:57 from nucleotide 287 to nucleotide 439;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:58;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:57.

67. A protein comprising an amino acid sequence selected from the group consisting of:



- (a) the amino acid sequence of SEQ ID NO:58;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo13\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

68. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:59;
- (b) the nucleotide sequence of SEQ ID NO:59 from nucleotide 96 to nucleotide 341;
- (c) the nucleotide sequence of SEQ ID NO:59 from nucleotide 174 to nucleotide 341;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:59.

69. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo14\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

70. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:61;
- (b) the nucleotide sequence of SEQ ID NO:61 from nucleotide 90 to nucleotide 599;
- (c) the nucleotide sequence of SEQ ID NO:61 from nucleotide 165 to nucleotide 599;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo15\_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo15\_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:62;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:61.

71. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:62;

(b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo15\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

72. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:63;

(b) the nucleotide sequence of SEQ ID NO:63 from nucleotide 209 to nucleotide 451;

(c) the nucleotide sequence of SEQ ID NO:63 from nucleotide 398 to nucleotide 451;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo16\_1 deposited with the ATCC under accession number PTA-366;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo16\_1 deposited with the ATCC under accession number PTA-366;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:64;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:63.

73. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:64;

(b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo16\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

74. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:65;
- (b) the nucleotide sequence of SEQ ID NO:65 from nucleotide 31 to nucleotide 231;
- (c) the nucleotide sequence of SEQ ID NO:65 from nucleotide 97 to nucleotide 231;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:65.

75. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;

(b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo18\_1 deposited with the ATCC under accession number PTA-366; the protein being substantially free from other mammalian proteins.

76. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:67;

(b) the nucleotide sequence of SEQ ID NO:67 from nucleotide 23 to nucleotide 736;

(c) the nucleotide sequence of SEQ ID NO:67 from nucleotide 83 to nucleotide 736;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo19\_1 deposited with the ATCC under accession number PTA-366;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-366;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo19\_1 deposited with the ATCC under accession number PTA-366;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:68;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:67.

77. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo19\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

78. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:69;
- (b) the nucleotide sequence of SEQ ID NO:69 from nucleotide 104 to nucleotide 1399;
- (c) the nucleotide sequence of SEQ ID NO:69 from nucleotide 158 to nucleotide 1399;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo22\_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo22\_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo22\_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo22\_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:70;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:69.

79. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:70;

(b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo22\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

80. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:71;

(b) the nucleotide sequence of SEQ ID NO:71 from nucleotide 174 to nucleotide 1595;

(c) the nucleotide sequence of the full-length protein coding sequence of clone vo23\_1 deposited with the ATCC under accession number PTA-366;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-366;



(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:72;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:71.

81. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:72;

(b) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo23\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

82. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:73;

(b) the nucleotide sequence of SEQ ID NO:73 from nucleotide 129 to nucleotide 311;

(c) the nucleotide sequence of SEQ ID NO:73 from nucleotide 195 to nucleotide 311;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:74;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:73.

83. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:74;

(b) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo24\_1 deposited with the ATCC under accession number PTA-366;

the protein being substantially free from other mammalian proteins.

84. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:75;
- (b) the nucleotide sequence of SEQ ID NO:75 from nucleotide 73 to nucleotide 798;
- (c) the nucleotide sequence of SEQ ID NO:75 from nucleotide 142 to nucleotide 798;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:75.

85. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;

- (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo25\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

86. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:27;
- (b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 26 to nucleotide 307;
- (c) the nucleotide sequence of SEQ ID NO:27 from nucleotide 101 to nucleotide 307;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:27.

87. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo26\_1 deposited with the ATCC under accession number PTA-366;
- the protein being substantially free from other mammalian proteins.

88. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:79;
- (b) the nucleotide sequence of SEQ ID NO:79 from nucleotide 43 to nucleotide 228;
- (c) the nucleotide sequence of SEQ ID NO:79 from nucleotide 94 to nucleotide 228;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp23\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp23\_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp23\_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp23\_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:80;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:79.

89. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:80;

(b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp23\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

90. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:81;

(b) the nucleotide sequence of SEQ ID NO:81 from nucleotide 245 to nucleotide 427;

(c) the nucleotide sequence of SEQ ID NO:81 from nucleotide 308 to nucleotide 427;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq7\_1 deposited with the ATCC under accession number PTA-368;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq7\_1 deposited with the ATCC under accession number PTA-368;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:82;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:81.

91. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:82;

(b) a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq7\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

92. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:83;
- (b) the nucleotide sequence of SEQ ID NO:83 from nucleotide 119 to nucleotide 475;
- (c) the nucleotide sequence of SEQ ID NO:83 from nucleotide 185 to nucleotide 475;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:84;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:83.

93. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:84;



- (b) a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq8\_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

94. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:85;
- (b) the nucleotide sequence of SEQ ID NO:85 from nucleotide 90 to nucleotide 323;
- (c) the nucleotide sequence of SEQ ID NO:85 from nucleotide 141 to nucleotide 323;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq9\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq9\_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:86;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:85.

95. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:86;
- (b) a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vq9\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

96. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:87;
- (b) the nucleotide sequence of SEQ ID NO:87 from nucleotide 18 to nucleotide 452;
- (c) the nucleotide sequence of SEQ ID NO:87 from nucleotide 72 to nucleotide 452;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq10\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq10\_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:88;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:87.

97. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:88;

(b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq10\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

98. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:89;

(b) the nucleotide sequence of SEQ ID NO:89 from nucleotide 196 to nucleotide 378;

(c) the nucleotide sequence of SEQ ID NO:89 from nucleotide 262 to nucleotide 378;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:90;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:89.

99. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:90;

(b) a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq13\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

100. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:91;
- (b) the nucleotide sequence of SEQ ID NO:91 from nucleotide 35 to nucleotide 718;
- (c) the nucleotide sequence of SEQ ID NO:91 from nucleotide 173 to nucleotide 718;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:92;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:91.

101. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:92;

- (b) a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq16\_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

102. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:93;
- (b) the nucleotide sequence of SEQ ID NO:93 from nucleotide 1 to nucleotide 762;
- (c) the nucleotide sequence of SEQ ID NO:93 from nucleotide 70 to nucleotide 762;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:94;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:93.

103. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vq19\_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

104. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:95;
- (b) the nucleotide sequence of SEQ ID NO:95 from nucleotide 106 to nucleotide 792;
- (c) the nucleotide sequence of SEQ ID NO:95 from nucleotide 172 to nucleotide 792;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq20\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq20\_1 deposited with the ATCC under accession number PTA-368;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:96;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:95.

105. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:96;

(b) a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq20\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

106. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:97;

(b) the nucleotide sequence of SEQ ID NO:97 from nucleotide 40 to nucleotide 315;

(c) the nucleotide sequence of SEQ ID NO:97 from nucleotide 124 to nucleotide 315;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq21\_1 deposited with the ATCC under accession number PTA-368;



(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq21\_1 deposited with the ATCC under accession number PTA-368;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:98;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:97.

107. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:98;

(b) a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq21\_1 deposited with the ATCC under accession number PTA-368;

the protein being substantially free from other mammalian proteins.

108. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:99;
- (b) the nucleotide sequence of SEQ ID NO:99 from nucleotide 70 to nucleotide 699;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vr2\_1 deposited with the ATCC under accession number PTA-368;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vr2\_1 deposited with the ATCC under accession number PTA-368;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:100;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:99.

109. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vr2\_1 deposited with the ATCC under accession number PTA-368;
- the protein being substantially free from other mammalian proteins.

110. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:101;
- (b) the nucleotide sequence of SEQ ID NO:101 from nucleotide 170 to nucleotide 394;
- (c) the nucleotide sequence of SEQ ID NO:101 from nucleotide 227 to nucleotide 394;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone PTA-1075 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:101.

111. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;

- (b) a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc69\_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins.

112. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:103;
- (b) the nucleotide sequence of SEQ ID NO:103 from nucleotide 43 to nucleotide 198;
- (c) the nucleotide sequence of SEQ ID NO:103 from nucleotide 85 to nucleotide 198;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:104;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:103.

113. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc71\_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins.

114. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:105;
- (b) the nucleotide sequence of SEQ ID NO:105 from nucleotide 260 to nucleotide 1552;
- (c) the nucleotide sequence of SEQ ID NO:105 from nucleotide 335 to nucleotide 1552;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:106;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:105.

115. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:106;

(b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo27\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins.

116. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:107;

(b) the nucleotide sequence of SEQ ID NO:107 from nucleotide 15 to nucleotide 320;

(c) the nucleotide sequence of SEQ ID NO:107 from nucleotide 72 to nucleotide 320;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;

(f) the nucleotide sequence of a mature protein coding sequence of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:108;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:107.

117. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:108;

(b) a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108; and

(c) the amino acid sequence encoded by the cDNA insert of clone vo31\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins.

118. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:109;
- (b) the nucleotide sequence of SEQ ID NO:109 from nucleotide 38 to nucleotide 1255;
- (c) the nucleotide sequence of SEQ ID NO:109 from nucleotide 86 to nucleotide 1255;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:110;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:109.

119. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:110;



- (b) a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vo32\_1 deposited with the ATCC under accession number PTA-1075;
- the protein being substantially free from other mammalian proteins.

120. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:111;
- (b) the nucleotide sequence of SEQ ID NO:111 from nucleotide 80 to nucleotide 1276;
- (c) the nucleotide sequence of SEQ ID NO:111 from nucleotide 131 to nucleotide 1276;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:112;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:111.

121. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:112;
- (b) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vo33\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins.

122. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:113;
- (b) the nucleotide sequence of SEQ ID NO:113 from nucleotide 202 to nucleotide 429;
- (c) the nucleotide sequence of SEQ ID NO:113 from nucleotide 292 to nucleotide 429;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:114;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:113.

123. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:114;

(b) a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq23\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins.

124. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:115;

(b) the nucleotide sequence of SEQ ID NO:115 from nucleotide 37 to nucleotide 1113;

(c) the nucleotide sequence of SEQ ID NO:115 from nucleotide 88 to nucleotide 1113;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;

(f) the nucleotide sequence of a mature protein coding sequence of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:116;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:115.

125. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:116;

(b) a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq24\_1 deposited with the ATCC under accession number PTA-1075;

the protein being substantially free from other mammalian proteins.

126. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:117;
- (b) the nucleotide sequence of SEQ ID NO:117 from nucleotide 40 to nucleotide 207;
- (c) the nucleotide sequence of SEQ ID NO:117 from nucleotide 103 to nucleotide 207;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq26\_1 deposited with the ATCC under accession number PTA-1075;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:118;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:117.

127. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:118;

(b) a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118; and

(c) the amino acid sequence encoded by the cDNA insert of clone vq26\_1 deposited with the ATCC under accession number PTA-1075; the protein being substantially free from other mammalian proteins.

Fig. 1A





Fig. 1B





## SEQUENCE LISTING

<110> Valenzuela, Dario  
 Yuan, Olive  
 Hoffman, Heidi  
 Hall, Jeff  
 Rapiejko, Peter

<120> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

<130> GI 6919X

<140>

<141>

<160> 194

<170> PatentIn Ver. 2.0

<210> 1

<211> 4177

<212> DNA

<213> Homo sapiens

<400> 1

```

gttaaagggt aaccccatgc cttggaatgg tattaaccct gtggtgcaat ttgtgctcca 60
gagcttcctc atgggtgaga cagaagcacg tgtcctgttg tgtccacaac tacactcagc 120
cttttttggc cattcaatcc tcattttggg ctatgtcttc agaaactaaa ccgaaggcac 180
tatccaaaaga ttacttgtgt atttcttata gaagccccc cccaccccc acacacagac 240
attcctcaaa taggttccta tgacctacca gttgaagccc aggtctctta tctggacatt 300
aagtctttgc atggtcagtc ggtctctgt ctttcaagat ttatttttca ctatcataca 360
ccatattaac ctactcaggg ctgcctactt tacattctct cccagccctg cacattcctt 420
ggaaaggcct cccctgcctc gcctgaatga tgaacactct attccttatt caagactcta 480
acatcacttg cctgtggaag gtcctcaga ttcctctagg ctccagtagg gccactcctc 540
agtcccttac agcatatggt gtcaatgaca acagtgatat gcttgcttga ttcctccact 600
taactatgaa ttttttgttg ttgttaataa aaacgtttat ttgtcatttt atacagaacc 660
gcctgaagtc tgcacatga cttgatggtt acccaggggt ttatgtgtgt ccaactgaac 720
accgtgggtc tgggattgag gatccgcaca aacacaggca ggagtttggg ggggaagaagg 780
gggcacagca ggaagtgtat tcaagccac ctcaaaggca gcagagcatt ccaagttgct 840
gaaaacctgt gcaaatgggg cctctaaaac aaacattgta ctgtgagctc tcggggaagg 900
ggcaggaaaa ggaagtgtgat aattaagggg gcattttctt gaggcggcag atgaggaggg 960
aggcttacta tttttaatcc cctcagcccc aggacatgat aatcagagga gatcatgtcc 1020
aatcttgaat caattccctg ttcaaaaaga ggtgctaata ccccagggac cagactctga 1080
aaataccatg aaaatactct gaaaataccg tgctgggtcat tcctgagtc ctgttctatg 1140
ccctacaaca tattagaagg gcaagggaag gtcagagggt agtggcttgt gttctgtgga 1200
agtaaaacac cttagtccat attagcact gataaaaccc aatgactggg accacagatc 1260
tggtgagagg tcactagtag ggaccaagag tgggagggtc ggaacatcct cagcttgaga 1320
tgccagtagg gtggaagtc tgggaggagg taaatattaa agtaatggga aacagaacac 1380
tcccaagcct taattttaat tggatagttt aaaaaattga aatagaatct aacattgggt 1440
gccagtcacc acaaatgcca tgcctttatg gtcacttggg atcataaaaa atttgccaaa 1500
geatgcctgt ctggctaata aaaagggtctt gctaggagggt gttcgtgtgc atgacagagt 1560
gccaaagggc ttgggtaccac tgactatcca acgtgattcc tatggaaaca gaagcggcag 1620
agtccttgggt agctggcttg ttgaggtttg gcagagaaga agcgaagct ccaaagtgc 1680
tcagattct ctgcaactgg ctttgaccga gagaagagg agccacattc tactctaaa 1740
gatagagagg gaccaaact aaccacagc acatgcatta gttctgttca tctacagt 1800
gatgggcaac taactctggg caggggaagg gagctgggta ggggaagaagg gctcattttt 1860
cattttttta agtcttaatt aatattttta cctcaaacag attatcagtg cctcagatac 1920
aatttaactt taagtcttta aaaggcccc tgaaacaaaa atatacattt ttttaaggca 1980
tcctcacttt ctgatagtc gtttccccc gtctccagtt ttacctgac ttagagtcca 2040
caaacttcat cagaccacct gtgctcatgg acgtgggtct ttctagaggg aagcctcggg 2100

```

```

ctcgggtcca gatgagctgt gccagtacac ccaatgcctg tgatgcccc aacaggaccg 2160
cgtagcaatg tatctctgtc atgctatagt acccaccatg tgcgtctaca ttgggcccag 2220
gattctcggc cttaccctac tgtcagagga tattgggcac aatctctgtat agctgagcaa 2280
ccaacttaaa catgggatca ttatgcaatg ttccagagca aactctcgct gacagggtaca 2340
ttgtggctca gtcttcttta gtactgcatg gccatagcct ggaacaaccc atctcgagtt 2400
cagtgtagtc ttgtaacttc tcatctgaca catctttgcc aacttccttc tgtagttgtg 2460
ttagccagac aagcacttcc tgatttgcca gtccatgtag aggaactgcc agccattcca 2520
tggctgctgc aaagggcagg gaagggctctg aaaggggtact gcctgccaaa tgcctggcat 2580
gagcacttat attgctgtgc tcatggtcac tgtggatggt gagctacagg tgcctgagct 2640
cagtgaactg agcatcagta tagcctaaca tgttgggtgaa actgtgggac cagtccagct 2700
tagagtcaag gaccccaata ccactgccct ctccacagag attttggtag accttttctg 2760
caatacaagg tagctttgtg attagacaca catgggtcttc ataaatcaac tcccagtact 2820
tggtttgact ggacccctct ggatatgtc gggcaaagtt gtttagggct gccatggctg 2880
cagtgggctc agcatgggt gtggattggt gggaaagttg tccagcatgg tgaccacatg 2940
ggaaggtaga gctgcctgcc ctctttgcc actctcttaa aagccaagat acctgttctc 3000
ctcttgggat ctgtccagtt accagcagcc aaaataagcc cctgggtagg ggttttcccc 3060
acccttagtc ttgggtagca gttttggtat tcagggatac tgaagcctcg gaaacagata 3120
acctcatcag gatcaagaac tgatgtttat ataccaatcc cttaatgcct ctcatgccgc 3180
catacattat gtctgcagtg atttggccca ccaccttctt gccatgttgc tgcctgaagg 3240
tcttaattct ggctgtctcc ttaactatca ggtcagccaa tatgtcttcc aaattcttgg 3300
aggaagcact ggcatgccag gcagcaagaa caagacaaga tgcactgttg cttccaaga 3360
gcctggcagc cacttgggta aaacatgggt gagtgaactc tgcgactctg tggagagggg 3420
agaaggggag agagctgtgg cagaacacag aaccaccact gcggcaccag aggccgcaat 3480
gggtggacaa ggttgaagg aggcagctga gaaacacat agatgaaccc gcagcaggtg 3540
cccagccagg ggttctcagc aaggccctgt gcactgccaa ctatgaactt attaagactg 3600
actttccaat agttcttagc agtgtgtgga gtagaagaca gaacaacgta taccttcaga 3660
atgcacagag ccaatgcatt tttggattac aaataaatca atttaagacc tattgttctc 3720
gacaatgatt tgcaaatga gaaaattctg gaaaaataga catccaacaa attaacagtg 3780
attataccta atgacagggt tacaggtggg gatttgttta tttgtatgtt cttatttgtt 3840
tcttgtttct tatgtcagca aattttaaat agaatatgca aaatgatgac ctcaataata 3900
aaaaatattg tgtttattca caagagttaa ataattgtta tagaaaggaa aagatgcgaa 3960
tggttttggg cagattagga tttaaacaaa aatgttaaac acagaataca aaatatgttt 4020
tttctcctgt tgtctagctt ttgtacagtg aacacatggt gcttttatag taagaaatat 4080
acattttaag tttcaaaatt actcagattt tgtaaacat ctaaaagaaa ataatgtgt 4140
ttggaataac caaaaaaaaa aaaaaaaaaa aaaaaaa 4177

```

&lt;210&gt; 2

&lt;211&gt; 78

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```

Met Val Leu Thr Leu Trp Cys Asn Leu Cys Ser Arg Ala Ser Ser Trp
  1             5             10             15

```

```

Val Arg Gln Lys His Val Ser Cys Cys Val His Asn Tyr Thr Gln Pro
      20             25             30

```

```

Phe Leu Leu Ile Gln Ser Ser Phe Trp Ala Met Ser Ser Glu Thr Lys
      35             40             45

```

```

Pro Lys Ala Leu Ser Lys Asp Tyr Leu Cys Ile Ser Tyr Arg Ser Pro
      50             55             60

```

```

His Ser Thr Pro Thr His Arg His Ser Ser Asn Arg Phe Leu
      65             70             75

```

&lt;210&gt; 3

&lt;211&gt; 1326

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

```

gggagatggc agtgagcag aggagggggc tcggccgcgg gagccccgcg gagtgggggc 60
agcggctact tctggtgctg ctgttgggtg gctgctccgg gcgcatccac cggttggcgc 120
tgacggggga gaagcgagcg gacatccagc tgaacagctt cggtttctac accaatggct 180
ctctggagggt ggagttgagc gtccctgcggc tgggcctccg ggaggcagaa gagaagtccc 240
tgctgggtggg gttcagtctc agccgggttc ggtctggcag agttcgctcc tattcaaccc 300
gggatttcca ggactgccct ctccagaaaa acagtagcag tttcctggtc ctgttcctca 360
tcaacaccaa ggatctgcag gtccagggtgc ggaagtatgg agagcagaag acgttggtta 420
tctttcccg gtcctcccg gaagcaccct ccaaaccagg gctcccgaag ccacaggcca 480
cagtcccccg caaggtggat ggcggaggga cctctgcagc cagcaagccc aagtcaacac 540
ccgcagtgat tcagggtcct agtgggaagg acaaggacct ggtgttgggc ctgagccacc 600
tcaacaactc ctacaacttc agtttccacg tggatgcgg ctctcaggcg gaagaaggcc 660
agtacagcct gaacttcac aactgcaaca attcagtgc aggaaaggag catccattcg 720
acatcacggt gatgatccgg gagaagaacc ccgatggctt cctgtcggca gcggagatgc 780
cccttttcaa gctctacatg gtcattgctc cctgttccct ggccgctggc agtggctgta 840
ccagctcttg gtggagggtc ccaccctgcc ttcttcgtgc tcacgggcta caagtccag 900
cccacaggaa acaaccgta cctgcagctg cccaggagg acgaggagga tgttcagatg 960
gagcaagtaa tgacggactc tgggttcccg gaaggcctct ccaaagtcaa caaaacagcc 1020
agcgggcggg aactgttatg atcacctca catctcagac caaagggtcg tcctcccca 1080
gcatttttca ctctgcctt tcttccacag cgtatgtggg gaggtggagg ggttccatgt 1140
ggaccaggcg cccagctccc cgggaccccg gttcccgac aagccattt ggaagaagag 1200
tcccttctc ccccaataa ttgggcagcc ctgtccttac cccgggacca cccctccctt 1260
ccagctatgt gtacaataa gaccaatctg tttggctaaa aaaaaaaaaa aaaaaaaaaa 1320
aaaaaa

```

&lt;210&gt; 4

&lt;211&gt; 440

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

```

Met Ala Val Ser Glu Arg Arg Gly Leu Gly Arg Gly Ser Pro Ala Glu
 1             5             10            15

Trp Gly Gln Arg Leu Leu Leu Val Leu Leu Leu Gly Gly Cys Ser Gly
      20             25             30

Arg Ile His Arg Leu Ala Leu Thr Gly Glu Lys Arg Ala Asp Ile Gln
      35             40             45

Leu Asn Ser Phe Gly Phe Tyr Thr Asn Gly Ser Leu Glu Val Glu Leu
      50             55             60

Ser Val Leu Arg Leu Gly Leu Arg Glu Ala Glu Lys Ser Leu Leu
      65             70             75            80

Val Gly Phe Ser Leu Ser Arg Val Arg Ser Gly Arg Val Arg Ser Tyr
      85             90            95

Ser Thr Arg Asp Phe Gln Asp Cys Pro Leu Gln Lys Asn Ser Ser Ser
      100            105            110

Phe Leu Val Leu Phe Leu Ile Asn Thr Lys Asp Leu Gln Val Gln Val
      115            120            125

Arg Lys Tyr Gly Glu Gln Lys Thr Leu Phe Ile Phe Pro Gly Leu Leu
      130            135            140

```

Pro Glu Ala Pro Ser Lys Pro Gly Leu Pro Lys Pro Gln Ala Thr Val  
 145 150 155 160  
 Pro Arg Lys Val Asp Gly Gly Gly Thr Ser Ala Ala Ser Lys Pro Lys  
 165 170 175  
 Ser Thr Pro Ala Val Ile Gln Gly Pro Ser Gly Lys Asp Lys Asp Leu  
 180 185 190  
 Val Leu Gly Leu Ser His Leu Asn Asn Ser Tyr Asn Phe Ser Phe His  
 195 200 205  
 Val Val Ile Gly Ser Gln Ala Glu Glu Gly Gln Tyr Ser Leu Asn Phe  
 210 215 220  
 His Asn Cys Asn Asn Ser Val Pro Gly Lys Glu His Pro Phe Asp Ile  
 225 230 235 240  
 Thr Val Met Ile Arg Glu Lys Asn Pro Asp Gly Phe Leu Ser Ala Ala  
 245 250 255  
 Glu Met Pro Leu Phe Lys Leu Tyr Met Val Met Ser Ala Cys Phe Leu  
 260 265 270  
 Ala Ala Gly Ser Gly Cys Thr Ser Ser Trp Trp Arg Ala Pro Pro Cys  
 275 280 285  
 Leu Leu Arg Ala His Gly Leu Gln Val Pro Ala His Arg Lys Gln Pro  
 290 295 300  
 Val Pro Ala Ala Ala Pro Gly Gly Arg Gly Gly Cys Ser Asp Gly Ala  
 305 310 315 320  
 Ser Asn Asp Gly Leu Trp Val Pro Gly Arg Pro Leu Gln Ser Gln Gln  
 325 330 335  
 Asn Ser Gln Arg Ala Gly Thr Val Met Ile Thr Ser Thr Ser Gln Thr  
 340 345 350  
 Lys Gly Ser Ser Ser Pro Ser Ile Ser His Ser Cys Pro Ser Ser Thr  
 355 360 365  
 Ala Tyr Val Gly Arg Trp Arg Gly Ser Met Trp Thr Arg Arg Pro Ala  
 370 375 380  
 Pro Arg Asp Pro Gly Ser Arg Thr Ser Pro Phe Gly Arg Arg Val Pro  
 385 390 395 400  
 Ser Ser Pro Gln Ile Leu Gly Ser Pro Val Leu Thr Pro Gly Pro Pro  
 405 410 415  
 Leu Pro Ser Ser Tyr Val Tyr Asn Asn Asp Gln Ser Val Trp Leu Lys  
 420 425 430  
 Lys Lys Lys Lys Lys Lys Lys  
 435 440

&lt;210&gt; 5

<211> 1280  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 aaaggggcta attggagatc tgtgagttta atgagcttga gtctcttctt cacaccagag 60  
 actgatgttt tctgggttagc aggggagcca gtaccttctt tttctcagta agtatctttc 120  
 tagtcacctg aggatatttg gagacagaat ggagaatgaa ttgattgcct tgggtgcctc 180  
 attcagggat gtgacagcag ctagtacaca ctgcatttgt gataaggctg atacgaggga 240  
 caacagaggg tacgttttct tgaagatgta tcttgtaaaa gaaaaaaatc ctttatttat 300  
 atatgatgga aagcttgga catagcacca ctccctttag aacttggtg tgaaaatgag 360  
 caattagtgg acatttagat ggtgcttaaa tgagcaagt ttcttgatta tagcctaaca 420  
 gagaaatcag aaagtgacca atataacatg ttttgttgac aaactaaaag catacacatt 480  
 agcagcatat ctctgtagaa ccttcacttc atgcttgact atcatgtatt ttgagtttgc 540  
 aaagaaatag ctggacagtc cagaagctat ctggctcaga catccattta tctcaggctg 600  
 tgtagagtgc tacacaggaa gaccacagca gcagcatcac ctggaacttc atagaaatgc 660  
 agattcttgg gctccattgc agacctaaaga aatcagaatc tgcattttta taagaccccc 720  
 ccaaagcaat ttgagtgcac attaaagctt aagaaacact ggttgggtcta ggccgggctg 780  
 ggtggcttat gcctgtgac ccagcacttt gggaggccaa ggcgggcaga tcacgaggtc 840  
 aggagattga gaacatcctg gctagcatgg tgaaccccg tctctgctaa aaatacaaaa 900  
 aattagccag gcgtgggtggc gggcgctgt agtcccagct acttgggagg ctgaggcagc 960  
 agaatggcat gaacccggga ggcagagctt tcggtgagcc cagatggcgc cactgcactc 1020  
 cagcctgggc tacagagccc gagactctgt ctcaaaaaac aaaaaaaaca aaaaaaaaca 1080  
 aaaaaaaat tagctgggccc tgatggcacg tgctgtagt cccagctact caggaggctg 1140  
 agggcaagaga attgcttgaa cctgggaggt ggaggttgca gtgagccgag attgcaccac 1200  
 tgcattctag cctggcgaca gagcgagact tcgtctcaa acaaaaacaa gcaaacaaaa 1260  
 aaaaaaaaaa aaaaaaaaaa 1280

<210> 6  
 <211> 58  
 <212> PRT  
 <213> Homo sapiens

<400> 6  
 Met Glu Asn Glu Leu Ile Ala Leu Gly Ala Ser Phe Arg Asp Val Thr  
 1 5 10 15  
 Ala Ala Ser Thr His Cys Ile Cys Asp Lys Ala Asp Thr Arg Asp Asn  
 20 25 30  
 Arg Gly Tyr Val Phe Leu Lys Met Tyr Leu Val Lys Glu Lys Asn Pro  
 35 40 45  
 Leu Phe Ile Tyr Asp Gly Lys Leu Gly Thr  
 50 55

<210> 7  
 <211> 1001  
 <212> DNA  
 <213> Homo sapiens

<400> 7  
 aggattgggc ttcaatgtga ctaacctaca attgcctcca ggtgctccac cactgagtc 60  
 cttgtgtctc tctgagggtc cttggagagt tactggagag ggctctgtgt cagatgacct 120  
 tgaggaggct ctgatttagc cttttgtaaa atgcaaagag ttgaggtctt ctccacgcaa 180  
 gagctcgctg atgtcaatga ggtattgagg atggggccat ctccatttct tgtggccagt 240  
 actgagtttt gttatccttc ctttaggtaa ggagtgccaa tggaccgatg cctgcctgtc 300  
 tcatccctgt gcaaatggaa gtacctgtac cactgtggcc aaccagttct cctgcaaatg 360  
 cctcacaggc ttcacagggc agaagtgtga gactgatgtc aatgagtgtg acattccagg 420

acactgccag catggtggca cctgcctcaa cctgcctggt tctaccagt gccagtgcct 480  
 tcagggtctc acaggccagt actgtgacag cctgtatgtg cctgtgcac cctgccttg 540  
 tgtaaatgga ggcacctgtc ggcagactgg tgacttcact tttgagtgc actgccttcc 600  
 aggttaaggag ctccctagtg tcccaggatt aggggacaaa cccctagcac aggaggtagt 660  
 ggggtgtggct caattgtttt ttttaggaag cgcaaggaaa aagggaagtg agaattttgt 720  
 ggggggtggg ttgctagtga gggaggagtt ttatggggcc actgtggtcc ataaactgag 780  
 caggggataa tttagcatgt cagggtttat gatgatgagt ggctagaaaa ttgtttattg 840  
 tcccttttgt agaaacagtg agaagaggaa cagagctctg ggaaagagac agggaagtct 900  
 ggaatggaaa agaacacgat gagaattaga cactggaaaa tatgtatgtg tggttaataa 960  
 agtgccttaa actgaaaaaa aaaaaaaaaa aaaaaaaaaa a 1001

<210> 8

<211> 114

<212> PRT

<213> Homo sapiens

<400> 8

Met Pro Ala Cys Leu Ile Pro Val Gln Met Glu Val Pro Val Pro Leu  
 1 5 10 15

Trp Pro Thr Ser Ser Pro Ala Asn Ala Ser Gln Ala Ser Gln Gly Arg  
 20 25 30

Ser Val Arg Leu Met Ser Met Ser Val Thr Phe Gln Asp Thr Ala Ser  
 35 40 45

Met Val Ala Pro Ala Ser Thr Cys Leu Val Pro Thr Ser Ala Ser Ala  
 50 55 60

Phe Arg Ala Ser Gln Ala Ser Thr Val Thr Ala Cys Met Cys Pro Val  
 65 70 75 80

His Pro Arg Leu Val Ser Met Glu Ala Pro Val Gly Arg Leu Val Thr  
 85 90 95

Ser Leu Leu Ser Ala Thr Ala Phe Gln Val Arg Ser Ser Leu Val Ser  
 100 105 110

Gln Asp

<210> 9

<211> 2058

<212> DNA

<213> Homo sapiens

<400> 9

ttttttatat atgacctttg atttctgttg tttgtatttt agcacagtgt atgcaccttc 60  
 atttaaatac atctgtgtgc atacagatac gcatatatgt gtgtgcgtat gcatatatct 120  
 ctcatctgta gtttccaaga gttcagctga agcagatgga gtccctgcagc ccaggagaca 180  
 ccctgcaccc ctgctaatag tgtttgccac aagtattagt gagtcttcct tattaatatt 240  
 ttcatttcag aagactgaag caaagctgat agtgtttgct gtttcttttg cagctaagtg 300  
 agggctcttg gatgacttgc tgtgttcctc aagctgcact ttggggccat ctctgcagta 360  
 ttagccccct ttttgcttg tggtactctg tctgtgcctg tgtgtgtgtg tgatagtcac 420  
 tcttgcatgg ctcccatgtc tggtttgttg catttgggga taaggtgctg aagccagagc 480  
 atttgcagtt tgtttgaggc ctctgtgcca atgatagatc actcctgttg acctggtatg 540  
 tctgcttgc tgcgtctttt ccttgctttc tcttggaaga ggaaaggact ctggtcaggc 600  
 ccaggctgag tgagatgagc tgcagctggc tcatggcctt cttagagcag agagaggagt 660  
 atgtcatttt actaagttcc taaacaaaca tttatgcagg caacactcct tgcagatcca 720

```

gaaactgagg cacaataggg ttatgacttg ctcaagaata thtagctgct aggggggtaaa 780
tcaaggcatc acaatttctg ttcagcgggc aggaataggg tgtgaattgc tagcactttt 840
tttttttaag caattacttt ttgacttggt cctctgaaag tgcaagaggg gtacaccttt 900
cccaaatgta gactagaatc tgcaggatgc caccactgt atagttctgc tttcccagag 960
aggaagaact tttagaaacc aaatgatctt aattgttatt gccacccct ggcttttccg 1020
ggtagaaaat tcacagtagg aatgattggt aagagagagt gcttggaaacc atgggttaac 1080
aggaaaggct acctaacttc acatatctgc aaccagagca gccaccaagc attacttagc 1140
agcaggaaaa tgattgtatt tgagttcctg tgtgtccaaa actgaggcac catgttcttt 1200
gaaaacatgc cactcaagg ctgggcggcg tggctcacac ctgtaatccc agcactttgg 1260
gaggccgagg cgggcggatc accggaggtc gggagtgtga gaccagcctg accaacaagg 1320
agaaacccca tctctactaa aaatacaaaa ttagccgggc gtggtggcat gcgcctataa 1380
tctcagctac ttgggaggct gaggcaggag aattgcttga acccaggagg cggagggtgc 1440
ggtgagtga gatcgtgcca ttgcactccg gcctgggcaa caacagcaaa actccgtctc 1500
aaaaaaaaa aaaaaaaaaa tgccacctca cagtctctat cgttggggcc tgtggttgca 1560
aagctgcctc ctgaattaag aaggaaaggg aaaagatata gctggaagcg agtgggaagag 1620
atgctgaggg ctggggctgc acagacgtgc tcagcagggc tacaggtgct caagccctat 1680
tggggatggg tggggagtgg ggcagcggcg tttgccacct taagaattgg ggccaaagcc 1740
actgatgttt atttgacagt gacactgcac tgggtactta aagaaattat ttcccgttgt 1800
aattataatt actgcttatt aaggaaaata tgggaatttt agaaagaatc aagtttgcca 1860
cccaaatgct accactgtta atcttttggg gttaaatgtt cccctagac atttctgtgc 1920
atagattttt ggtgtgttta catagtcgtt attctgtata tacaatttta tgtccctttt 1980
gtacttaacg tataaacatt tttccatgt taaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040
aaaaaaaaa aaaaaaaaaa                                     2058

```

&lt;210&gt; 10

&lt;211&gt; 96

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 10

```

Met Thr Phe Asp Phe Cys Cys Leu Tyr Phe Ser Thr Val Tyr Ala Pro
 1             5             10            15

Ser Phe Lys Tyr Ile Cys Val His Thr Asp Thr His Ile Cys Val Cys
      20             25            30

Val Cys Ile Tyr Leu Ser Ser Val Val Ser Lys Ser Ser Ala Glu Ala
      35             40            45

Asp Gly Val Leu Gln Pro Arg Arg His Pro Ala Ser Leu Leu Ile Val
      50             55            60

Phe Ala Thr Ser Ile Ser Glu Ser Ser Leu Leu Ile Phe Ser Phe Gln
      65             70            75            80

Lys Thr Glu Ala Lys Leu Ile Val Phe Ala Val Ser Leu Ala Ala Lys
      85             90            95

```

&lt;210&gt; 11

&lt;211&gt; 1498

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

```

ggacttcctc taccatcctc ggggtggacat tttgtttgct ttccattaa tccactgggt 60
tagtttccag aacgcaatgt cagacctgag gtgacagaaa agggaaaacg tgagagagtg 120
tcttcttccc tgaagagagg tgggcctgag gcttgacagt ggcgctgcct gagtgcacta 180
gccgctgggc tgggcccctg acagctgac ccaggttcct gcacttgcg ccacccttcc 240
gtagtccaca ggcctgatgg cggcggcggc tgagccgatg ggcccggcac aggttcccat 300

```

gaactcagaa gtaattgtgg accctataca ggggcagggtg aactttgagg acgtgttcgt 360  
 gtacttctcc caggaggagt ggggtgctcct tgatgaggct cagaggctcc tgtaccgtga 420  
 tgtgatgctg gagaactttg cacttatggc ctctctgggg atacctcaga caatggcagc 480  
 ttttgattg aaatatttgc tgaatgacac tggatacaca agcagcaagt ccaacacaat 540  
 cacagcaaca gactctcctg cagaccttcc caggaagaca gagccacaca caccatcatg 600  
 gtcctggtga gtcctctttt tgttttctat tcagtccata gttttctgac aatttggaca 660  
 actgtagttg caaacccagg ccagtggata gtgaccaact ctgtgttggc cgcctcatgt 720  
 tccccagcac gcagcccttt tgcctcctc atgagtata ctcatatctc tcagttctgt 780  
 tttgcctgca ggacaaggaa aacactcttt cctaactctg tttgtcatgcc atgagtcttt 840  
 tctcttcatg gaattcagct atttatcata actcttgcta agatttagga aatattaact 900  
 actagttatt tgtgatagca acatacacat ggcagtaat gctctgtcc aggaagatct 960  
 aataccagag ctaaaaatga aagtcattga tactgttaca caaagaacac tctatataac 1020  
 tgtgttaagt ccttcagaca agttcaggaa atcaaaaagt ttaaaaaggg aattcttttag 1080  
 agatttaggg gagatttctc atttttgtac tatgaagaat catggaatgt tttaaaaata 1140  
 tttttaagc acagtttgat tcagggtgctt cttgaacagc ataaatcccc tggagagtcc 1200  
 acatgtaaga aagacatgtg caggccgggc acagtggctc acgcctgtaa tcccaccgtt 1260  
 ttgggaggct gaggcactcc caaatgcctc aagtgatcca ctcaagatga cttgaggcca 1320  
 ggagttcgag accagcctgg ccaacatggc aaaaaccctg tctcaaatac aaaaatttag 1380  
 caaccatgtg gcacacacct gtagtcccag ctactccaga gggatgaagca cgagaattac 1440  
 ttaccagctt gggatgacga gggagactca aaaaaaaaaa aaaaaaaaaa aaaaaaaa 1498

&lt;210&gt; 12

&lt;211&gt; 117

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

Met Ala Ala Ala Glu Pro Met Gly Pro Ala Gln Val Pro Met Asn  
 1 5 10 15  
 Ser Glu Val Ile Val Asp Pro Ile Gln Gly Gln Val Asn Phe Glu Asp  
 20 25 30  
 Val Phe Val Tyr Phe Ser Gln Glu Trp Val Leu Leu Asp Glu Ala  
 35 40 45  
 Gln Arg Leu Leu Tyr Arg Asp Val Met Leu Glu Asn Phe Ala Leu Met  
 50 55 60  
 Ala Ser Leu Gly Ile Pro Gln Thr Met Ala Ala Phe Gly Leu Lys Tyr  
 65 70 75 80  
 Leu Leu Asn Asp Thr Gly Tyr Thr Ser Ser Lys Ser Asn Thr Ile Thr  
 85 90 95  
 Ala Thr Asp Ser Pro Ala Asp Leu Pro Arg Lys Thr Glu Pro His Thr  
 100 105 110  
 Pro Ser Trp Ser Trp  
 115

&lt;210&gt; 13

&lt;211&gt; 1718

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 13

ttctgctct cgctggaggc caggccgtgc agcatcgaag acaggaggaa ctggagcctc 60  
 attggccggc cggggcgcc ggcctcgggc ttaaatagga gctccgggct ctggctggga 120



```

ccccaccgct gccggccgcg ctcccgtgc tcctgccggg tgatggaaaa cccagcccc 180
gccgccgccc tgggcaaggc cctctgcgt ctctcctgg ccactctcgg cggcccggc 240
cagcctcttg ggggagagtc catctgttcc gccggagccc cggccaaaata cagcatcacc 300
ttcacgggca agtggagcca gacggcttcc ccaagcagta cccctgttc cgccccctg 360
cgcagtggct ttcgtgctg ggggccgcgc atagctccga ctacagcatg tggaggaaga 420
accagtacgt cagtaacggg ctgcgcgact ttgcggagcg cggcgaggcc tgggcgctga 480
tgaaggagat cgaggcgcg ggggagggcg tgcagagcgt gcacgcggtg ttttcggcg 540
ccgccgtccc cagcgccacc gggcagacgt cggcgagct ggaggtgcag cgcaggcact 600
cgctggtctc gtttgtggtg cgcacgtgc ccagccccga ctggttcgtg ggcgtggaca 660
gcctggacct gtgcgacggg gaccgttggc gggaacaggc ggcgtggac ctgtaccct 720
acgacgccgg gacggacagc ggcttcacct tctctcccc caacttcgcc accatcccc 780
aggcacgggt gaccagata acgtctctct ctcccagcca cccggccaac tccttctact 840
accgcgctc gaaggccctg cctcccatcg ccagggtgac actggtgcgg ctgcgacaga 900
gccccagggc cttcatcct cccgccccag tcctgccag cagggaacaat gagattgtag 960
acagcgctc agttccagaa acgccgctgg actgcagggt ctccctgtgg tcgtcctgg 1020
gactgtgcgg agggcactgt gggaggctcg ggaccaagag caggactcgc tacgtccgg 1080
tccagcccgc caacaacggg agcccctgcc ccgagctcga agaagaggct gagtgcgtcc 1140
ctgataactg cgtctaagac cagagcccc cagccccctg gggcccccg agccatggg 1200
tgtcgggggc tcctgtgcag gctcatgctg caggcgccg agggcacagg gggtttcg 1260
ctgctcctga ccgcggtgag gccgcgccga ccactctgc actgaaggc cctctggtg 1320
ccggcacggg cattgggaaa cagcctctc ctttcccaac cttgcttctt aggggcccc 1380
gtgtcccgtc tgctctcagc ctctctctcc tgcaggataa agtcatcccc aaggctccag 1440
ctactctaaa ttatgtctcc ttataagta ttgtgctcc aggagattgt ccttcacgt 1500
ccaggggcct ggctcccacg tggttgcaga tacctcagac ctggtgctct aggcgtgct 1560
gagccactc tcccagggc gcattcaagc gggggccact tgagaagtga ataaatggg 1620
cggtttcgga agcgtcagtg tttccatgtt atggatctct ctgcgtttga ataaagacta 1680
tctctgttgc tcacaaaaaa aaaaaaaaaa aaaaaaaa 1718

```

<210> 14  
 <211> 105  
 <212> PRT  
 <213> Homo sapiens

<400> 14  
 Met Glu Asn Pro Ser Pro Ala Ala Ala Leu Gly Lys Ala Leu Cys Ala  
 1 5 10 15  
 Leu Leu Leu Ala Thr Leu Gly Ala Ala Gly Gln Pro Leu Gly Gly Glu  
 20 25 30  
 Ser Ile Cys Ser Ala Gly Ala Pro Ala Lys Tyr Ser Ile Thr Phe Thr  
 35 40 45  
 Gly Lys Trp Ser Gln Thr Ala Ser Pro Ser Ser Thr Pro Cys Ser Ala  
 50 55 60  
 Pro Leu Arg Ser Gly Leu Arg Cys Trp Gly Pro Arg Ile Ala Pro Thr  
 65 70 75 80  
 Thr Ala Cys Gly Gly Arg Thr Ser Thr Ser Val Thr Gly Cys Ala Thr  
 85 90 95  
 Leu Arg Ser Ala Ala Arg Pro Gly Arg  
 100 105

<210> 15  
 <211> 847  
 <212> DNA  
 <213> Homo sapiens

&lt;400&gt; 15

acacacacat ctgcacctca accacagact acacttgctg aactggctcc tggggccatg 60  
 aggtgtgtcac tgccactgct gctgtgtgctg ctgggagcct gggccatccc agggggcctc 120  
 ggggacaggg cgccactcac agccacagcc ccacaactgg atgatgagga gatgtactca 180  
 gccacatgc cgcctcacct gcgtgtgat gcctgcagag ctgtggctta ccagatgtgg 240  
 caaaatcttg caaaggcaga gaccaaactt catacctcaa actctggggg gcggcgaggag 300  
 ctgagcgagt. tggcttacac ggatgtcctg gaccggagct gctcccggaa ctggcaggac 360  
 tacggagttc gagaagtgga ccaagtgaac cgtctcacag gcccgaggact tagcgagggg 420  
 ccagagccaa gcatcagcgt gatggtcaca gggggccctt ggctaccag gctctccagg 480  
 acatgtttgc actacttggg ggagtttggg gaagaccaga tctatgaagc ccaccaacaa 540  
 ggccgagggg ctctggaggc attgctatgt gggggacccc agggggcctg ctcagagaag 600  
 gtgtcagcca caagagaaga gctctagtcc tggactctac cctcctctga aagaagctgg 660  
 ggcttgctct gacggtctcc actccgctct gcaggcagcc aggagggcag gaagcccttg 720  
 ctctgtgctg ccctcctgcc tccctcctcc agcctcaggg cactcggggc tgggtgggag 780  
 tcaacgcctt cccctctgga ctcaataaaa acccagtgc ctcaaaaaaa aaaaaaaaaa 840  
 aaaaaaa 847

&lt;210&gt; 16

&lt;211&gt; 189

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

Met Arg Leu Ser Leu Pro Leu Leu Leu Leu Leu Gly Ala Trp Ala  
 1 5 10 15  
 Ile Pro Gly Gly Leu Gly Asp Arg Ala Pro Leu Thr Ala Thr Ala Pro  
 20 25 30  
 Gln Leu Asp Asp Glu Glu Met Tyr Ser Ala His Met Pro Ala His Leu  
 35 40 45  
 Arg Cys Asp Ala Cys Arg Ala Val Ala Tyr Gln Met Trp Gln Asn Leu  
 50 55 60  
 Ala Lys Ala Glu Thr Lys Leu His Thr Ser Asn Ser Gly Gly Arg Arg  
 65 70 75 80  
 Glu Leu Ser Glu Leu Val Tyr Thr Asp Val Leu Asp Arg Ser Cys Ser  
 85 90 95  
 Arg Asn Trp Gln Asp Tyr Gly Val Arg Glu Val Asp Gln Val Lys Arg  
 100 105 110  
 Leu Thr Gly Pro Gly Leu Ser Glu Gly Pro Glu Pro Ser Ile Ser Val  
 115 120 125  
 Met Val Thr Gly Gly Pro Trp Pro Thr Arg Leu Ser Arg Thr Cys Leu  
 130 135 140  
 His Tyr Leu Gly Glu Phe Gly Glu Asp Gln Ile Tyr Glu Ala His Gln  
 145 150 155 160  
 Gln Gly Arg Gly Ala Leu Glu Ala Leu Leu Cys Gly Gly Pro Gln Gly  
 165 170 175  
 Ala Cys Ser Glu Lys Val Ser Ala Thr Arg Glu Glu Leu  
 180 185

<210> 17  
 <211> 1448  
 <212> DNA  
 <213> Homo sapiens

<400> 17  
 atttcatccc tgcgctgtcc tagcaccctg tgcgtggagt tggcatgcag agactttgtc 60  
 aggatgggag aggccgcagt gtatcggatc tgcgttctg gaaagaagac cttctatatt 120  
 agagtagaaa caaacgaaac ttctaaggta tcatctgtgt taagtgatga gaccatattt 180  
 ctttgatgtt tctgaacatc aaagctgatt cagtactggt agatgtgtc attctccctg 240  
 aaacataccc atcatatttc ctattataat tacatctcat tgcctgtgg aggtggacat 300  
 gataaacatt atcttttgtt ttcttgtttt gttttgtttg agacgggtc attctgtcac 360  
 ccagactgga gtgcagtgcc acaatcatgg ctaccgcgat tgacctcctt ggctcaaggc 420  
 atctctccac ctccagcttc tgactagctg ggactactgg tgtgcaccac cacaccagc 480  
 taattttcaa tttttcatag agacagggc tccactgtgt gtccaggctg gtcttgaact 540  
 cctgggctca agccaccgc ccacctgggc ctcccaaagt gctgggatta caggcatcag 600  
 ccatcacacc catccataaa cattatatta atgtcacatt acaaaaactga gacctaaagt 660  
 gcttaggata aaatgaaatt gaaagactag ctaacatgaa aatttatatt ttggcttttt 720  
 catgtttttt gataaaacca gtgtatttga atgattcttt tgatgtttag taatggtttt 780  
 ttgtttgttt tttgtttttt ttttttgag acggagtctc actctgtcgc cagtctggag 840  
 tgctagtggc gcgactctgg cttactgcag cctccacttc ccaagttcaa gcgattctcc 900  
 tgctcagcc tcccgagtag ctgggactac aggcgcagc caccacgccc ggctaatttt 960  
 tgtattttta gtagagacgg agtttcactg tgttgccag gatggctctg atctcgacct 1020  
 cgtgatcacc caccctggcc tctcaaagt ctgggattac aggcgtgagc caccacgcct 1080  
 ggcctatgtt taataatgtt gaaataggat ggaatatttt gttaaattaa cattttaaaa 1140  
 ttagaagaca ccgttttaatt ttttaaacc ttctctctct cattgtaacg aaattaattc 1200  
 cagctgcagt gagaaaactt aaaaatcatg atacaaaatg aaacaatc tgaaagtagt 1260  
 tttataaaac tgaaattgct gttaaagaga atgtgttagt gacttaacca tttgctctat 1320  
 gtgatgttta ttatcaaata catataattt tgaagatttt aatgaatggc ttaagatttt 1380  
 atctttgtgt agaatgtggc taaagaaacc ttagttgaga ttcaaaaaaa aaaaaaaaaa 1440  
 aaaaaaaaaa 1448

<210> 18  
 <211> 106  
 <212> PRT  
 <213> Homo sapiens

<400> 18  
 Met Val Phe Cys Leu Phe Phe Gly Phe Phe Phe Leu Arg Arg Ser Leu  
 1 5 10 15  
 Thr Leu Ser Pro Val Trp Ser Ala Ser Gly Ala Ile Leu Ala Tyr Cys  
 20 25 30  
 Ser Leu His Phe Pro Ser Ser Ser Asp Ser Pro Ala Ser Ala Ser Arg  
 35 40 45  
 Val Ala Gly Thr Thr Gly Ala Cys His His Ala Arg Leu Ile Phe Val  
 50 55 60  
 Phe Leu Val Glu Thr Glu Phe His Cys Val Gly Gln Asp Gly Leu Asp  
 65 70 75 80  
 Leu Asp Leu Val Ile Thr His Leu Gly Leu Ser Lys Cys Trp Asp Tyr  
 85 90 95  
 Arg Arg Glu Pro Pro Arg Leu Ala Tyr Val  
 100 105

<210> 19  
 <211> 2166  
 <212> DNA  
 <213> Homo sapiens

<400> 19  
 atgatttttg gagcttgctt cctcattgac ttcattgtatt ttgaaagatc tccaccacaa 60  
 cgttcaagat agagagacac agcaagccat caactatggt taattttgaa aaatggaaaa 120  
 gttggatttg gcttacagtc agcactcagt tatattgcaag tgtatttctt tgctttgtag 180  
 agtattttta ttgggtgtta actttgacag ctgagagtgg gcttgcaaga acacaatcta 240  
 aaagtgtgtt tcaattgagt atctctctag tagaatagga gttcatcctg aaaagctgtg 300  
 actcattaac ccagtaaaca tatacaaagt aagcttaaaa cactataaac atgagataag 360  
 ggaaaaatgaa tccagagttc tcatattaat aggtagtga acaataaggc tttttagagc 420  
 agactttgtt ggcataaaat aacctggctt ctatccctaa ccctttccta cctttcctct 480  
 ccgtcaacat gtcctcatac tgaagacaaa cttgtttcaa tgatagtctt cattttcaca 540  
 aacaaaaagg caggcagaca gaaataatga tgttttcttg cactaagaag gtactacttg 600  
 tacacatata tcaaaacctc attctgcaaa gtttttgaa gtttcaatgg gaaatttgat 660  
 tttattacaa aataaaacat tttttaatgt taaagtttat atattccatg cttgttttct 720  
 cattcactgg catggatgat caggagctgc ctatatatga aggcagaatc agactatcag 780  
 gaaaggagct gccagggcc acagccagtc aagatctctg agcaacttag agacattggc 840  
 gtcattatat gaagcttgca ttaatacat ttatacataa tacatttgta catttaattc 900  
 ataacgtctc ttggtcacag atgccttata tataaaataa gttgccagat ctctaagatt 960  
 gcttagtaca cctttgtatc tcatttgatg tgataccag aagagatcat tgttttttgt 1020  
 ttttgttttt gtttttttca agaagatcct tcgtgatcac catgctgttc tcatggtaag 1080  
 aactggagtt atgtttttta atttggaaat atgacatttt atgtagcact ttataaaaag 1140  
 tgaagcgcg aaattccacc gctgcttaat actgctttgc ttctttttat tgacatgata 1200  
 gataaataatg tatctacaca gagtaataat aataaaacac agtaaacatt ctatttctct 1260  
 atgggtctaca gcatgccagt aaataatatg taggaccaat aataaattat caattacaca 1320  
 tttttgtgtt aactaattaa aagcatagtg tataagttag tacactctaa ttaacttgct 1380  
 tctgttgac tttagttttc tacctgcata tggactgcat tttttttttt aacacagtca 1440  
 gtatgtagaa tgggatgtat tcttctgctg ctgcttatta aataaagaaa gcctgagtgt 1500  
 tcttagatgg ggttattctg agatgagggc cttagcctac agttcttttt gaaatgaaag 1560  
 gtgctttgtt ttttaattat attcatcttt tcagggtaaa tttgtttttc tgagtttctc 1620  
 gtaatgctca tttttacatg ctgctactag cttttttttt taaaaaaagt aaaagtgtgt 1680  
 gctttctaaa atattaattg ccttatattt gaaagtgcc ttgcaatcgt aagtagacta 1740  
 tgattttcct ataattgatg ctgatattta aataggaaat cagacaaaca atattcagaa 1800  
 agtttaagca tataaacttt ttatttttaa cttgcctaaa tccctgtatt ccaaaacctg 1860  
 ctgcatcata ataatatat ctatatatat ttagcataag acgtgatatt tttaatttct 1920  
 tttttaaaaa attatatttg tctcttagag ttaaaatttt ctttatataa tattgtcata 1980  
 tgtcatagtt ttaatacaat tcacatgatt tctatgtttc ttaatgatat tttgttgtgt 2040  
 aaaattgatc ggattgatta aaaaacaaat tctctggaat ttgtgcgttc atgctttttc 2100  
 gtattcttta tggcttttaa ataaatatac aatgggttaat agtaaaaaaa aaaaaaaaaa 2160  
 aaaaaa 2166

<210> 20  
 <211> 60  
 <212> PRT  
 <213> Homo sapiens

<400> 20  
 Met Val Asn Phe Glu Lys Trp Lys Ser Trp Ile Gly Leu Thr Val Ser  
 1 5 10 15  
 Thr Gln Leu Phe Ala Ser Val Phe Leu Cys Phe Val Glu Tyr Phe Tyr  
 20 25 30  
 Trp Val Leu Thr Leu Thr Ala Glu Ser Gly Leu Ala Arg Thr Gln Ser  
 35 40 45

Lys Ser Val Phe Gln Leu Ser Ile Ser Leu Val Glu  
50 55 60

<210> 21  
<211> 1833  
<212> DNA  
<213> Homo sapiens

<400> 21  
gtgtgtgtaa tatcaacgca gtgtatttaa tttggggaag tctactagggtt agaactgtag 60  
gcatactcag tggagaggaa gctacctatt ctattctaac tattctgagg tcttcctgga 120  
gagggattgt tttagcagct cctctccaga gggggctgga agtccagttt caccaatgac 180  
caagttttat ttccttgctc ttgttttagt tttttcattt tgttttcagt ccaggtgcct 240  
cgcagctctt ttggaatggg ccagcattag tctaatttta agcgtatgt gttttgtacc 300  
cttgcaaaact tgccttctct ttcttcttgc tgttgccctt tgaatgaagc cttgtgtgct 360  
tgcattgatca gacctgttaa taggatgaat ttccatgcac ttaacatgaa gagatacaaa 420  
aagaattctag agagagatat atatttttaa tgaaatagaa gcttggcata gaaacattga 480  
tttagatggt atgagttgct ggccagagat tcttcagata aaagagccta gaaaattgag 540  
ttttgagcgg gcatggtggc tcacgcctgt aatgccagta gtttgggagg ctgaggcggg 600  
tggatcacct gaggtcagga gttcgagatc agcctggcca acatgggtgaa accccgcctc 660  
tactaaaaaa tatgcaaaat tagccgggcyg tgggtgtgca ggctgtaat ccagctact 720  
cagaggctga ggcaggagat tcacttgagc ccaggaggtyg gaggttgtag tgagccaaga 780  
tcgcgccatt gcactccagt cagggaaca agagtgaac tccgtctcaa aatataaaaag 840  
aaaattgagt gtcccaccca aatttttagg cgtacctcct tccatttgat aagaagatgc 900  
tatttccagt gtccctggga gaattcttcc atccagttta tatggtgaat tgccaccagc 960  
gttccctaaa ggaagagaga acttagctta tttctgtctg tgacttttat cagctagatc 1020  
agttaattgt atttttaaat taaaaattag taactagagg aagagactaa gagaactatt 1080  
tttacctgag taaatgctct cctcttcttg tgtccaaatc taatgcagag gtgaaaacac 1140  
ttgatgagaa aatcctttgt ggcttgaggg gaatcatggt tctttcagga cgtgtgtgtt 1200  
ccctgaagtc tatgtcacac tgagaacttt ttctgactgc actgcttttg agctattcat 1260  
ttaaattgag tttggcctag ttttccctgt ataaaacagg aaatctacat tgttccctct 1320  
ttctctccct tccatccctt cccctgcctt tctatacac gctggtgtgc ttttgcctc 1380  
agaaattctt ggtttgggag ttatgtattt ttgagtgaac gaggttatga catacttgga 1440  
gcggcatgat ttcaagaagc cattccaaat tcttatagct ccttagctct ggttttactc 1500  
tcgtattttc cccaagaaaa ttttttggtc attcaaagaa agcacaaaat atgtgatgtg 1560  
gaactcatga aaatagggac ttttgatata tctttaaata ttccctgact cacggaaaag 1620  
atgtcctctt ttggtgtata cattgtgaaa atctacccat ttatctctct ctctctctt 1680  
tcttttaaat aaattatctg ataaggccgg caaggtggct cagcctgta atcccagcac 1740  
tttggggagg cgaggcgagc ggatcacgag gtcaggagat cgagaccgcy gtgaaacccc 1800  
gtctctacta aaaaaaaaaa aaaaaaaaaa aaa 1833

<210> 22  
<211> 55  
<212> PRT  
<213> Homo sapiens

<400> 22  
Met Thr Lys Phe Tyr Phe Leu Val Leu Val Leu Val Phe Ser Phe Cys  
1 5 10 15  
Phe Gln Ser Arg Cys Leu Ala Ala Leu Leu Glu Trp Ala Ser Ile Ser  
20 25 30  
Leu Ile Leu Ser Ala Met Cys Phe Val Pro Leu Gln Thr Cys Phe Leu  
35 40 45  
Phe Leu Leu Ala Val Ala Leu  
50 55

<210> 23  
 <211> 1504  
 <212> DNA  
 <213> Homo sapiens

<400> 23  
 tctcattgct ggcattgcca ttcctgccat ggtcattggc attcctgttt atgttggaag 60  
 gaagattcac agcaggtatg aggggaaggaa aacctccaaa cacaagagga atttggttat 120  
 cactggagga gtgactttgt cggctattgc atccccagtt attgctgcag ttagtggttg 180  
 tattggtgtc cccattatgc tggcatatgt ttatgggggt gtgcccattt ctctttgtcg 240  
 tggaggcggc tgtggagtta gcacagccaa cggaaaagga gtgaaaattg aatttgatga 300  
 agatgatggt ccaatcacag tggcagatgc ctggagagcc ctcaagaatc ccagcattgg 360  
 ggaaagcagc attgaaggcc tgactagtgt attgagcact agtggaagcc ctacagatgg 420  
 acttagtggt atgcaaggte cttacagcga aacggccagc tttgcagccc tctcaggggg 480  
 cacgctgagt ggcggcattc tctccagtgg caagggaata tatagcaggt tagaagttca 540  
 agccgatgtc caaaaggaaa ttttcccaa agacacagcc agtcttggtg caattagtga 600  
 caacgcaagc actcgtgcta tggccggttc cataatcagt tcctacaacc cacaggacag 660  
 agaatgcaac aatatggaat tccaagtga cattgaagcc aaaccaagcc actatcagct 720  
 ggtgagtggg agcagcacgg aggactcgct ccatgttcat gctcagatgg cagagaatga 780  
 agaagaaggt agtgggtggc gaggcagtga agaggatccc ccctgcagac accaaagtctg 840  
 tgaacagaaa gactgcctgg ccagcaaacc ttgggacatc agcctggccc agcctgaaag 900  
 catcgcgagt gacctagaga gttctgatgc acagtcagac gatgtgccag acatcacctc 960  
 agatgagtgt ggtccccc gctccatac tgcagcctgc ccctcgacc ccagagccca 1020  
 aggtgcaccg agcccaagtg cccatatgaa cctctctgcc ctagecagg gacaaactgt 1080  
 cttgaagcca gaaggtggag aagccagagt atgaagtggg atgaatgctc ctgttctgag 1140  
 aagcacactt gtaactgcat cttttggaat tttttttttt ttttttccaa ggggtagaga 1200  
 tttatgtatt ttatttcaca gattctctgg tcacaggttt ttgcccagg aaattctgag 1260  
 aaattcacia tttcttacca gataaaacat gaaaagttt ccgttagttc ccctcccctc 1320  
 ccctcccctc ttttagtttt aatttattgg ttaaaactgat ggcagcaatc catgaggtgt 1380  
 gtcaaagagt gtacatatgt atgtgtgtat attgaatgct aaacatatta ctgaaagaca 1440  
 cattttaata aagatttctg tcataattca aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1500  
 aaaa 1504

<210> 24  
 <211> 361  
 <212> PRT  
 <213> Homo sapiens

<400> 24  
 Met Val Ile Gly Ile Pro Val Tyr Val Gly Arg Lys Ile His Ser Arg  
 1 5 10 15  
 Tyr Glu Gly Arg Lys Thr Ser Lys His Lys Arg Asn Leu Ala Ile Thr  
 20 25 30  
 Gly Gly Val Thr Leu Ser Val Ile Ala Ser Pro Val Ile Ala Ala Val  
 35 40 45  
 Ser Val Gly Ile Gly Val Pro Ile Met Leu Ala Tyr Val Tyr Gly Val  
 50 55 60  
 Val Pro Ile Ser Leu Cys Arg Gly Gly Gly Cys Gly Val Ser Thr Ala  
 65 70 75 80  
 Asn Gly Lys Gly Val Lys Ile Glu Phe Asp Glu Asp Asp Gly Pro Ile  
 85 90 95  
 Thr Val Ala Asp Ala Trp Arg Ala Leu Lys Asn Pro Ser Ile Gly Glu  
 100 105 110

Ser Ser Ile Glu Gly Leu Thr Ser Val Leu Ser Thr Ser Gly Ser Pro  
 115 120 125  
 Thr Asp Gly Leu Ser Val Met Gln Gly Pro Tyr Ser Glu Thr Ala Ser  
 130 135 140  
 Phe Ala Ala Leu Ser Gly Gly Thr Leu Ser Gly Gly Ile Leu Ser Ser  
 145 150 155 160  
 Gly Lys Gly Lys Tyr Ser Arg Leu Glu Val Gln Ala Asp Val Gln Lys  
 165 170 175  
 Glu Ile Phe Pro Lys Asp Thr Ala Ser Leu Gly Ala Ile Ser Asp Asn  
 180 185 190  
 Ala Ser Thr Arg Ala Met Ala Gly Ser Ile Ile Ser Ser Tyr Asn Pro  
 195 200 205  
 Gln Asp Arg Glu Cys Asn Asn Met Glu Ile Gln Val Asp Ile Glu Ala  
 210 215 220  
 Lys Pro Ser His Tyr Gln Leu Val Ser Gly Ser Ser Thr Glu Asp Ser  
 225 230 235 240  
 Leu His Val His Ala Gln Met Ala Glu Asn Glu Glu Glu Gly Ser Gly  
 245 250 255  
 Gly Gly Gly Ser Glu Glu Asp Pro Pro Cys Arg His Gln Ser Cys Glu  
 260 265 270  
 Gln Lys Asp Cys Leu Ala Ser Lys Pro Trp Asp Ile Ser Leu Ala Gln  
 275 280 285  
 Pro Glu Ser Ile Arg Ser Asp Leu Glu Ser Ser Asp Ala Gln Ser Asp  
 290 295 300  
 Asp Val Pro Asp Ile Thr Ser Asp Glu Cys Gly Ser Pro Arg Ser His  
 305 310 315 320  
 Thr Ala Ala Cys Pro Ser Thr Pro Arg Ala Gln Gly Ala Pro Ser Pro  
 325 330 335  
 Ser Ala His Met Asn Leu Ser Ala Leu Ala Glu Gly Gln Thr Val Leu  
 340 345 350  
 Lys Pro Glu Gly Gly Glu Ala Arg Val  
 355 360

&lt;210&gt; 25

&lt;211&gt; 2350

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 25

agtcccgtag taatgtacca tcagcttctg aagttcctga taatagggca tctgaagctt 60  
 ctcagggatt tcgatttctt aggcgaagat ggggtttgtc atctcttagc cacaatcata 120  
 gctctgagtc agattcagaa aattttaacc aagaatctga aggtagaaat acaggaccat 180  
 gggtatcttc ctcaactaga aatagatgca cacctttgtt ctctagaagg aggcgagagg 240

```

gaagagatga atcttcaagg atacctacct ctgatacatc atctagatct catattttta 300
gaagagaatc aaatgaagtg gttcaccttg aagcacagaa tgatcctctt ggagctgctg 360
ccaacagacc acaagcatct gcagcatcaa gcagtgccac aacaggtggc tctacatcag 420
attcggctca aggtggaaga aatacaggaa tatcagggat tcttcctggg tctttattcc 480
ggtttgcagt cccccagca cttgggagta atttgaccga caatgtcatg atcacagtag 540
atattattcc ttcaggttgg aattcagctg atggtaaaag tgataaaact aaaagtgcgc 600
cttcaagaga tccagaaaga ttgcagaaaa taaaagagag cctcctttta gaggactcag 660
aagaagaaga aggtgactta tgtagaattt gtcaaatggc agctgcatca tcatctaatt 720
tgctgataga gccatgcaag tgcacaggaa gtttgagta tgtccaccaa gactgtatga 780
aaaagtgggt acaggccaaa attaaactctg gttcttcatt agaagctgta accacctgtg 840
aactatgtaa agagaagttg gagcttaacc tggaggattt tgatattcat gaactacata 900
gagctcatgc aaatgaacaa gctgagtatg agtttatcag ctctgggtctc tacctagtgg 960
tgttattgca tctgtgcgaa caaagctttt ctgatatgat gggaaatata aatgaaccaa 1020
gcacacgtgt ccgacttcag aggatgattc cgaagaagac ggagaccata acaggacatt 1080
tgatattgcc taacttcata taagacagat ggatgatctg tgaacataag tgtttattaa 1140
aaatggcaat taaatataaa ttacttttgt gggggaatgc ctaataaata cattgactat 1200
atataaaatg aatatataca tacacatgta tgcctgtata tatatatcca tctccagtgt 1260
ttgtgtaatt aaaattctgc tggacttttt aacatagcaa atccgatgtt tataaactgg 1320
taatcaaaaa gggtttttct tttaggtgag tgggaaagta ttacccttgt tttaaatatc 1380
taagcaatgc ctatcaaccc ttttttgtgt tatgattact gtagtcatat ttatgaaaaa 1440
aggtttgtgt tttactcttg ctagtggaga aagtgggaca aaatatactt ttgaaataaa 1500
atgctatatg gcacctaatt attttttctt ttaaaatgcc ttaagttgca gtctcatttt 1560
gataatcatt tgcttccagt gtttaaaaat taaaaaaga atggggagaa ggttatgaga 1620
agagcattat taagtttcca aatttaattt gaattccaaa ttcacctagc aataaaatct 1680
aatttttaaa aagtatataa atataaaatg tataaatgat ggatagattt ttgtattgat 1740
ttgcaaatg cagattatat ttgataggct atagtatgta gatattcctt ttaggaatat 1800
tacagctgta aattatatga gacttgccag tcaaatgcta tttggttaa aaaaattatt 1860
gcaatctcaa gttaatggaa tatttttaaa tcccacattc agagtttaaa acactggttt 1920
tcaatgtgtt ttttagtgtt gtcacttgtt tatagataaa tatataaata acctgtttgg 1980
atcctgggcc tttttaactg ttccttggtg attctgagca tttatttgat gacttaatat 2040
ttttcactac ctttggagaa cagatgaaca ttattcacca tgaatggatc tatactgtgt 2100
ggtcagtgtg tgtgtatact tccataaacac tgtatttttc ttctgtcagt acccttagga 2160
tacactttaa aacaccttaa ggtctgatgt tatggcaaca aactactttt tcaaacctaa 2220
ataggaacca tgcaatttct caaaagtgat tgaacagttt gccacactt agtttgttgg 2280
tcttatgtaa aacattggct caaaataaag tacacactga tttataaaaa aaaaaaaaaa 2340
aaaaaaaaaa 2350

```

&lt;210&gt; 26

&lt;211&gt; 167

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 26

```

Met Tyr His Gln Leu Leu Lys Phe Leu Ile Ile Gly His Leu Lys Leu
  1             5             10             15

```

```

Leu Arg Asp Phe Asp Phe Leu Gly Glu Asp Gly Val Cys His Leu Leu
  20             25             30

```

```

Ala Thr Ile Ile Ala Leu Ser Gln Ile Gln Lys Ile Leu Thr Lys Asn
  35             40             45

```

```

Leu Lys Val Glu Ile Gln Asp His Gly Tyr Leu Pro His Leu Glu Ile
  50             55             60

```

```

Asp Ala His Leu Cys Ser Leu Glu Gly Gly Glu Arg Glu Glu Met Asn
  65             70             75             80

```

```

Leu Gln Gly Tyr Leu Pro Leu Ile His His Leu Asp Leu Ile Phe Leu
  85             90             95

```



Glu Glu Asn Gln Met Lys Trp Phe Thr Leu Lys His Arg Met Ile Leu  
100 105 110

Leu Glu Leu Leu Pro Thr Asp His Lys His Leu Gln His Gln Ala Val  
115 120 125

Pro Gln Gln Val Ala Leu His Gln Ile Arg Leu Lys Val Glu Glu Ile  
130 135 140

Gln Glu Tyr Gln Gly Phe Phe Leu Val Pro Tyr Ser Gly Leu Gln Ser  
145 150 155 160

Pro Gln His Leu Gly Val Ile  
165

<210> 27

<211> 1635

<212> DNA

<213> Homo sapiens

<400> 27

```

gaaaatcaca gttgtaaaag cagtctccag ggccagggtat atttagtata gtttttagaa 60
acagcagcca gaaggataat gggaaataaaa gttagaagcc tctctctctc tctctctctc 120
tctctctctg tgtgtgtgtg tgtgtgtgtg tgtgtttgtg tttgtatgtt tgtcttgtct 180
ctatcatcca tccctatggt gattgggaga caggatgctt taatcaagcc ccaggggagc 240
aggggttttag tcttacagca tccctgtgcta acttgctgtg tgacctaga atatttcttg 300
gcatctctgg gcttcaggcg ctgtctctat acattagtgt gttactgac ttgtaagtag 360
gaatttagag gcccaggag ggaagggct tgcttcagtc agttagctta gcctctacac 420
tgagcactct ttaggtgtc tagttttctc cttcttaaat ggaaagaaag aggcctgacc 480
agtcatagg tgcaactctt cttgaaatat aaagcactgc aaatacacaa ggctgaagaa 540
aagaatcaca aagcttttgt gattgggtcat agccattttc ataaagcagc agccagactt 600
gccagctaac ttattaccta tgggtgaaac ccgttagttc tgaaatgtta cactgttctt 660
tgaccactat ttgagaatta agtgctttgt gattaaaaac aacagcaaca atgctgaatc 720
ttgtaattat ttcataatta agtaaataaa agctgaggtg acctctcgg attttgcaa 780
gccctgaatg taacttgaa ggggaaaccc attcatttgg cctgaacggg agattacctt 840
gtaggggagc tgcgtgggtga atatcaaggg tcttttgaag aaaaacagag tagaagagca 900
attgggtaga tacacaagtc aggaaaaatg caaagtgact tagtttgatg cctccaaatg 960
tgtgagttgc tagaaagagg gctcctcact atggctgatc tcttgggggt agaggaaaga 1020
ggagatctga agtaagtga gtgtaaatag aagggtttgt acattcttat ttgggagttg 1080
aagccaagtg gaggagtgt gtattttagg agaagtggc atataaaatg tgatcttaaa 1140
actaaataat ttgagataag ttaattgcag tacaaaaaga tttttacctt actaaaggga 1200
ttcatagtga gtctcctttt aatagctaatt tccgttatta tatttcaaat tatttgatga 1260
acacaatcat gatagaccct tttgggagcc tcacctgtta tgcaaaacca aggacttttc 1320
ggttgggcac ggtggctcac ccctgtagtc ccagcacttt gggaggccga ggcaggtgga 1380
tcacgaagtc aagagattga gaccatcctg gccaatatgg tgaaacccc tgtctactaa 1440
aaatacaaaa aaaaaaatta gctgggcatg gtggcgctg cctgtagtcc cagctactcg 1500
ggaggctgag gtaggagaat ggctcgaacc tgtagggcag aggttgagc gagttgagat 1560
gccgccactg cactccagcc tggggaaaga gcaagactcc ttctcaaaaa caaaaaaaa 1620
aaaaaaaaa aaaaaa
1635

```

<210> 28

<211> 89

<212> PRT

<213> Homo sapiens

<400> 28

Met Gly Ile Lys Val Arg Ser Leu Ser Leu Ser Leu Ser Leu  
1 5 10 15

Ser Val Cys Val Cys Val Cys Val Cys Val Cys Val Cys Met Phe Val  
 20 25 30

Leu Ser Leu Ser Ser Ile Pro Met Leu Ile Gly Arg Gln Asp Ala Leu  
 35 40 45

Ile Lys Pro Gln Gly Ile Arg Gly Leu Val Leu Gln His Pro Val Leu  
 50 55 60

Thr Cys Cys Val Thr Leu Glu Tyr Phe Leu Ala Ser Leu Gly Phe Arg  
 65 70 75 80

Arg Cys Leu Tyr Thr Leu Val Cys Tyr  
 85

&lt;210&gt; 29

&lt;211&gt; 3415

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 29

atatacctaa aactgattaa tgttacgggc ctattgaatg tttgtttctg cttgtgtgtt 60  
 ttcttttttc tatgtttatg aaaatatata catcatcagt tctctgttc cgagggatgt 120  
 tgtcatgctt ggcattgtct tgttggtgta gtttgtctgc acccctcagc tctgtgggta 180  
 atggtgattg tgggagctgc ctacagtatct ctgacagttc taatgatacg ggaaagtaga 240  
 actatctgct taggatagat tttaggatta gggttttctg tgtttatgtg aagtattttt 300  
 atgtgttgag gtataactaa aatcatctaa ggctaaatgt aatgaaacag ctcataacag 360  
 atgaaatgta catgaataga ttatcctgca gggggagcaa gaggcagacg ggttaaaagt 420  
 ctgttgggct tttccccaga acaaaacagt aggccttcag gcctgtcact catacagaat 480  
 gaatcacaaag tattttcagg agataagtgt gggtaatatc attcattcgt tgtcattatg 540  
 gttgccacca agaataaggga gctattttaa tgtatattaa attaataaaa attaaggaaa 600  
 ccttaaaatt tagctctca atggcactag tacattctaa gtgtgcaata gccccatgtg 660  
 tgtgtgtgtt atattagcaa tacagatacg gagagtctt atcatcgagc aagtctatga 720  
 aacagtaccg gttttgtcag actgttataa acctttgtgt cttaatgttc gtttattgat 780  
 ttatttaaac agtggtaata tatagagttt aacaagggga gttatcagtt aacaagttcc 840  
 tgctcatgca caaagaagaa atcaagtagc ggtgtgatgt tagcttgtaa agaaatcatg 900  
 gatctgcatt agtaagtcac aggtactcaa ggaccctgg gagtacttgt tttggcagag 960  
 ttgcctggca gtaagggcac caaaatagct atgggaagga ggcagtttta ctacttctgt 1020  
 agatcaggaa atggtcttag tgatacttgg acttggtcac agatacttct gttggtagaa 1080  
 ttcaggactc atgaatattt tagtataagc cttttttctt cctcagtcctg tgtgagcccc 1140  
 atgcaggact agggaaagt gtaaggagga cctcggctct tgtgtgttctc aggagtctct 1200  
 tggctgatta atcatgttgt tactcatatt gagtaatact aagcccttga agacttcagg 1260  
 gtgggtatacc tggcattgtc cttgatttta aaatatcttg gaatctatta taagaagatt 1320  
 aggatcatta gcgaaagtac tcattgatgg tcaaaataca ttaaagagct ggaaaaggaa 1380  
 actgtgaggt gtgatctctc tctctgaatt tttcccctgc ttgtttggat gaatgaatag 1440  
 aaggcatatt tataaagttt gcagaagaca actaaaacag ttgagagggc tatgttgata 1500  
 ctgacctgtg cttctcttgc ttttttattt gctgcttttc agtaaaatat gaaagaatca 1560  
 aatttctggt gattgctttg aagagttctg tggaaagtcta tgcgtgggca ccaaagccat 1620  
 atcacaaatt tatggccttt aaggtaacaa catcaagtga atttaaaagt agtattggcc 1680  
 attcaagctg caaccaagag tcagggaatg tgtttaaaaa gtctgaatgt taaaattgct 1740  
 aatataaaaag ctatgtgcta atatagcata taactttatc ataaaccatt tctaattgta 1800  
 taagcttagt taagctgctt tctaagccca cagtgagaag gagagagaga taaatgttgg 1860  
 gtagacactt taatcgatgt ggcaatgtgt tcacagagga aaagagaaca gtacttcac 1920  
 ccttcagtta aaaaggtgac cttcacctga gtcattggaag cgtgtaaaga tttagatgtg 1980  
 tttttgataa caaaactgtg tctatcgggc agttttaaga tatatctgtt cataaaatac 2040  
 taattaaaaa taaattaca gaaattctga tgacaacatt atatactaag tgaaaaaagt 2100  
 taaaatattt catatgatc catttttgtt tcaataaaaa actcagctat aacatctgaa 2160  
 ctaatgtaca aattaagatg tttttgccat tttgcatgta tacagtttta ggaaagtaaa 2220

```

tgatggagta cttagtctta aaattaggac tgttttcatt tgtgagttca caaaaatact 2280
catgaaattt acaaatatac ctcacattgc ctgggtgattg gctttttaga atagtttttt 2340
tatattttat tgaaggggta gggtttcattt atttgcaaaa tttgtgtttt tggattgctt 2400
actgcttgat ttcccagtga agcaggatag atggagtcac aatattcgct taaaaataa 2460
tattcactta aaaaaataat ccaagtgtta ctgaataaag agaattgggt atacagttaa 2520
attatcttct gagatctggc cttaatatcc tttatatacc aggtaccgta ctagtgtgtt 2580
ttatacatat taccttattt aaagctgctg tttcattatc gtagtctcgt gaactgtggg 2640
tagtgatgac agtgaaaaat ggagaccacc agcacaatcc aggctgttgt agcacatata 2700
gcctttttcac catttttagt tagtcagaaa attagagacc tttatagtcc aaatagctac 2760
tagtatgatg atagtgtata aattttcagt gtgtgactcc tacaactcct ctgctctac 2820
tgtgcatttg aatagttaga tagcattttt aggaaagtcc tcactatttt actttgcatg 2880
atttctgat caagcgagcc aaaagcacag taaatgacag agcagaaatc ttgatctgga 2940
aaggggagatt tggacatat cttctggaag aagtgtcttc tagatgctaa ttaacaggca 3000
aaaacgtaat aaagactaat tttgtagagt attgttgctt tacggttggt gccagtgtgg 3060
ctcagtaatt gcataactga gtatgttggg tcttctctag tttgatctat tagaagtaag 3120
ttctccggcc gggcgtgggt gctcacgcct gtaattccag cactttggga ggtcgaggtc 3180
aggagatcaa gaccatcctg gctaacatgg tgaaaccccg tctctactaa aaatacaaaa 3240
aattagctga gtgtggtggc gggcacctgt agtcccagct actcgggagg ctgaggcagg 3300
agaatggtgc gaacctggga ggcggagctt gcagtgagcc gagatggcgc cactgcactc 3360
cagcctgggt gacagagcga gactccgtct caaaaaaaaa aaaaaaaaaa aaaaa 3415

```

&lt;210&gt; 30

&lt;211&gt; 55

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 30

```

Met Phe Met Lys Ile Tyr Thr Ser Ser Val Pro Leu Phe Arg Gly Met
  1             5             10             15

```

```

Leu Ser Cys Leu Ala Leu Ser Cys Trp Cys Ser Leu Ser Ala Pro Leu
      20             25             30

```

```

Ser Ser Trp Val Asn Gly Asp Cys Gly Ser Cys Leu Ser Ile Ser Asp
      35             40             45

```

```

Ser Ser Asn Asp Thr Gly Lys
      50             55

```

&lt;210&gt; 31

&lt;211&gt; 2967

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 31

```

gatgaagtgg agcagtgagt gtgagcctca acatagttcc agaactctcc atccggacta 60
gttattgagc atctgcctct catatcacca gtggccatct gaggtgttcc cctggctctg 120
aaggggtagg cagcatggcc aggtgcttca gcctggtgtt gcttctcact tccatctgga 180
ccacgaggct cctggtccaa ggctctttgc gtgcagaaga gctttccatc caggtgtcat 240
gcagaattat ggggatcacc cttgtgagca aaaaggcgaa ccagcagctg aatttcacag 300
aagctaagga ggctgttagg ctgctgggac taagtttggc cggcaaggac caagttgaaa 360
cagccttgaa agctagcttt gaaacttgca gctatggctg ggttgagat ggattcgtgg 420
tcattctctag gattagccca aaccccaagt gtgggaaaaa tggggtgggt gtccgtgattt 480
ggaaggttcc agtgagccga cagtttgcag cctattgtta caactcatct gatacttggg 540
ctaactcgtg cattccagaa attatcacca ccaagatcc catattcaac actcaaactg 600
caacacaaac aacagaattt attgtcagt acagtaccta ctcggtggca tccccttact 660
ctacaatacc tgcctcact actactctc ctgctccagc ttccacttct attccacgga 720
gaaaaaaatt gatttgtgtc acagaagttt ttatggaaac tagcaccatg tctacagaaa 780
ctgaaccatt tgttgaaaat aaagcagcat tcaagaatga agctgctggg tttggaggta 840

```

```

aatggtttct gtgtctatgt tacgtgtgta tgtatgttca tgtataacag ataattcata 900
gaatcctgcc tgtctttaga tgaggcgctc aggacctgtc aaatactaata aaaagcttaa 960
cacaactca actttcatga agtcatagaa tctgagagct gtaaagaacc ttagtgatta 1020
agctaaaaag tgtaagtata aaagaatttc actgttgaaa actcaagact tttaaaaata 1080
aatcactatc tgttctctatg cataagttct ttttaaaata tataccctaa aggcctatga 1140
ctttattcta gtcattctctt tctcaggcag gcttgttgac tagagtaggt ctcaaggcca 1200
tggaggagaa atgaactctg tcatattgca actgatagaa ttgagagtgt gagaagcaag 1260
gccaatgatt ctacaagcat gaagcaaaga agctgggtgtg ggcccaatat cataaagggc 1320
tacagagaag gtttttctcc tgtactgtct actgtctcaa tcacctggat catgcaatga 1380
gcattacaca caatgtttaa gtgaaggacc aacaggcctt gtctctgttc cttgcagggtg 1440
tccccacggc tctgctagtgt cttgctctcc tcttctttgg tgctgcagct ggtcttggat 1500
tttgctatgt caaaaggatg gtgaaggcct tcccttttac aaacaagaat cagcagaagg 1560
aaatgatcga aaccaagta gtaaaggagg agaaggccaa tgatagcaac cctaattgagg 1620
aatcaaagaa aactgataaa aaccacagaag agtccaagag tccaagcaaa actaccgtgc 1680
gatgcctgga agctgaagtt tagatgagac agaaatgagg agacacacct gaggctggtt 1740
tctttcatgc tctttacctt gcccagctg gggaaatcaa aaggggccaaa gaaccaaaga 1800
agaaagtcca cccttggttc ctaactggaa tcagctcagg actgccattg gactattggag 1860
tgcaccaaag agaattgccct tctccttatt gtaaccctgt ctggatccta tctcctacc 1920
tccaaagctt cccacggcct ttctagcctg gctatgtctt aataatatcc cactgggaga 1980
aaggagtttt gcaaagtga aggacctaaa acatctcatc agtatccagt ggtaaaaagg 2040
cctcctggct gtctgaggct aggtgggttg aaagccaagg agtcaactgag accaaggctt 2100
tctctactga ttccgcagct cagacccttt cttcagctct gaaagagaaa cacgtatccc 2160
acctgacatg tcttctctgag cccggtaaga gcaaaagaat ggcagaaaag tttagccctt 2220
gaaagccatg gagattctca taacttgaga cctaattctt gtaaagctaa aataaagaaa 2280
tagaacaagg ctgaggatac gacagtacac tgtcagcagg gactgtaaac acagacaggg 2340
tcaaagtgtt ttctctgaac acattgagtt ggaatcactg tttagaacac acacacttac 2400
tttttctggt ctctaccact gctgatattt tctctaggaa atatactttt acaagtaaca 2460
aaaaataaaa ctcttataaa tttctatttt tatctgagtt acagaaatga ttactaagga 2520
agattactca gtaatttgtt taaaaagtaa taaaattcaa caaacatttg ctgaatagct 2580
actatatgtc aagtgtgtg caaggtatta cactctgtaa ttgaatatta ttcctcaaaa 2640
aattgcacat agtagaacgc tatctgggaa gctatttttt tcagttttga tatttctagc 2700
ttatctactt ccaaactaat ttttattttt gctgagacta atcttattca ttttctctaa 2760
tatggcaacc attataacct taatttatta ttaacatacc taagaagtac attgttacct 2820
ctatatacca aagcacattt taaaagtgcc attaacaat gtatcactag cctcctttt 2880
tccaacaaga agggactgag agatgcagaa atatttgtga caaaaaatta aagcatttag 2940
aaaaaaaaa aaaaaaaaaa aaaaaaa 2967

```

&lt;210&gt; 32

&lt;211&gt; 250

&lt;212&gt; -PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 32

```

Met Ala Arg Cys Phe Ser Leu Val Leu Leu Leu Thr Ser Ile Trp Thr
  1             5             10             15

```

```

Thr Arg Leu Leu Val Gln Gly Ser Leu Arg Ala Glu Glu Leu Ser Ile
      20             25             30

```

```

Gln Val Ser Cys Arg Ile Met Gly Ile Thr Leu Val Ser Lys Lys Ala
      35             40             45

```

```

Asn Gln Gln Leu Asn Phe Thr Glu Ala Lys Glu Ala Cys Arg Leu Leu
      50             55             60

```

```

Gly Leu Ser Leu Ala Gly Lys Asp Gln Val Glu Thr Ala Leu Lys Ala
      65             70             75             80

```

```

Ser Phe Glu Thr Cys Ser Tyr Gly Trp Val Gly Asp Gly Phe Val Val
      85             90             95

```

Ile Ser Arg Ile Ser Pro Asn Pro Lys Cys Gly Lys Asn Gly Val Gly  
 100 105 110  
 Val Leu Ile Trp Lys Val Pro Val Ser Arg Gln Phe Ala Ala Tyr Cys  
 115 120 125  
 Tyr Asn Ser Ser Asp Thr Trp Thr Asn Ser Cys Ile Pro Glu Ile Ile  
 130 135 140  
 Thr Thr Lys Asp Pro Ile Phe Asn Thr Gln Thr Ala Thr Gln Thr Thr  
 145 150 155 160  
 Glu Phe Ile Val Ser Asp Ser Thr Tyr Ser Val Ala Ser Pro Tyr Ser  
 165 170 175  
 Thr Ile Pro Ala Pro Thr Thr Thr Pro Pro Ala Pro Ala Ser Thr Ser  
 180 185 190  
 Ile Pro Arg Arg Lys Lys Leu Ile Cys Val Thr Glu Val Phe Met Glu  
 195 200 205  
 Thr Ser Thr Met Ser Thr Glu Thr Glu Pro Phe Val Glu Asn Lys Ala  
 210 215 220  
 Ala Phe Lys Asn Glu Ala Ala Gly Phe Gly Gly Lys Trp Phe Leu Cys  
 225 230 235 240  
 Leu Cys Tyr Val Cys Met Tyr Val His Val  
 245 250

&lt;210&gt; 33

&lt;211&gt; 2926

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 33

tttagtctgga aatatgctag tagtaatcat tggaaaagct gatgaagcta caactgtggt 60  
 acgtgtgtgt gattttatctt cttattggca tttctctgct ggcagttgga actgacagtg 120  
 atgggaatgt ggcaactgaa tgttttctct ctttcttggt tccttccatt tttatctcaa 180  
 cttttctagt gtgtgtgtccc cttttctaaa tggcattgac aaactcatca agatccagag 240  
 ataaacagtc catcgtttaa gaatatttta ttttgcattg agaggaaaaga attaaccaaa 300  
 gacgtttctg ctctgatact ttctaagaa aacagactaa ttatatataa gtggtatcaa 360  
 taggtctgct tataagtatt gtcctatgta gagatttact ttttggtttt tacaagcata 420  
 gtttgcgctt ttagatgtct ctcaggtgaa aaatagaagt tggcagccct gagcaagcac 480  
 ctgtgacaca gggattttta ttttagtttt aaagtatgat atagagaatg aggttttttg 540  
 tttgatttgg tttggttggt ttttttagtt ctttattttt tatttggggg ttctcttagt 600  
 gatggcatat ttcaggtttc agtgacatag atcaagtaat tttaaaccga tttggttatt 660  
 ctatctaaga tgagggacat ggctattaag gtcaagccaa actatactaa aagtagtata 720  
 gggcagcagt taaaattatc ttttgaaaat caagtattac ctggtttctt ctgtcacaac 780  
 agaatactgt gttacctagt cagtcacagt tgccttgcc ttcccttggt agtccctgga 840  
 ggtacttgag tggacacagaa ggcagaatta gcaacagctc aatcacttta ggtagcattt 900  
 ctctgaatt ctgctgcaa atcctcaggg tctatggatt ggttgaaata gtaaaatcac 960  
 acatagtgat ttcttgaca gcgtagaggc gtttaatttt agtgatatag gaacaaagaa 1020  
 ggaaaactgg gtcataataa atttaacctt aaaacacaga ttataaatat ctttgaatgt 1080  
 taagccctag gcagacagac actctcattc ttaactctgg tatgcaagct ggccttgctg 1140  
 acagtggcat agttacatta actgtttttt aagttgcatg ttgtgatgtg aatgtttttt 1200  
 gtaaaagttg ccaagttaat cgagtcctaa ataagtagcc aaatggtaac tgagaccac 1260  
 tgtagacatt actcgcttta tattatttgc tttcaaatac taaaacctca gctgctaatt 1320

```

cagcgctcta ggtgtacagt gtctccttcc cttggctcct agcaatgtag ggaacttgag 1380
gttcttcagt atcttcagag attccttcca ggccacatat ttttatgcta atacttctat 1440
agttgcattc aatctacaag gagaaagtgt acattggtgc atgttttata aaccagaca 1500
cagatagcta aaaccataat tagtttttct ataccagag ctttgaatta aaaacagatg 1560
cttttttttt ccccaagagc agacagtctt acattcctaa cattaggaat tgggtgatacg 1620
agaagcagga tccagaattc agaattgatt agtgagactc ggtgaaaaaa atgcattttc 1680
ccctggctgt ttgaaaattt acttatttgc agataagtct agatttagtc ttggagatca 1740
aagtccttta tattttaaaa acttatttct tatattgatc aaacatggca tatgttagag 1800
aaccacttct tctgtcatgt ttatgtattt tggaattaag ttgtttgcat tcactttcaa 1860
aatctgcccc tttctgttta tgtgcactta ccacagatgt gtcgggactt tgcctcaggg 1920
gagagggtact ttagcacctg tgtcactgag gagatggagt ggttgacaag tactgttgcg 1980
ctgtgtaact tggggttttg ccctgtggac aatatattag cagaatgata ccacacaaaa 2040
gtattacagg attaaggcat gtaacttcta tggtagtctt tatgtatcag cgtataacca 2100
agttcgaaaa ccacaggtgc atttttagac ctttacttag agaactaaag gcagttccaa 2160
ccctcagccc atatggcggg attaatgcat gaaaaccctc agagggtgtt gggacatcct 2220
acttccctgt cctcaccagc tggaactctg gtgtgtgcct tgaggataag gaagtagagt 2280
ggaaactcat cctatcattg agtattctca atattttggc cttccctctg gaattatgag 2340
aaatttaaca aagtcctcagg aacctttaga atccattgtc caacactgct agaaaaactg 2400
taggaggtac atggagaatt cctatagttc ttaggtaagt gcaagacatg gcacagggat 2460
ccctatccac ataaagggga atctggatgc tgcacacctc aattctgaga aatccctgac 2520
tgaacttgga attatgacag taaagttttc gtcctttagt tttctagagc agtcacaga 2580
aatttttaaaa agtaaaacaa ggccaggcgc agtggctcat gcctgtaate ccagctcttc 2640
gggaggctga ggcgggcaga tcacgaggtg aggagatcga gaccatcctg gctaacaggg 2700
tgaacccccg tctctactga aaatacaaaa aattagctgg gcattggtggc gggcgctgt 2760
agtcacagat gctcaggagg ctgaggcagg agaatcgctt gaacctggga ggcagaagat 2820
tacagtaagc caagatcgcc ccaactgcact ccagcctggg cgacagagtg agactccgtc 2880
tcaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 2926

```

&lt;210&gt; 34

&lt;211&gt; 55

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

```

Met Lys Leu Gln Leu Trp Tyr Val Cys Val Ile Leu Phe Leu Ile Gly
  1              5              10              15

```

```

Ile Ser Leu Leu Ala Val Gly Thr Asp Ser Asp Gly Asn Val Ala Thr
      20              25              30

```

```

Glu Cys Phe Leu Ser Phe Leu Val Pro Ser Ile Phe Ile Ser Thr Phe
      35              40              45

```

```

Leu Val Cys Cys Pro Leu Phe
      50              55

```

&lt;210&gt; 35

&lt;211&gt; 3283

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 35

```

aaaagctgaa tcagaaatga agttaattaa acacatgtac acctgccaca aatttgtctc 60
tgcactttta cttgctgttg ttcaattcac agtgtccaag agaaaaataa attatttctc 120
agaaaaagtt catccattct tagtacttag aacaatcaaa atacattatg atgaattgta 180
tttaatatat gtctattttg atactggatt taaagaacat ttaagggaag aatgttctct 240
tgatttgcta aactaacagt aggaaccaag tcaatgcgta ttgaatataa caaaattatt 300
cctttatgtt tccagaagca gcatcaaaaa taattgtatt tctaagtcac atacagcaaa 360
ccattatata tttgcatttc atatttttaa aagtactcta taggaaattt ttaatgtatc 420

```

```

ctacatacac atagctgtat gaaaatagtt tattcatgga ggaattttcc tgtagaatac 480
ttatatatct gccttacaat taaaaccaa catgtattac tttttttgtt acagaagctt 540
tatatagaaa atacagttaa attcatcaa ataaaaaatt ttaattctga gtctgggaaa 600
tgtatacaaaa ggacttcatt gagccattct cttgactttt gaatatattt gaaaattttc 660
gtatgaattt tcagtaacca catgtattcc tttcagatca tactagaatg cagttcactt 720
tcctatttga taaactcatg ctcatTTTTt tgaatttgat acggtatttc caaaatatta 780
tctgattgga aataataaac atttgcttaa actgtaacga aggccaaagt gctgggatta 840
caggcatgag ccaccgcgcc agggccaaaa ctaaaagtaa tttttatgtt caatacctta 900
ttgaaacatc ttgactctca taaataaaca ttgccaaatt ttaacatact gtctaaagtt 960
agcaatgaat aagcatatgt ttcttaatct gtggcgggtg aaaaatttgt tatatttgtt 1020
cagtcaataa gatttttagt gtaaatgtgg gtagattcat gaaacgatca aatttctgga 1080
caattaacat tgcacaaaat taatggcata aaacatattt acaaagaact aaacatcaaa 1140
tgtgaggtca tctgagcata tagtagacct tactctttct ttcactttct ataaacatc 1200
tgcttaggga aaacctagac ttagaaataa taaaaatgca gagaccaaag actggatttt 1260
tttaaatgaa cttttcttag aaaaaagatt ctgcaaaatg tcagtgtattg aatgtatttt 1320
taacgtttga gcacagcatg gcacaataaa acaaatgtca gccatctgac ttgggttgta 1380
ctttacaagt ttatgactat tgcaagagga caaaatactg aggttgctga aggtaaagt 1440
ttttgtctcg tgggccttta gcctttctcg ttttgatctt cctctttaat ccatagggtg 1500
tctgatgggt ggagacctca cgtctgtgac aggcaccagt gcaggtaagc ccagactcca 1560
tactgcagta tgtatcattg ttctgagttc ttctactagg tgatttgtag gacaagtct 1620
tttgaacaaa taaagttaag agatctgtga gaggcattga aaaatgacac caagaatttt 1680
taaagcactt ttctgaagca ggagtggta aatagtaact tagaagaaaa ctatttatta 1740
actagaacct tccttaataa agcttcaact ggtagcaga gaattaagggt ctctgatcca 1800
atcaacttaa tcctgggata catccttgga cacaagagta cctttagcaa accatacttt 1860
ttgtgtctc tgtaaatcat ttacacattaa atatgaatca tttccagcc aaagaagtat 1920
tgaacttcta tgcatttcta aagaagagta cataaagaaa aggaggggtt aagattttta 1980
ttattcaaca atactgttaa taaaataaaa tcacattggc attaaaatga gctcttagac 2040
aacatgaata taatgcattc ccaaaagacc ctaaaatata tttcagctaa gaacttctta 2100
cactgtgacc aataattgtc tcagtccatc ttctccaggt ttccgtgtta atttgtaata 2160
ttgagattat tgagatatcg atgtcatttt caatgcagag gcaagatagc acaatacttt 2220
caaatgccaa ctgcagttca ctattggcat aacaagtaac catggttaatt tctgcagctg 2280
gaacatgcta tacattttaa atagtaggag gcaggcatga atgacacaag gttatgtttg 2340
aaatgtcaaa aaaaatcaag tttattgaaa tagacatgga tagcctaatt aaaaattttt 2400
gccagggtca agtagaatag cctgctaatt ataagaaaga agataaagaa aattggttta 2460
caaatacatt tttataaata tatctatttg agggtagagt ttctctgttc tgatgctggc 2520
tcttcactaa ttcactgaat aatgaagaaa aagtattaga cctacacagt gattagcaaa 2580
caagatgaat aacactagct tcctatttta tatatctttt gaaaattagg tcgagccaaa 2640
ttagtgtgta aactcacgta gtattttgct ggacatggaa atgaaatttt ttccctcaaa 2700
aaatactttg tggtttgtag cctcaaaatg agagaaccag atcaaatcca gaccaatgtt 2760
tcttggtacc aagtctctac caaacactat gattagaaaag aatggtgagg actgagaatt 2820
gggtgggtcac aaagaccaag atataatata agaaaaatgg tttagatgtt atgatgtttt 2880
acccaatata tcactgcacc gatttgcttg ccattatgga aacagctagt agtggttctg 2940
aaagtcatta aaagtagtaa aaaagggcgc ggcggtgtg ctacgcctg taatccagc 3000
actttgggag gccgagggcg gcggatcacg aggtcaggag atcgagacca tcccggtcaa 3060
aacggtgaaa ccccgctctc actaaaaata caaaaaaatt agccgggctg agtggcgggc 3120
gcctgtagtc ccagctactt gggaggctga ggcaggagaa tggcgtgaac ccgggaggca 3180
gagcttgtag tgagccgaga tcccgccact gcactccagc ctgggcgaca gagcgagact 3240
ccgtctcaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 3283

```

<210> 36

<211> 79

<212> PRT

<213> Homo sapiens

<400> 36

Met Lys Leu Ile Lys His Met Tyr Thr Cys His Lys Phe Val Ser Ala  
1 5 10 15

Leu Leu Leu Ala Val Val Gln Phe Thr Val Ser Lys Arg Lys Ile Asn  
20 25 30

Tyr Phe Ser Glu Lys Val His Pro Phe Leu Val Leu Arg Thr Ile Lys  
 35 40 45

Ile His Tyr Asp Glu Leu Tyr Leu Ile Tyr Val Tyr Phe Asp Thr Gly  
 50 55 60

Phe Lys Glu His Leu Arg Glu Glu Cys Ser Leu Asp Leu Leu Asn  
 65 70 75

<210> 37  
 <211> 2248  
 <212> DNA  
 <213> Homo sapiens

<400> 37  
 tgatgttcct cgctgtctc tgcctggaaa actggtcttc ccaagctcca ctggcagcca 60  
 cttctccatg ttgggcatcg gagacatcgt tatgcctggg ctccactat gctttgtcct 120  
 tcgctatgac aactacaaaa agcaagccag tggggactcc tgtggggccc ctggacctgc 180  
 caacatctcc gggcgcatgc agaaggtctc ctactttcac tgcacctca tcggatactt 240  
 tctaggcctg ctactgcta ctgtggcgtc tcgcatcac cggggcgcgc agcccgccct 300  
 tctctatttg gtgccattta ctttattgcc actcctcacg atggcctatt taaagggcga 360  
 cctccggcgg atgtggtctg agcctttcca ctccaagtc agcagctccc gattcctgga 420  
 agtatgatgg atcacgtgga aagtgaccag atggcgtca tagtctttt ctctcaactc 480  
 atggtttgtt tcctcttaga gctggcctgg tactcagaaa tgtacctgtg ttaaggaac 540  
 tgcctgtgta ctggatttgg catttaaagg gagctcgttt gcaggagaga ggtgctggag 600  
 ccctgttttg ttctctctct tcctgctgat gtagaggtgg ggcccttcc aagagggaca 660  
 ggctctctcc cagcgcgctt tcctcccacg ttttatgga tctgcaccag actgttacct 720  
 tctgggggag atggagattt gactgtttaa aaactgaaaa cagcgaggag tctttctaga 780  
 acttttgaa actaaaagga tgaaaaaat tagcaaacg aagtttcttc aatgacctct 840  
 cgagaacttt gggaccagtt tcctataggg gactcagttt cagagaactg agacagaagc 900  
 tcttctgtcg ttatattctt ctttctttt tttggattta ttaaataatt tctgtggtgt 960  
 gaagtgactt attaaatcca cagacattga gtgacttctt acaacatcca cataagaatt 1020  
 tgttgaatg agttcatgtc caccagatg ttgtgttggc agtgaacaag ggcacggttt 1080  
 ttatacatac gtacatata atataaaac acacacatag atatataga ataaacaaaa 1140  
 atgaaatcct gctaagatca cgctgtgtag ctgacagggg cttgctgtcg ttttgagcat 1200  
 gtcgagcagt ttactgtggc ttcttgtat atggataagc tgcgtctctt ccccttcaca 1260  
 actgaccccg cagttacaaa ctagtatagc atttgtgctg attgatgata gactcatgga 1320  
 cttcaggagc cttactttgg ttttgatcag tgtagcaaat tagggatgaa gagttcaaac 1380  
 cttttggccc tttctttctt ttctaggctt ctccctcgca ggggtgtccg tagtttcttc 1440  
 tcgagccaat gcatgtatta tagcagcagg tgtcttctgt ctttctcctc atagtaacgt 1500  
 actacttgta aatacatctt tctattttct atttttttgt attttttttt ttgacatttt 1560  
 gtttcatagg tgtgctgtat attttccatg ccctcactcc tttaagaaaa aaaaaaagga 1620  
 aaaaagcaac acaatcctgt cttgctgtt gtgattatag tcttggttta cctgtggtga 1680  
 caaccgggtg ttggggacac atgtcaaatg cccctctgag atgggccccta aattccagta 1740  
 actggggaaa gaaccagctg ctgtgtcctg agagcctggc cctgtgctgt ggtctctgct 1800  
 gcaagcccag atttctggga gtaactagtg ttaggtctgc tgacctttac ctaagcagcc 1860  
 cctgcctggg aagaaggtgc ccattgttca gaggcaaaga gaagcctgcg gttggcatga 1920  
 ggatgcctga caacaaaggc tggagaaggg ccctgagttc cagcctctcc ccaaggggtcc 1980  
 ccgccccagt ggtgctctct gtcttgacct gtgtaatgaa ttagtgtgct gtgtcactgt 2040  
 ggcttgaagt cactgtggat cgagctcaca gggggcacc atcctgttgt caacagagtc 2100  
 tgaagcagtc aggtgtttgg attcctctgt tgttgcctc aattcctgct gaggggtttc 2160  
 tgggggtttt tttttaataa atgactcctt tgtaaaaaa aaaaaaaaaa aaaaaaaaaa 2220  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2248

<210> 38  
 <211> 119  
 <212> PRT  
 <213> Homo sapiens



&lt;400&gt; 38

Met Leu Gly Ile Gly Asp Ile Val Met Pro Gly Leu Leu Leu Cys Phe  
 1 5 10 15

Val Leu Arg Tyr Asp Asn Tyr Lys Lys Gln Ala Ser Gly Asp Ser Cys  
 20 25 30

Gly Ala Pro Gly Pro Ala Asn Ile Ser Gly Arg Met Gln Lys Val Ser  
 35 40 45

Tyr Phe His Cys Thr Leu Ile Gly Tyr Phe Val Gly Leu Leu Thr Ala  
 50 55 60

Thr Val Ala Ser Arg Ile His Arg Ala Ala Gln Pro Ala Leu Leu Tyr  
 65 70 75 80

Leu Val Pro Phe Thr Leu Leu Pro Leu Leu Thr Met Ala Tyr Leu Lys  
 85 90 95

Gly Asp Leu Arg Arg Met Trp Ser Glu Pro Phe His Ser Lys Ser Ser  
 100 105 110

Ser Ser Arg Phe Leu Glu Val  
 115

&lt;210&gt; 39

&lt;211&gt; 931

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 39

tggggccgct tgggctgttg gtctccagag cagggccact gggcactctg tgatggggga 60  
 gcctttgtct gaaagcacag cccctcgc cttcctctcc ccattggctt cccttcattg 120  
 gcattaatct gggcaccagc tctctccata gcagtgaact ccctcaccac tctcatctct 180  
 cagccttgcc ttttcttctt gacactgtcg cccctcctc tcaggagaca ctgccgaggg 240  
 ccacctggca gaaggctgag ttaggcagca gggccgggag cgtctgccct ccacaggggtg 300  
 ggggacagat aggctaagcg actcccagct tgctaccctc agtggccagt gtgggcgttg 360  
 gcggtttggg gcgcttggtt ggtgggtggc actgcatccc ttaatttatt tctctgctgt 420  
 ttctgttctt gagaaatttg ggtggggagt cctacacaga ggctgcccc accctcacct 480  
 gagttgtaca tttttttgtg atgggtttta ttttttatta ttttatttta tttttttttt 540  
 tttttgattt atgatgactc caccctctt catcaccccc gctcccaggc caggctcagc 600  
 gattaagccg agcccttgcg tcctaggaag gggccttgcc aacctcagcc ctctgcccc 660  
 acactcctac tgcggctcag accaagggtt cccctcctt cccttcccc ctctgccc 720  
 atggaacagc ccgggtgctc tgagggggct gggagggcat ggcttggtc ccaaaggggg 780  
 taggggccc gggcaccag gcaaggtggc cctccccgt ctagccccct cctcccaac 840  
 cctgcactta gtttctctc tggatcaaac acgtaataaa gagaatgttt ggaatctgaa 900  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 931

&lt;210&gt; 40

&lt;211&gt; 53

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 40

Met Ala Ser Pro Ser Leu Ala Leu Ile Trp Ala Pro Ala Leu Ser Ile  
 1 5 10 15

Ala Val Thr Ser Leu Thr Thr Leu Ile Ser Gln Pro Cys Leu Phe Phe

20

25

30

Leu Thr Leu Ser Pro Pro Pro Leu Arg Arg His Cys Arg Gly Pro Pro  
 35 40 45

Gly Arg Arg Leu Ser  
 50

&lt;210&gt; 41

&lt;211&gt; 3625

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 41

tggctaggct ggtcttgaat tectgacctt gtgatccacc cacctcggcc tcccaaagtg 60  
 ctgggattgc aggcattgagc caccacgccc ggccgttgga acatattttt atttaattta 120  
 attaatattt ttatttattt ttgagatgga gtcttgctct gtcgcccagg caggagtga 180  
 gtggcgcgat ctcggtcac tgcaagctcc acctcctggg ttcacacat tctcctgcct 240  
 cggcctccca agtagctggg actactcgcc accactccag ctaatttttt tgtattttta 300  
 gtgagacagg ggttttacca tgttgccaa gctggtctcg atctcttgac cttgtgatcc 360  
 acccgctca gcctcccaa gtgctgggat tacaggaatg agccaccgct cccggctgct 420  
 ggaattttta ataagcacgt gagctgggtg gatgagaacc ccattgtagga gtttttaccc 480  
 atttcttctt cggcagcctt gttgagtgc tactgtgcgg caggctttgc tgattcagca 540  
 gtgagtccac caagcaagcc ctgcccgcag gatgagtcca gcaggggagg tggcagcgag 600  
 gcagccttca ggccacagcg gctggcctct gtccttggtg ctgaggctgc gctggtggga 660  
 aaaatcatcc agcatcaaaa acccccacc gtcaaaaagc ttgtgtctta gcaagtggga 720  
 gacagctgag tgggcccctcc ctgagttagt gattcctaac ctagtctctc gacctttctc 780  
 tttgtttgat accaagttta ttttttttaa tttttttatt atctatttat ttatttctgg 840  
 gacggagtca cgtctgtggt cccagactgg atttcagtgg tgtgatctcg gctcactgcy 900  
 gctctgcct cctgggctca agcgattctc ctgccttgcc ctcccagta gctgggatta 960  
 taggcgtgca ctaccatgcc cagctaattt ttgtattttt ggtagagaca gggtttcacc 1020  
 atgttggtcca ggatggtctt gatctgtga cctcgtgatc caccacctc ggccctccga 1080  
 agagctggga ttatgggaat gagccaccgc acctggcct cggcctccc aagagctggg 1140  
 attacgggaa tgagcgaccg cgcccggccc cagattttcc tttggttgcc atctttgtat 1200  
 acttcttaac tctcgtttg atcaaaagat agttttggct gggcatggtg gccacgcct 1260  
 gtggtcccgg tacttcggga ggccgaggtg ggaagatggc ttgaggccag gattccata 1320  
 ccagcctggg ctgcagagca agacctgtc ttacaaaaa ataagaaaac ttagctgagt 1380  
 gtggtgacct acgcctgtgg ctgcggctac tcgggaggct caggcaaaag gatcccagga 1440  
 gtttgaggct gcagttagcc gtgatcgct cactgcactc cagcttggtg gacaggcatg 1500  
 cgtggagtgg cagtgtaac ggctggtgtc tcgactgtt ggctgtgaa ggtacgtgaa 1560  
 gctgaaagcc tggaatggct ggaaaggggt catcaggcag cgggcccctg ctgctggggc 1620  
 tgcgtggtggc cgtagccact gtccacctgg tcatctgtcc ctacaccaa gtggaggaga 1680  
 gcttcaacct gcaggccaca catgacctgc tctaccactg gcaagacctg gagcagtacg 1740  
 accatcttga gttccccgga gtcgtcccca ggacgttctt cgggcccagt gtgatcgag 1800  
 tgttctccag ccccgcggtt tacgtgcttt cgctgttaga aatgtccaag ttttactctc 1860  
 agctaatagt tagaggagtg cttggactcg cggtgatttt tggactctgg acgttacaaa 1920  
 aggaagttag acggcacttc ggggccatgg tggccaccat gttctgctgg gtgacggcca 1980  
 tgcagtcca cctgatgttc tactgcacgc ggacactgcc caatgtgctg gccctgcctg 2040  
 tagtcctgct ggcctcgcg gcctggctgc ggcacagtg ggcctgctt atctggctgt 2100  
 cagccttcgc catcatcgtg ttcaggggtg agctgtgctt gttcctgggc ctctgctgc 2160  
 tgctggcctt gggcaaccga aaggtttctg tagtcagagc ccttcgccac gccgtcccgg 2220  
 cagggaacct ctgttttaga ctgacgggtt ctgtggactc ttatttttgg cggcagctca 2280  
 cttggccgga aggaaggtg ctttggtaac aactgtctt gaacaaaagc tccaactggg 2340  
 ggacctcccc gctgctgtgg tacttctact cagccctgcc ccgcgccctg ggctgcagcc 2400  
 tgctcttcat cccctgggc ttggtagaca gaaggacgca cgcgccgacg gtgctggcac 2460  
 tgggcttcat ggactctac tccctcctgc cacacaagga gctacgttct atcatctatg 2520  
 ccttcccctat gctcaacatc acggctgcca gaggctgtc ctacctgtg aataactata 2580  
 aaaagtcttg gctgtacaaa gcgggggtctc tgcttggtat cggacacctc gtggtgaatg 2640  
 ccgctactc agccacggcc ctgtatgtgt ccattttcaa ctaccagggt ggcgtcgcaa 2700

```

tgcagaggct gcaccagctg gtgccccccc agacagacgt ccttctgcac attgacgtgg 2760
cagccgcccc gacaggtgtg tctcggtttc tccaagtcaa cagcgcttgg aggtacgaca 2820
agagggagga tgtgcagccg gggacaggca tgctggcata cacacacatc ctcattggagg 2880
cggcccttgg gctcctggcc ctctacaggg acacacaccg ggtcctggcc agcgtcgtgg 2940
ggaccacagg tgtgagtctg aacctgaccc aactgcccc cttcaacgtc cacctgcaga 3000
caaagctggg gcttctggag aggtctcccc ggccgtcctg agggggacca ggcagccctc 3060
agcagccaca ggccttccag gagctgttat cactaccagt ttctggcaca attccagcac 3120
aattatgaca attcagagaa gcaagtcaaa ggactgggca cctgcctctg acagacacca 3180
gaccaggtct tgttggcacc ccgggagcca ctgcccaggg tgatggtggc cagctcaggg 3240
cttctctggg gtgactgtcg cccagaccag gtgccattca tgactaatca ggagcagcgg 3300
gtcaccacag gcacctgtct gccaggaggc cagctgtgtc ctgcccaccc agggggagct 3360
gtattttggc agcaccacac gcttgcctgc cgagggcctc ttggggcacc taagacagca 3420
ccccctctca ggggagacca tgggtggcccc ggccgcaccc cccacacctg gtgccaccac 3480
tgcaactttt gtattcacag gcatcccatc tccatcacag ataaaatctt aggagataaa 3540
cacattcaaa aaggaaatgag ataaaagaa taaggcaata aatgttgatt ggaacctctc 3600
aaaaaaaaaa aaaaaaaaaa aaaaaa 3625

```

&lt;210&gt; 42

&lt;211&gt; 488

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 42

```

Met Ala Gly Lys Gly Ser Ser Gly Arg Arg Pro Leu Leu Leu Gly Leu
  1             5             10             15

Leu Val Ala Val Ala Thr Val His Leu Val Ile Cys Pro Tyr Thr Lys
             20             25             30

Val Glu Glu Ser Phe Asn Leu Gln Ala Thr His Asp Leu Leu Tyr His
             35             40             45

Trp Gln Asp Leu Glu Gln Tyr Asp His Leu Glu Phe Pro Gly Val Val
             50             55             60

Pro Arg Thr Phe Leu Gly Pro Val Val Ile Ala Val Phe Ser Ser Pro
             65             70             75             80

Ala Val Tyr Val Leu Ser Leu Leu Glu Met Ser Lys Phe Tyr Ser Gln
             85             90             95

Leu Ile Val Arg Gly Val Leu Gly Leu Gly Val Ile Phe Gly Leu Trp
             100            105            110

Thr Leu Gln Lys Glu Val Arg Arg His Phe Gly Ala Met Val Ala Thr
             115            120            125

Met Phe Cys Trp Val Thr Ala Met Gln Phe His Leu Met Phe Tyr Cys
             130            135            140

Thr Arg Thr Leu Pro Asn Val Leu Ala Leu Pro Val Val Leu Leu Ala
             145            150            155            160

Leu Ala Ala Trp Leu Arg His Glu Trp Ala Arg Phe Ile Trp Leu Ser
             165            170            175

Ala Phe Ala Ile Ile Val Phe Arg Val Glu Leu Cys Leu Phe Leu Gly
             180            185            190

Leu Leu Leu Leu Leu Ala Leu Gly Asn Arg Lys Val Ser Val Val Arg

```

195	200	205
Ala Leu Arg His Ala Val Pro Ala Gly Ile Leu Cys Leu Gly Leu Thr 210 215 220		
Val Ala Val Asp Ser Tyr Phe Trp Arg Gln Leu Thr Trp Pro Glu Gly 225 230 235 240		
Lys Val Leu Trp Tyr Asn Thr Val Leu Asn Lys Ser Ser Asn Trp Gly 245 250 255		
Thr Ser Pro Leu Leu Trp Tyr Phe Tyr Ser Ala Leu Pro Arg Gly Leu 260 265 270		
Gly Cys Ser Leu Leu Phe Ile Pro Leu Gly Leu Val Asp Arg Arg Thr 275 280 285		
His Ala Pro Thr Val Leu Ala Leu Gly Phe Met Ala Leu Tyr Ser Leu 290 295 300		
Leu Pro His Lys Glu Leu Arg Phe Ile Ile Tyr Ala Phe Pro Met Leu 305 310 315 320		
Asn Ile Thr Ala Ala Arg Gly Cys Ser Tyr Leu Leu Asn Asn Tyr Lys 325 330 335		
Lys Ser Trp Leu Tyr Lys Ala Gly Ser Leu Leu Val Ile Gly His Leu 340 345 350		
Val Val Asn Ala Ala Tyr Ser Ala Thr Ala Leu Tyr Val Ser His Phe 355 360 365		
Asn Tyr Pro Gly Gly Val Ala Met Gln Arg Leu His Gln Leu Val Pro 370 375 380		
Pro Gln Thr Asp Val Leu Leu His Ile Asp Val Ala Ala Ala Gln Thr 385 390 395 400		
Gly Val Ser Arg Phe Leu Gln Val Asn Ser Ala Trp Arg Tyr Asp Lys 405 410 415		
Arg Glu Asp Val Gln Pro Gly Thr Gly Met Leu Ala Tyr Thr His Ile 420 425 430		
Leu Met Glu Ala Ala Pro Gly Leu Leu Ala Leu Tyr Arg Asp Thr His 435 440 445		
Arg Val Leu Ala Ser Val Val Gly Thr Thr Gly Val Ser Leu Asn Leu 450 455 460		
Thr Gln Leu Pro Pro Phe Asn Val His Leu Gln Thr Lys Leu Val Leu 465 470 475 480		
Leu Glu Arg Leu Pro Arg Pro Ser 485		

<210> 43  
 <211> 2861  
 <212> DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

```

ttgttttggg gaagtaaaat ccttaaggga ctaaattaat gcttgggtgc attaaaaaga 60
acaaaacatt cccacatgtt ggggtcattg ggagatgccc ggttttgcgg gttttatttg 120
tttaatttta ttctgtgttt tctcttggct ctttgggtct ttcccgggta cactagatgg 180
ctccatccca aggcattctt tcataaaaca gctttccccc accccatata atgggaaaag 240
ggggagaaaat atagccccta gcctaataac ttatcatttg taaaatgact tataaaaaata 300
ttacctcaat ggtaggagac atccagactt gtatatttca gtggaaatac aaaaccactt 360
cagagaccag ggtatctcct ctggaaggat ctaagagaag gtaagacaga ttaggacata 420
gaaaaggagg atggagccag gtgccatggc ttgagcctat aatccgaggc tgagggtggga 480
ggatcacttg agcccaggag tttgagggtg cagtgaagct tgatcacacc actgcactcc 540
agcctgggtg acagagtggg actctgtctc aattaatttt ttttttttaa aggaggagga 600
tctccatggg taagtgtgtt ctaccgcgat gggtagagtt ctgcctctgg tccttctcag 660
ggggcacttt caaccaagagc agtgtaatat tctctgaaag agcaagtacg cttgtgcgcg 720
atccccaacc aatccacagc ctggagtacc tttcaagggt aaagtgcata gccagctcca 780
ttgagacatt ccatttcaaa gcaccgtgct gacagatata aaagtactct agcagggaaa 840
ataatttgtt tgctgtgtaa ggaagaatgt agacaagaca gataaatctg aagggtcatgt 900
ggcatcaggg aaagggcata gctgtgtctt ttgcacccaa tatgaaacat cttctcccaa 960
cactgcttta atggaagtgc taggaaccaa tttagctcag gcatttgact cctacagcag 1020
aagtctctag cctgaccaca gatggtgtgt aatctatcaa acacacccct ggccaagttg 1080
ggtcctatag caccgtgtac tatgtactat tgtaacttct agttccctaa gaggtacctg 1140
ttttcagtaa aaaggggtcc tgagtctgtg gcagggtgaa gagctacccg agaactacct 1200
gagtctcttg caggtagagt cccatttctt atgggacctg tgtgctcctg agaactctta 1260
cttgagacat caaaaagaag cagcaagagc ttctgggaca gagactgctt ggccagcttt 1320
gtaagtaagt ggtgcctcc aatgtgatgt gagtacctgt tgggcagctt cactgtccta 1380
aggatgtctt tctttccacc tcccactgcc cctcccctgc caccatcaa tgatgccttg 1440
gttcagtcac tagaaatctg ttgctttgag ttctgaaata ttttcacctt aaaaaaatg 1500
ctgaaaatac acattctcct gggaagacga taaacagcta gctaagaagc cgagggttcag 1560
tggtggcagc aggaaggaca ctgccacaaa ttttgtctat ttcataattg tcccctagag 1620
ccagccctag caaatgtgtg agttgggagt agttaatagt aaataagact ctgactttac 1680
acaagctaca cattttatac ttttcataaa ccacaagatc tctctagaat tttttctgcc 1740
ttcactaaaa ttggactgta gccaaagatat aaagcaagtc atttggaaac tgccgagtga 1800
gcactgaagc tactttatca tgagatgtgt gttaaagaag ctgcagccca caggagtcca 1860
gggaaggcgg ggaccacaga ggcacagagt ccagcacttg gccgctcatg ggccttcttt 1920
ctgcctcaga ggacgggggc agagaagtga tgaagggaaa tgttcttaga ggaggaaata 1980
tcctttgtcc tgctcagaga gaccagggcc ctaccattag gcatactttc agaagcaacc 2040
tgagaaacag ctatcaatca tattcaaaac cagtacaaga actgctgcct ggtacctgtt 2100
gagtcatttc tatgaaattc catataaaga atgatgataa gtttacacac tgtgcaatct 2160
cacaatctga aaataaagtt gagttggctg tgttttctct gctcttgtca gaacattggg 2220
acaattgggt gttcaaaaac attcatcctc ttactgcaag ttatctggg tacttttacc 2280
tgtgtgttca aaggcatttc ttttcagcag tgatcattat aacttcacaa aaaaagatgc 2340
tgacggattt acttacaggg ctttaattgt attttgtccc agccaacacc ctctagggtc 2400
taaaagtcaa ggtacttcag tttatttggc aaacatgaca acattttttt tggccctggg 2460
cccaacagtt gtactttcat gaaacatatt gtacatttta catagttaa tttaaaaaat 2520
accttttaag ctagtgtatc tttgactgtc ttattttatc taacctttca gcacattcca 2580
aggttttagt tactcaggaa ggagttaatt aaaatgattt tattttgggt tgatggatgt 2640
tttttaaaag gaaaattatt attatgaacc ttcagcctac tttcttgagt gccgtaaaag 2700
tgcttgtaaa tctttttttt tttttaagaa gaaagaaaaa aatggtgttt gacgttgatg 2760
gaaattcaaa aatatatatg gaactgaaac attaacttag ctaaaataaa agcaatctgt 2820
gtttgaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 2861

```

&lt;210&gt; 44

&lt;211&gt; 84

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 44

Met Lys Phe His Ile Lys Asn Asp Asp Lys Phe Thr His Cys Ala Ile

1

5

10

15

Ser Gln Ser Glu Asn Lys Val Glu Leu Ala Val Phe Ser Leu Leu Leu  
 20 25 30

Ser Glu His Trp Asp Asn Trp Ser Phe Lys Asn Ile His Pro Leu Thr  
 35 40 45

Ala Ser Leu Ser Gly Tyr Phe Tyr Leu Cys Val Gln Arg His Phe Phe  
 50 55 60

Ser Ala Val Ile Ile Ile Thr Ser Gln Lys Lys Met Leu Thr Asp Leu  
 65 70 75 80

Leu Thr Gly Pro

<210> 45  
 <211> 1556  
 <212> DNA  
 <213> Homo sapiens

<400> 45  
 acttgtcctg cctgctgctg gggccctgg gctctatgtg catcctcttc actatctact 60  
 ggatgcagta ctggcgtggt ggctttgcct ggaatggcag catctacatg ttcaactggc 120  
 acccagtgtc tatggttgct ggcattggtg tattctatgg aggtgcgtca ctggtgtacc 180  
 gcctgccccg gtcgtgggtg gggcccaaac tgcctggaa actcctccat gcagcgctgc 240  
 acctgatggc ctctgccttc actgtttgtg ggctgggtgc tgcctttacg ttccacaacc 300  
 atggaaggac tgccaacctc tactcccttc acagctggct gggcatcacc actgtcttcc 360  
 tcttcgcctg ccagtgggtc ctgggctttg ctgtcttcct cctgccctgg gcgtccatgt 420  
 tcttcgcctg ccagtgggtc ctgggctttg ctgtcttcct cctgccctgg gcgtccatgt 480  
 ggctgcgcag cctcctaaaa cctatccacg tcttttttgg agccgccatc ctctctctgt 540  
 ccacgcgcatc cgtcatttgc ggcattaatg agaagctttt ctccagtttg aaaaacacca 600  
 ccaggccata ccacagcctg cccagtggag cggctcttgc caacagcacc gggatgctgg 660  
 tgggtggcctt tgggctgctg gtgctctaca tccttctggc ttcatcttgg aagcgcccag 720  
 agccggggat cctgaccgac agacagcccc tgctgcatga tggggagtga agcagcagga 780  
 aggggctccc aagagctcct ggtggtgcag cctgtgctcc cctcagaagc tctgctcttc 840  
 ccagggtccc cggctgggtt cagcaggcga ctttcttcca atgctgggccc cagacttctt 900  
 gcctgggtgc tggcctgccc tctccggcgg cttgctgcct gtctgcttcc cttgggtggc 960  
 ttgctgggtg gctgggctgc cctctccggc ccgcttgctg cctgtctgct ttcttggtg 1020  
 gctttgcctg ggtgctgggc ctgcctcttc tggctgctg ctgctgtct actgccacct 1080  
 tcagaggggt tccttgctga gacccacatt gcttcacctg gccccaccat ggtgcttgc 1140  
 ctggcccaac ctacgcttct gtgccatgct agagcttgag ctgttgctct tcttcagggg 1200  
 aggaatatag gtggagagcg ggaagggtct tgctcctaag tgttgctgct gtggcttttt 1260  
 tgccttctcc aaagacgcac tgccagggtc caagcttcag actgctgtgc ttagtaagca 1320  
 agtgagaagc ctgggggttg gagccacct actctctggc agcatcagca tcctactcct 1380  
 ggcaacatca ggccaacgct caccacagcc tcacattgcc agatgttggc agaagggcta 1440  
 atattgaccg tcttgactgg ctggagcctt caaagccact gggatgtcct ccaggcacct 1500  
 gggteccatg accagctccc cgtctccata ggggtaggca ttctactggt ttatgaagct 1556  
 cgagtttcat taaatatgtt aagaatcaaa aaaaaaaaaa aaaaaaaaaa aaaaaa

<210> 46  
 <211> 224  
 <212> PRT  
 <213> Homo sapiens

<400> 46  
 Met Cys Ile Leu Phe Thr Ile Tyr Trp Met Gln Tyr Trp Arg Gly Gly  
 1 5 10 15

Phe Ala Trp Asn Gly Ser Ile Tyr Met Phe Asn Trp His Pro Val Leu

20 25 30  
 Met Val Ala Gly Met Val Val Phe Tyr Gly Gly Ala Ser Leu Val Tyr  
 35 40 45  
 Arg Leu Pro Gln Ser Trp Val Gly Pro Lys Leu Pro Trp Lys Leu Leu  
 50 55 60  
 His Ala Ala Leu His Leu Met Ala Phe Val Leu Thr Val Val Gly Leu  
 65 70 75 80  
 Val Ala Val Phe Thr Phe His Asn His Gly Arg Thr Ala Asn Leu Tyr  
 85 90 95  
 Ser Leu His Ser Trp Leu Gly Ile Thr Thr Val Phe Leu Phe Ala Cys  
 100 105 110  
 Gln Trp Phe Leu Gly Phe Ala Val Phe Leu Leu Pro Trp Ala Ser Met  
 115 120 125  
 Trp Leu Arg Ser Leu Leu Lys Pro Ile His Val Phe Phe Gly Ala Ala  
 130 135 140  
 Ile Leu Ser Leu Ser Ile Ala Ser Val Ile Ser Gly Ile Asn Glu Lys  
 145 150 155 160  
 Leu Phe Phe Ser Leu Lys Asn Thr Thr Arg Pro Tyr His Ser Leu Pro  
 165 170 175  
 Ser Glu Ala Val Phe Ala Asn Ser Thr Gly Met Leu Val Val Ala Phe  
 180 185 190  
 Gly Leu Leu Val Leu Tyr Ile Leu Leu Ala Ser Ser Trp Lys Arg Pro  
 195 200 205  
 Glu Pro Gly Ile Leu Thr Asp Arg Gln Pro Leu Leu His Asp Gly Glu  
 210 215 220

&lt;210&gt; 47

&lt;211&gt; 2446

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 47

atgtttctgt ccttgctgtt gactcctggt tcgcccagga gatcctgggt caccagggga 60  
 tcaccaggg tggatgacag agggaaatatt tactctgggt tcactgctgg agctccagt 120  
 ccctgggagg ggagaggaga agcctgacct tgagcacagc tccagcctcc actccccact 180  
 cctatcgag gctctgggct gtggtttcat ttccccagt tctctaacta cttaggaagc 240  
 ccagagtttc tccctaaaga aagggtggcc tgcaactctt cctctcctcc aaaattaggc 300  
 ttcagtaatt gtcctctaag ttgatattta gaacttaaaa tctagacttc ttggatcacc 360  
 aagaaatatg caggaagtcc cacgtgatct catgtgtgat agcacgtaac ttagtgctac 420  
 aggtgacagt ttatttttg gaggagtcct attttacata gggaaataga aaactaaatg 480  
 gccctccttg aggcagccca agggctccac tggaccaga actctgagtt ctgatgctgc 540  
 ctatcatttg ctttttctca tgagctcctg gaccacctca tatcaccagc accctctacc 600  
 caaggtcctg gagccaacac tggagtaaaa gagcacaaga gaactcctta gaggtggaag 660  
 aagaacacag gtgttttgt cagacagcca gggccaatc ctgcttctgc ttagtgcaa 720  
 ttggctggaa gttgcagagg ttccagaagc tcatccaaga ttctggagcc cccgtctcca 780  
 gccctaattg tgtccgcctt ttcaccagc caccacctgt ttcaggagaa cacacagaag 840  
 tgatgacctt ttctggacag gataccctct gaggtcaaat cattgggctt gtggccatga 900

```

gaccacagac tttgcctggc ttccagtgac tgtccgagat gacctgtgga gtgcaccgcc 960
tgtacttgga catcactgtc ttttccgac catacccgtc atccttggaa tttccctgt 1020
atccaaacct gttcctgata cagataaccc taaggaatat cttggatgtc ttcacggggc 1080
ttatgactta ttctgaggtt tacaaaatag agggcccccag ctgagcaaag agagggctgc 1140
agcgccgagg gaacagctgc tggactcacc ggagaggcct cctggagcac gccgggaccc 1200
cctccacccc ctggcctttg ctgcatccat ggctagtttg cctgaactat atctgagcca 1260
ctgagagtat ttattaagca gagaataatt ttgagtttca cttttataac ttttgtttca 1320
aaagtgtgtt tggataaatt aatgaaatga ctgtaaacca gaattccctt tttcattatt 1380
tccttgtgtc gatataaaaa gagttgatga ttttaaaagc cagtggatg tccctgtctg 1440
tggatgtcag cgggagcgtg gggctctgaag cctttctctg gtggctgtcc aggaagattg 1500
ccccagcttg cactcccttt tggcggttc cagagccttc catttgaaag gagcatctct 1560
ccccactctg cccacccag caccagctct tctggaaccc actgacttcc taaacacttt 1620
cttcctcctt cacctcattg aattcaagac tttagaaaac agccaaaaac tctagaggtg 1680
ggggtgggtg tgacttgaa atccctgtg taatctttt aagggaatcc actggaaatt 1740
attaagctat gttttaaata aaaccattcc ttacttcaca gaattattaag ccactcactt 1800
tttaacagaa ctttgagata taattaaaat gccgtaaatt tcacccccac acagtgtaca 1860
gttccaaggt ttccaatata gtcacagaag tctctgcaac aattaccaca attttagaca 1920
tgaagaccat ttgttgagc ccctacctgt atctatgtgt atatatataa ttttaattt 1980
ggattagatg ttatagtatt taatattgta ctgttaccta tttgtgtctt tatattgctg 2040
ttttttattc aggtgaataa taattatctc catatagtca tacatttata tagggttaaa 2100
acataccttt ttacaattca ataaattaaa ttggccaggc gtgggtggctc atgcctgtaa 2160
tcccagcat ttgggaggcc gagggggggg tggatcacct gaggttgggg gttcgagacc 2220
agcctgacca acatggagaa accctgtctg tactaaaaat acaaaaaatt agccgggtgt 2280
ggtggctcat gcctgtaatc ccagctactt gggaggctga ggcaggagaa tcactagaac 2340
ccaggaggcg gacgttgagg taagccggga tcgcaccatt gcactccagc ctgggcaaca 2400
agacaaaaac tccatctcaa gaaaaaaaaa aaaaaaaaaa aaaaaa 2446

```

&lt;210&gt; 48

&lt;211&gt; 74

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 48

```

Met Thr Glu Gly Ile Phe Thr Leu Gly Ser Leu Leu Glu Leu Gln Cys
  1             5             10            15

```

```

Pro Gly Arg Gly Glu Glu Lys Pro Asp Leu Glu His Ser Ser Ser Leu
      20             25            30

```

```

His Ser Pro Leu Leu Ser Gln Ala Leu Gly Cys Gly Phe Ile Phe Pro -
      35             40            45

```

```

Ser Ser Leu Thr Thr Gln Glu Ala Gln Ser Phe Ser Leu Lys Lys Gly
      50             55            60

```

```

Gly Pro Ala Leu Phe Pro Leu Leu Gln Asn
      65             70

```

&lt;210&gt; 49

&lt;211&gt; 1231

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

```

ttggcgacac cccgtaccac ctgtatccgg cctatggctg ggtgatgttc gtcgctgtct 60
tcctctggct ggtgacaatc gtccctcttca acctctacct gtttcagctg cacatgaagt 120
tgtacatggt tccctggcca ctgggtgtaa tgatctttaa catcagcgcc accgttctct 180
acatcaccgc cttcatcgcc tgctctgcgg cagttgacct gacatccctg aggggcaccc 240
ggccttataa ccagcgcgcg gctgcctcgt tctttgcgtg tttggtgatg atcgctatg 300

```



```

gagtgaagtgc cttcttcagc taccaggcct ggcgaggagt aggcagcaat gcggccacca 360
gtcagatggc tggcggtat gcctaaacca cctgtgccac ggccccctct ggggctgaag 420
ccgccgctgg gtcacagagc aggggtcacc tgcaagcctg aagctgggga gccctgcgtg 480
gagtcagccc aacagggact gcatttgctc ctctctgccc gtcagacata agctctcaca 540
gcgctaagga agcaggccca ggctggcagg catctcggct tgcaggaggc caactgctga 600
gacctcttct ccatccccct tattcagtgg aagatgacgg gggatctgag gctgtgtctc 660
tgccttgtct ttagaggact tcagcgtcca agactggggc ccacccttct caccagcact 720
aaatgcacta acaggactcc agacctgcag ccccagaccc gccgtagtat aagcctaaca 780
agcaacacgt agcaccttag tctttgttcc aggagagctg agcaagctgg tgaaaccact 840
ctccttcctt taaacaccgt ttcaaccaac ctctccctgg agccaacctg taaaaagtgg 900
gttgattgct gacagcatgg tcttccctcc ctgcatttca gacataccag ttactgaaaag 960
caaatcagtt ttaagtgatt tctcagtgtc gaaaagcctg tccaggtttc cttecccttc 1020
ccaagcctct ctctgtaata ctccctttgg gcgaagctaa catcggtgcc tccccgacct 1080
tgctgagctg gacatggga cgcaaaggag ggaggggaagc aaggccttgc ctggcgagtt 1140
gtcatgtggt tgggtgtgac tggtttattt tttttaataa aaataaagat gagagaaatt 1200
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa a                                     1231

```

&lt;210&gt; 50

&lt;211&gt; 113

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 50

```

Met Phe Val Ala Val Phe Leu Trp Leu Val Thr Ile Val Leu Phe Asn
  1              5              10              15

```

```

Leu Tyr Leu Phe Gln Leu His Met Lys Leu Tyr Met Val Pro Trp Pro
      20              25              30

```

```

Leu Val Leu Met Ile Phe Asn Ile Ser Ala Thr Val Leu Tyr Ile Thr
      35              40              45

```

```

Ala Phe Ile Ala Cys Ser Ala Ala Val Asp Leu Thr Ser Leu Arg Gly
      50              55              60

```

```

Thr Arg Pro Tyr Asn Gln Arg Ala Ala Ala Ser Phe Phe Ala Cys Leu
      65              70              75              80

```

```

Val Met Ile Ala Tyr Gly Val Ser Ala Phe Phe Ser Tyr Gln Ala Trp
      85              90              95

```

```

Arg Gly Val Gly Ser Asn Ala Ala Thr Ser Gln Met Ala Gly Gly Tyr
      100             105             110

```

Ala

&lt;210&gt; 51

&lt;211&gt; 3290

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 51

```

tctctgtcaa ctgcttggac cagcatgttc ggaagtcccc cgagagcgtt gctttgatct 60
gggagcgcga tgagcctgga acggaagtga ggatcaccta cagggaacta ctggagacca 120
cgtgccgcct ggccaacacg ctgaagaggc atggagtcca ccgtggggac cgtgttgcca 180
tctacatgcc cgtgtcccca ttggctgtgg cagcaatgct ggctgtgcc aggatcggag 240
ctgtccacac agtcattctt gctggcttca gtgcagagtc cttggctggg aggatcaatg 300
atgccaagtg caaggtggtt atcaccttca accaaggact ccgggggtggg cgcgtggtgg 360

```

```

agctgaagaa aatagtggat gaggtctgtga agcactgccc caccgtgcag catgtcctgg 420
tggtctcacag gacagacaac aaggtccaca tgggggatct ggacgtcccg ctggagcagg 480
aaatggccaa ggaggaccct gtttgcgccc cagagagcat gggcagttag gacatgctct 540
tcatgtctgta cacctcaggg agcaccggaa tgccaagggt catcgtccat acccaggcag 600
gctacctgct ctatgccgcc ctgactcaca agcttgtgtt tgaccaccag ccaggtgaca 660
tctttggctg tgtggccgac atcggttggg ttacaggaca cagctacgtg gtgtatgggc 720
ctctctgcaa tggtgccacc agcgtccttt ttgagagcac cccagtattat cccaatgctg 780
gtcgggtactg ggagacagta gagagggtta agatcaatca gttctatggc gcccacacgg 840
ctgtccggct gttgctgaaa tacggtgatg cctgggtgaa gaagtatgat cgctcctccc 900
tgccggaccct ggggtcagtg ggagagccca tcaactgtga ggcctgggag tggcttcaca 960
gggtgtgtggg ggacagcagg tgacagctgg tggacacctg gtggcagaca gaaacagggtg 1020
gcatctgcat cgcaccacgg ccctcggaag aaggggaggga aatcctccct gccatggcga 1080
tgaggccctt ctttggcatc gtccccgtcc tcatggatga gaagggcagc gtcgtggagg 1140
gcagcaacgt acgtggggcc ctgtgcattc ccaggccctg gccggggcatg gccaggacca 1200
tctatggcga ccaccagcg tttgtggacg cctacttcaa ggcctacca ggctattact 1260
tcaactggaga cggggcttac cgaactgagg gcggtatta ccagatcaca gggcggtatg 1320
atgatgtcat caacatcagt gccaccggc tggggaccgc agagatttag gacgccatcg 1380
ccgaccaccc tgcagtacca gaaagtgtctg tcattggcta cccccacgac atcaaaggag 1440
aagctgcctt tgccttcatt gtggtgaaag atagtgcggg tgactcagat gtggtggtgc 1500
aggagctcaa gtccatgggt gccaccaaga tcgccaata tgctgtgctt gatgagatcc 1560
tggtgtgtgaa acgtcttcca aaaaccaggt ctgggaaggt catgcggcgg ctcctgagga 1620
agatcatcac tagtgaggcc caggagctgg gagacactac cacttggag gacccagca 1680
tcatcgaga gatcctgagt gtctaccaga agtgcaagga caagcaggct gctgctaagt 1740
gagctggcac cttgtggggc tcttgggatg ggcgggcacc caagccctgg cttgtccttc 1800
ccagaaggta cccctgagggt tggcgtcttc ctacgtcca gaagcagccc ccacccaca 1860
catgacccac accgcccctca cgtgaagctg ggctgagagc cctttctccc atccattgga 1920
ggtcccagga gtgtcaccca tggagaggct atgcgacatg gctagggtg gttctgcat 1980
ctgagtttgg tttcctggaa tgaaaaggca ttgccatctc cattcctctg cctcttgag 2040
ccagcacagg aaggtgaggc cctgggtag cgcgcctgct cagataaac agagctagt 2100
agctagtagc aaccgtgtt tctccagatc tgtctagata caaaggctcag aaatcttatt 2160
tttatacttt tatattgttg aagaacagca tgcaaacctc acatgtagt tgtggattta 2220
cttgaacatg ttctttttaa catgtagtta tgaaaatctc cttttttgcc tctactggtg 2280
aggaacatg aggatcagag gccacatttt taattattgt tagtgtattt ggaagtctga 2340
attggagatg tttgtacctc tgtctaaaca gtcccttga gaacttcaa gcctccgga 2400
tcttttctctg ttgagtggtt ctctgtgct tgggtgtgta taatggagct aactcctaag 2460
cgggtgggtg aatgtggccg ccttagttct gaagctactc cagttatgtt ctgtttcttc 2520
aagctgtgat ccagaaagat ttttgtgccc ccagatgcct cttgatagga gaggcaacat 2580
actccaaata gttgggttct tcagggaagc tattagaaac tcaggtagt tgttagagca 2640
ctaacttggc cagagccaaa tcctggcaaa cgtgcctga ccttactct gtggttgggg 2700
cagtgaagac cactgaggtc caatgatgag acttggaggt ctggatccag tctctctttg 2760
ttttaatgtg acttaggtgc tgtcaacatt agcaagataa tggaaatcac gacgccagt 2820
ggtgcttacc tccctgctag gcatgcaggg gctggcggtt ggcaggggaa ggaggccag 2880
tgagccgggt cccttagggg agggagagtt tgtcctcttt gccccacagt ctacccttca 2940
gggccttgtg gcagtccag tgttcggggg gtgtctgggc cactgagtac ccactcggtc 3000
gtggttgtgc tggcctcttg ggtgagtga cctgtgaagc ccaggaggtg gtgttggtg 3060
cagggtacac aaatactgag tgggtgtctt ttgttacagg cttagcaaca aagctgtgcc 3120
ctgggcatgg ggggtctgtag tgtagctaca gttgtgcgtt tgtgaaatg cttagcttct 3180
catgttgctg agaggaacct ggacatggc cgggcatct gaatgatctg taggggagg 3240
agttcaataa aagctttatt ttgttccaaa aaaaaaaaaa aaaaaaaaaa 3290

```

&lt;210&gt; 52

&lt;211&gt; 518

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 52

Met Pro Val Ser Pro Leu Ala Val Ala Ala Met Leu Ala Cys Ala Arg

1

5

10

15

Ile Gly Ala Val His Thr Val Ile Phe Ala Gly Phe Ser Ala Glu Ser

20	25	30
Leu Ala Gly Arg Ile Asn Asp Ala Lys Cys Lys Val Val Ile Thr Phe 35 40 45		
Asn Gln Gly Leu Arg Gly Gly Arg Val Val Glu Leu Lys Lys Ile Val 50 55 60		
Asp Glu Ala Val Lys His Cys Pro Thr Val Gln His Val Leu Val Ala 65 70 75 80		
His Arg Thr Asp Asn Lys Val His Met Gly Asp Leu Asp Val Pro Leu 85 90 95		
Glu Gln Glu Met Ala Lys Glu Asp Pro Val Cys Ala Pro Glu Ser Met 100 105 110		
Gly Ser Glu Asp Met Leu Phe Met Leu Tyr Thr Ser Gly Ser Thr Gly 115 120 125		
Met Pro Lys Gly Ile Val His Thr Gln Ala Gly Tyr Leu Leu Tyr Ala 130 135 140		
Ala Leu Thr His Lys Leu Val Phe Asp His Gln Pro Gly Asp Ile Phe 145 150 155 160		
Gly Cys Val Ala Asp Ile Gly Trp Ile Thr Gly His Ser Tyr Val Val 165 170 175		
Tyr Gly Pro Leu Cys Asn Gly Ala Thr Ser Val Leu Phe Glu Ser Thr 180 185 190		
Pro Val Tyr Pro Asn Ala Gly Arg Tyr Trp Glu Thr Val Glu Arg Leu 195 200 205		
Lys Ile Asn Gln Phe Tyr Gly Ala Pro Thr Ala Val Arg Leu Leu Leu 210 215 220		
Lys Tyr Gly Asp Ala Trp Val Lys Lys Tyr Asp Arg Ser Ser Leu Arg 225 230 235 240		
Thr Leu Gly Ser Val Gly Glu Pro Ile Asn Cys Glu Ala Trp Glu Trp 245 250 255		
Leu His Arg Val Val Gly Asp Ser Arg Cys Thr Leu Val Asp Thr Trp 260 265 270		
Trp Gln Thr Glu Thr Gly Gly Ile Cys Ile Ala Pro Arg Pro Ser Glu 275 280 285		
Glu Gly Ala Glu Ile Leu Pro Ala Met Ala Met Arg Pro Phe Phe Gly 290 295 300		
Ile Val Pro Val Leu Met Asp Glu Lys Gly Ser Val Val Glu Gly Ser 305 310 315 320		
Asn Val Ser Gly Ala Leu Cys Ile Ser Gln Ala Trp Pro Gly Met Ala 325 330 335		
Arg Thr Ile Tyr Gly Asp His Gln Arg Phe Val Asp Ala Tyr Phe Lys		

340 345 350  
 Ala Tyr Pro Gly Tyr Tyr Phe Thr Gly Asp Gly Ala Tyr Arg Thr Glu  
 355 360 365  
 Gly Gly Tyr Tyr Gln Ile Thr Gly Arg Met Asp Asp Val Ile Asn Ile  
 370 375 380  
 Ser Gly His Arg Leu Gly Thr Ala Glu Ile Glu Asp Ala Ile Ala Asp  
 385 390 395 400  
 His Pro Ala Val Pro Glu Ser Ala Val Ile Gly Tyr Pro His Asp Ile  
 405 410 415  
 Lys Gly Glu Ala Ala Phe Ala Phe Ile Val Val Lys Asp Ser Ala Gly  
 420 425 430  
 Asp Ser Asp Val Val Val Gln Glu Leu Lys Ser Met Val Ala Thr Lys  
 435 440 445  
 Ile Ala Lys Tyr Ala Val Pro Asp Glu Ile Leu Val Val Lys Arg Leu  
 450 455 460  
 Pro Lys Thr Arg Ser Gly Lys Val Met Arg Arg Leu Leu Arg Lys Ile  
 465 470 475 480  
 Ile Thr Ser Glu Ala Gln Glu Leu Gly Asp Thr Thr Thr Leu Glu Asp  
 485 490 495  
 Pro Ser Ile Ile Ala Glu Ile Leu Ser Val Tyr Gln Lys Cys Lys Asp  
 500 505 510  
 Lys Gln Ala Ala Ala Lys  
 515

<210> 53  
 <211> 1467  
 <212> DNA  
 <213> Homo sapiens

<400> 53  
 tcgcctgaac aaggaccatg ctgtcctgca cgctgggtct gaccgtctgc cctctctccc 60  
 cagcaccaag cgtgaccttg gctgtggcgc tcaacggcca gctccggcgg cccctctgct 120  
 gctcctcggc ttcccccga gtgggagagc ctgcctggcc tcggcctttg tccagcgacc 180  
 aggctctgtc cccgagaagc tacggccgac ctgggtctgg tgttgggacg catggaccgg 240  
 gctggggagg tgcacagagt gatgttaact ttttcccgtg ttagatatg tacagccaaa 300  
 gggctcgtgta aatgttctgc aaaagtgggt ctatacagag tgaaagctat ttattttgtg 360  
 cagagaaaaa agtctggagg gatggaacct tcagggttta ttcattatta agatgtagct 420  
 tttgttgtt tcaggcatta tgtataaagc aacgattatt ttatggacca agttttcatg 480  
 taactgttg agtgaaagt caatatctga ccccccgtct cccagcagga agttgcttg 540  
 cccgacaatc acagccccg tcaggggccc tgtggccagt gcctcctcct ctcttgccc 600  
 caccttatec tgtcttgcc gctgcctggg agaccagcca tccagagaag cacctggaag 660  
 agtctcgggc cctcctgcaa taaaggccgg gagccctgt gggcagtggg ctccagcctct 720  
 cccaggggg gcagctcccc cacggctcct cactccccgc ctgctgccc agccgcccagc 780  
 catgccaagg acaacagcaa tagtcccctg gggctctccc agcggccctc agccatagat 840  
 ggcaagggtg gcagcctgcc ccccatggg aagtctcttc tgtatccagg tctgctttcc 900  
 acctccctc agattcctt tggcacattc tcctcttgag gaagtaccag tctttctgaa 960  
 actaagagag ggaggggcag gtcctttaa aataccaaaa atgtttacag agttgggtgc 1020  
 tgagctgcag ggctcaggcc tgaccagtca taaccaaagg gtgaggcagg ccttgctgac 1080

```

tgccaccccc caggcctgtt agaatagaag ccttagtccc actcccacca cacccccacg 1140
ccccaccacc tgcctttctt ttgatttcta aagagggatt cagcagagac cccccaaccc 1200
tccctggctc ggtctgagtc ccaactgccc ccccatcaca gccttcacgt ctcagccctt 1260
cccgctctgt ctgtccgtgt gccgtctgtt tctctgggcc atgtgtgagc agtgtcccca 1320
tctcccatc cgtccctgct gtccccgc attgggcct gagtgtgctc tgtatacaac 1380
gtcatgtctg ttacaccaat taaagaagcg ggaaggcttc aaaaaaaaaa aaaaaaaaaa 1440
aaaaaaaaaa aaaaaaaaaa aaaaaaa

```

1467

&lt;210&gt; 54

&lt;211&gt; 132

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 54

```

Met Tyr Lys Ala Thr Ile Ile Leu Trp Thr Lys Phe Ser Cys Asn Cys
  1             5             10             15

Cys Ser Glu Ser Ala Ile Ser Asp Pro Pro Ala Pro Ser Arg Lys Leu
      20             25             30

Leu Gly Pro Thr Ile Thr Ala Pro Val Arg Gly Pro Val Ala Ser Ala
      35             40             45

Ser Ser Ser Leu Gly Pro Thr Leu Ser Cys Leu Ala Cys Cys Leu Gly
      50             55             60

Asp Gln Pro Ser Arg Glu Ala Pro Gly Arg Val Ser Gly Pro Pro Ala
      65             70             75             80

Ile Lys Ala Gly Arg Pro Cys Gly Gln Trp Ala Gln Pro Leu Pro Arg
      85             90             95

Gly Ala Ala Pro Pro Arg Leu Leu Thr Pro Arg Leu Pro Ala Gln Pro
      100            105            110

Pro Ala Met Pro Arg Thr Thr Ala Ile Val Pro Trp Gly Ser Pro Ser
      115            120            125

Gly Pro Gln Pro
      130

```

&lt;210&gt; 55

&lt;211&gt; 943

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 55

```

tatacttgct taagcaatct tgatttgagt aagggtcttg atttgtgcta ttatgttctg 60
ttagtttttg catgaatata ctaaagcttt tttttttttt tctagcatgt gtttttctct 120
ctttggttct ctttgtattt actacttttc ttttttttct gtgttttttt ttctctgttt 180
ttgtttttgt tgggtgtttt ttcctgtctt cattgtttca ggtattttct taccctctgt 240
gattccccac gggctggatc gagatgggcc agttatgccc agctccttcc tctcctctct 300
cctcctctgg tagagcactc ttgcgatgct gacactgcca acctccagta toctcaccct 360
cgcagacgat atctctctcg gcctcttaat cccttacctg agaatgaagg gattttaaac 420
actgatttaa cattgaaagg ccttattcaa gtgcttgtaa atgctttcat ttctggctgc 480
tttttgtttt tcattttctt tcagaagatt tttctaactt agggctctgtc ttgcattgat 540
tacaaccaga atacagtgtt tggaaacctaa atctgtttgt gcgtctgcat caaaggaaca 600
tttgcttcac tgggtgataa ctttgatga aatgagatat gtccaagtaa cgtaactgt 660
gaagttacac acagtagctg acttcaaagt gcctgttctg taaattttat tttaaactgt 720

```

taccatagtc ttaagttggt tatgctttat cagactggct aatgtgaaag cataatatta 780  
 tgaagtttat tctgccttat gagaccttaa aaaatggatt tcattttaca ggctaattgtt 840  
 gtaactgact agtatgtaaa ataaatcatt cctgtgtata aagcagcaaa acctaaaaaa 900  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 943

<210> 56

<211> 86

<212> PRT

<213> Homo sapiens

<400> 56

Met Asn Ile Leu Lys Leu Phe Phe Phe Phe Leu Ala Cys Val Phe Ser  
 1 5 10 15

Ser Leu Val Leu Phe Val Phe Thr Thr Phe Leu Phe Phe Leu Cys Phe  
 20 25 30

Phe Phe Pro Val Phe Val Leu Phe Gly Val Leu Phe Leu Ser Ser Leu  
 35 40 45

Phe Gln Val Phe Leu Tyr Pro Ser Gly Phe Pro Thr Gly Trp Ile Glu  
 50 55 60

Met Val Gln Leu Cys Pro Ala Pro Ser Ser Ser Ser Ser Ser Ser Gly  
 65 70 75 80

Arg Ala Leu Leu Arg Cys  
 85

<210> 57

<211> 1032

<212> DNA

<213> Homo sapiens

<400> 57

aaaacgaatt gattaatgaa gagttgggaa tagcttatcc aatcattgat gggatcccta 60  
 atatgatacc acaggcagct aggatgacac gtcaaagtaa gaagcaagaa gaagtggagc 120  
 agcgctagtt cataatttaa aaaaattaaa aaaacgcaac agccaacttt tcttaatacc 180  
 atataccttt taaaacacag tggcaggtaa taagtggaag agaagaatgt ttctgtctct 240  
 tctacgttg actgttctta ttccactggg ttcttttagca ggactgttct actcagcctc 300  
 tgtggaagaa aacttccac agggctgcac tagcacagcc agcctttgct ttacagcct 360  
 gctcttgcc attaccatac cagtgtatgt attcttccac ctttggaactt ggatgggtat 420  
 taaactcttc aggcataatt gatgcaacta gagtcaatat gctgtatata ttaatgatag 480  
 ctcttgggca tcgatctctg aaagctcaaa tggatggaat ttagtttgcg ggaaagaggc 540  
 tttgctttgc gcatatcagg cttaggactg tgggaggctt aagttgcaga tgcttctttt 600  
 attgtactct tgttctgccc ttgttttttg aaggctctga cttataactg ctgtatcaga 660  
 agaaacattt tgacagtgtc ttggttgag atgaacatcc ctaattgaca tgtgatgact 720  
 atttcttatt ccattcatct aagagtcatt gaaattttgt tttgcttggt tgtttagctt 780  
 caaggctctt ggtaaagtca catgttaagg atgactgaaa taattccaaa ggagtgatgt 840  
 tggaatagtc cctctaagg agagaaatgc atttgaacga atgtgatata aaaccacata 900  
 atcaaataga aacttcatgt acttacaata actgagtttg taaaattacc ttcatttctt 960  
 tgacattaaa tgcttatatt agcaataaac atgttgacac tttcctataa aaaaaaaaaa 1020  
 aaaaaaaaaa aa 1032

<210> 58

<211> 71

<212> PRT

<213> Homo sapiens

&lt;400&gt; 58

Met Phe Leu Ser Leu Pro Thr Leu Thr Val Leu Ile Pro Leu Val Ser  
 1 5 10 15

Leu Ala Gly Leu Phe Tyr Ser Ala Ser Val Glu Glu Asn Phe Pro Gln  
 20 25 30

Gly Cys Thr Ser Thr Ala Ser Leu Cys Phe Tyr Ser Leu Leu Leu Pro  
 35 40 45

Ile Thr Ile Pro Val Tyr Val Phe Phe His Leu Trp Thr Trp Met Gly  
 50 55 60

Ile Lys Leu Phe Arg His Asn  
 65 70

&lt;210&gt; 59

&lt;211&gt; 1564

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 59

gtagcatttg tctccttctt tcaatttggt aacacatcag atagtagatt tttcgtagta 60  
 agaatgaaat gtttttactt ctcctttaat attttatggt gaatgtcaga gcctccatat 120  
 tggaaattaa aaggcacctt ttactgcatt ctctctgtgc tttgtctagg tcccttctag 180  
 agccttcagg aatcggtttg tgggaattgca ctttgatgag ttacctagct ccgagttgga 240  
 aacaatcttg cacaagcggg gtagtttgcc accctcctat tgcagcaagt tggtaaagt 300  
 catgctggat cttcagggtac ttagtttcaa atggaatgta ttgattatac ttgggtgatca 360  
 aaatttcatg gcttcttttg ttagaaaaa gaatctgtct gttgatgtat ttttccaagt 420  
 agagattttt ttttattttt tattgttttt aattttttta ttattttatt tttgcaagac 480  
 acagtctcac tctactctcc tggctggagt gtagtggcac aatcatagct cattgcagcc 540  
 ttgacctcct gggctcagg gacctcccca tctctgacta ctactcctt agtagctggg 600  
 accacagatg tgcaccgtca tgcccggcta catttttttt tttgtagaga caaggtttcg 660  
 ccatgtttgc cggctgggtc tcgaactcct gggtcaagg gatccacca ccttgatctc 720  
 ccaaagtttt gggattatag gtgtgaacca cagttcttgg ccttcaagta gaggtctttt 780  
 aaatgcttcc caataatata tatttttagt tttttttttt aatcaatcaa caattttaca 840  
 tcattaaaat ttttctgata ttacatatg aggtttttgt atttgtgtat aatattgaag 900  
 cattttgttt agaatgtaat ttggcttagg gaaaactggt ttttggtttt taaatggata 960  
 tagtccctga atggacataa ttaaagagat ccattctttg tggctccccc tagtcaactg 1020  
 caaagctaca gctgatcttt aaaaattcca ttatcttat ggggttaaaa gttttcttta 1080  
 tcagaaaagta tacattgtca aacttctttt tagaaaagcta ataataatga agactcctac 1140  
 cattatgggt ccacattggt tcaactaact tctgcatct ttgccaccat ttgcattact 1200  
 ttttgtttgt tttttgagac agagcctcac tctgtcgcca ggctggagt cagtggcgct 1260  
 atctcagctc actgcaacct ccgcctcccg ggttcaagca attcccttgc ctcagcctcc 1320  
 tgagtagctg cgactatagg cgtgcaccac caggcttggc taattttttg tctctataaa 1380  
 aaatttttaa agttggccag ggggtgatggc tctgectgta gtcccagcta ctcgaaaggc 1440  
 ccaggtggga ggactgcttg agcctaggtg ttccaggctg caggagagctg tgatagctcc 1500  
 actgcactcc accctgtgta ataaaggag actgtatctc aaaaaaaaaa aaaaaaaaaa 1564  
 aaaa

&lt;210&gt; 60

&lt;211&gt; 82

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 60

Met Val Asn Val Arg Ala Ser Ile Leu Glu Ile Lys Arg His Leu Leu  
 1 5 10 15

Leu His Ser Ser Cys Ala Leu Ser Arg Ser Phe Leu Glu Pro Ser Gly  
 20 25 30  
 Ile Gly Leu Trp Asn Cys Thr Leu Met Ser Tyr Leu Ala Pro Ser Trp  
 35 40 45  
 Lys Gln Ser Cys Thr Ser Gly Val Val Cys His Pro Pro Ile Ala Ala  
 50 55 60  
 Ser Trp Leu Lys Ser Cys Trp Ile Phe Arg Tyr Leu Val Ser Asn Gly  
 65 70 75 80

Met Tyr.

<210> 61

<211> 2800

<212> DNA

<213> Homo sapiens

<400> 61

taatagatta ttatcaacta gagatattta taaatgaagt atatactttt aaaacatgta 60  
 agttataacta attttgtgtt taatttcata tgttttcatg taatgaaaat tctatttttt 120  
 ttagaattgg ttttgttttt attctattat cctttatttc cagttgtcaa actttaaatg 180  
 gctatgtctg cattctaata acactgttca gtctgttatg gaaaaggaga actagggaac 240  
 aaatgttgct cagggctggt gtcagtgaga aaaatttatc aatgcttttt aatgtgtttt 300  
 tacccttgcc tactctgtg tgtgtcactt tctataatat aaagaaatac tataatattt 360  
 ctagaatctg gaactgtcac catgatgaat ggccctttca atgcatagtt acagaaattc 420  
 ctgaagattc ccaggactt caatttcatt ggtttttatt tgcagttttt agttgtctgt 480  
 attgttgctg tttccagtct aaaggacctc ctttggtaaa ggtgaacaag acctctcccc 540  
 tatgctaccc agccagattt tgtgtgtgta atgggtggc ccaagagtgt tcttttactt 600  
 agaactcctt ttgattttgc tttctctgc ccctagaaat tttagggaca aagacatttt 660  
 ggaaattgtc agttactttt agaaagaaaa cactgcagaa atatttaaca gattacttct 720  
 tgataaaatt taaatgggat tatatgaatt tgtaatgcc aattgaaaata ttatatacac 780  
 aaaatttgct attttcactt aaagtaagca tttttaaagg atttatttta gaatacaatc 840  
 tatgcaatcc tcgaactcag tgtgcttttc cctgacagga tatataaaaa ggtagattag 900  
 gtcaaccctc ctttaagctt cctctcccc cactttttct tttcactgta tataaacggg 960  
 actttctgtg attcatacca acagtcctaa actgtcattt tctaaatctg ggatttaaaa 1020  
 tcttttaaca atatttaaaa tcttgatttt ttttaaagt cagcagatac aatttggaag 1080  
 attgcatctg ttagaagta atgcaggta aagatctaga gtttttctt agactcttta 1140  
 gtctttctg aattaaagaa taagtgtct taaatccata ttttatctg tcaaagtctg 1200  
 aaatgtgtc attgatgatt tggtagacat taaaaaattg tatttagttg gcttggctca 1260  
 cccctcttaa aaatacttag ccactctctg gtgggagtgt gtgggggtat taaggctctca 1320  
 gagttattta tttctcacc taatttcaaa cttttataaa cttgggggtt tgttgttgat 1380  
 ggtggtagta tttttgtta agttgttta aaggctgact tgattggagc ctctctact 1440  
 tttcttgag ggttgcaagt attgtatggt gggggagggg ggccttaagg caaactgtct 1500  
 aaaaagtgac tatccaatta atttaactag taaattagta agaattatc ttagtattca 1560  
 ttcagagttt aagatgtgaa aaatctgcac tttgtttatc actctaaatg taaagaactt 1620  
 tttgatagtg ttatttcacc actccctcag ttaaaggcta ttatttaaac tgtttactgg 1680  
 agaaaaacct cgctcatgtc catttattgt tttttctgt actgtgattt gtttcaagct 1740  
 taggaaaaat agtatattag agtatgttct aggaatttaa aagatctggt tagagtaaaa 1800  
 agttcttttt aaggttctta actaattttt tcacaactaa gaaaataaat gaagtattct 1860  
 taggtgaaa ttcactctat tttatcataa attagattgt aggggcagcc tacatttttg 1920  
 tgtatgtgtt tttatttctt aaatgattgt gtgagcctgg tgacatttta tggttcttgt 1980  
 gatctaaact gtttttccaa ttcacatctt ttgtgtgaa gtgatattat actagagtac 2040  
 tgtttgcatt gtaaaaatgc tttgtgtgtg ctctggcatt ttgtctttat ctcatcact 2100  
 aatttaaaaa ccagcactt gataatataa ctgacagaaa tgattgtacc cactgatgaa 2160  
 gtaatgaaaa tgaagaaaag gaaaatattc cttccttcaa tggcgtaagt ttatttttta 2220  
 cttaagtggc atctatgtaa gttagacaaa ctatatatta aaattggtaa actttgaatg 2280



```

tttctcctgt tttgattctt agattatgag gaggatcaga cagaataaga ctccacactt 2340
ttgaaatttg ataagtcaaa acgtttttatt ttgacatttt taacattgtg aattattcga 2400
tatttgtaat ctatgtatat gaaactctgc catgtttttc aactttgcct atgtgccact 2460
gtatataaga aaggacagct atttgatctt tcatcagcat tgccttcagc tgaaaatttg 2520
ggaacagttc acagtaaacc tctcagtagt ttgcaaactg aggtacctgc agaaaccata 2580
gctactcacc aaaatacaag attaggtgca tggatcgtg gaaattattt ggtaaaaaaa 2640
cacacacaca aatctatgtg tactgtattt taaggaatat ctacaggtgc agctttgatt 2700
ttgaacagaa ttccaaataa tttactgttg taccaccag aatctagaaa atgtgaatta 2760
aagtacatgt ttctgcaaaa aaaaaaaaaa aaaaaaaaaa 2800

```

&lt;210&gt; 62

&lt;211&gt; 170

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 62

```

Met Phe Ser Cys Asn Glu Asn Ser Ile Phe Phe Arg Ile Gly Phe Val
  1                      5                      10                      15

```

```

Phe Ile Leu Leu Ser Phe Ile Ser Ser Cys Gln Thr Leu Asn Gly Tyr
      20                      25                      30

```

```

Val Cys Ile Leu Ile Thr Leu Phe Ser Leu Leu Trp Lys Arg Arg Thr
      35                      40                      45

```

```

Arg Glu Gln Met Leu Leu Arg Ala Gly Val Ser Glu Lys Asn Leu Ser
      50                      55                      60

```

```

Met Leu Phe Asn Val Phe Leu Pro Leu Pro His Ser Val Cys Val Thr
      65                      70                      75                      80

```

```

Phe Tyr Asn Ile Lys Lys Tyr Tyr Asn Ile Ser Arg Ile Trp Asn Cys
      85                      90                      95

```

```

His His Asp Glu Trp Pro Phe Gln Cys Ile Val Thr Glu Ile Pro Glu
      100                      105                      110

```

```

Asp Ser Pro Gly Leu Gln Phe His Trp Phe Leu Phe Ala Val Phe Ser
      115                      120                      125

```

```

Cys Cys Asn Cys Cys Cys Phe Gln Ser Lys Gly Pro Pro Leu Val Lys
      130                      135                      140

```

```

Val Asn Lys Thr Ser Pro Leu Cys Tyr Pro Ala Arg Phe Cys Val Cys
      145                      150                      155                      160

```

```

Asn Gly Leu Ala Gln Glu Cys Ser Phe Thr
      165                      170

```

&lt;210&gt; 63

&lt;211&gt; 2056

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 63

```

acaccatgtc atgccctttg ccactgttga tticagccat agctgcagtg ggtagttcca 60
tgcaaaactca cgctcgagcc agttttgctg cagggccttc tcagaagact tctcagccca 120
tttgggcccag gataacaca gcccagaagt gcaggggcat taccacgcca ggggcaatcc 180
tccagcagtg ggagatacaa gcctatggat gaaggttccc acctcccatc actcagatga 240

```

```

gaaacaccaa gaggtcttct gtacctttct cagaggccac agcaggatca atccccatt 300
gcacacagca gccatctcaa taatgcacca ttctattagt gggatatatgc ataacaggg 360
attccttggt gcctctcttg gtttttcaag ctccagcaata gtggagtggg tgcacagcca 420
ggggctagca atgggaagcac acaaaagggc ttgaaactgct caccaagtta ctgggttact 480
tcacctggta aacaacctaa atcacatagg cgccactaag agtaggagaa atctaaaatg 540
ggtaatgggg aagggagatg aggaattaat tacagctggc ttaatcagaa ctgcctgcat 600
ccgctggtag ttttagcatg gacatgtact tcatgttgcc caggctggcc tcgaactcct 660
gggctcaagc gatcctctca cctcagcctc tggagttagt gggactatag gcacatgcc 720
ccatattcag caggacatat acttttataa gtgtgggac tgctgtcacc ttgaggatgt 780
actaactggg ttgacttcac gtagggcatg ggtagatctg agcagggcaa agtagactgc 840
agggatgtgt atggagtcac cacaccatgt catgcccttt gccactgttg atttcagcca 900
tagctgcagt gggtagttcc atgcaaaactc acgctcgagc cagttttgct gcagggcctt 960
ctcagaagac ttctcagccc atttgggtcca ggattttcct ccactttaa gtaactgctc 1020
ccaagtctg ccccatgttc tactttcagg agttcccaaa ctaaggcaat agctgcacct 1080
gcaaagaaag cgactacttg ttaatatgtg atataatcca gaaaagcaag agtccaccca 1140
aagtcagaag agaagaattc agaagagtaa ggtgggtgtg ggcataatgat acgtatgagg 1200
gggcatttga taacttcaac ctaccaagtg cagttcattc agtaagccag taatatagg 1260
ttttagagct gacaatgtga ggtataatat acaaaagcct agaaaggcat agggagacat 1320
ctatcttttt cctttgcttt gagggcatgt gcttgcata gctctgtgtg tgtgtgtgtg 1380
tgtgtgtgtg tgtgtatata cacacataca ctgactata aaaagaaatc acgacatcca 1440
tacggaggaa cctgggtatt tgaatcccag aaactttctc aaggcctcac tagggaaaga 1500
tagttcagtt acaagccaaa gactcaagag gaaaaagccc ctggagtga taaatagtct 1560
gataaaccta gacatatgtg gtaactgcag aaataagtcc tttttcattt atagaggagg 1620
agaagttgca tatattaatc atgtctgctc ttgccccac tttcaggaa cacttatatg 1680
ccttatgaaa catcaatgtt ccatgaaaca cagtttgaaa acctgagcta acctctcacc 1740
tttcttctag agcgtctcca cttggtcatc tctaacagac cggacccgc cactttgcta 1800
ggatatccca gaggagacca gagcagaatt ccctgtgaag atattgcaaa tgctgcacca 1860
ctttgacaca gacgaatatt gccttaatgc tttctcattg cagctccatt ctcatcctc 1920
ttgttcta at ctactggga acagtaagaa gctgccatca ttttatgtta tacttttaca 1980
tactttaagg ttattactga ttaatgagaa ttaaaggaag ctgagaatat ataaaaaaaa 2040
aaaaaaaa aaaaaa 2056

```

&lt;210&gt; 64

&lt;211&gt; 81

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 64

Met Lys Val Pro Thr Ser His His Ser Asp Glu Lys His Gln Glu Ala

1

5

10

15

Ser Cys Thr Phe Leu Arg Gly His Ser Arg Ile Asn Pro Pro Leu His

20

25

30

Thr Ala Ala Ile Ser Ile Met His His Ser Ile Ser Gly Tyr Met His

35

40

45

Asn Arg Val Phe Leu Gly Ala Ser Leu Gly Phe Ser Ser Ser Ala Ile

50

55

60

Val Glu Trp Leu His Ser Gln Gly Leu Ala Met Glu Ala His Lys Arg

65

70

75

80

Ala

&lt;210&gt; 65

&lt;211&gt; 581

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 65

```

aaaatatata tgaatagata aatataacta atgtgcgtga gaatgggcct ggcacacctg 60
gggtgctcgg gggtagggct gccatgctgc agagctgatg gagtgatgtt gtgggcacaa 120
ggcatctgtc agctgcctcc tttaacagcc atgccttctg gaatctggaa gaggacacca 180
ctcctgcagt cactgggcag ccacatagca gctgcaggtc ccaggagggc ctgagggtcaa 240
agctgtggaa ggccctcttg gtatcagggc tctctcccca tggcctcttc cccacctctg 300
attgctggtt gtggcccat agctcatcac ccctgtctcc tgctgaatgt gcagggttga 360
tgtggcactg aggagctgac tttggtcaaa ggccaggctc tgcccagcag gggacttgac 420
tgccctgggc gccctgccc tcccgctgcg tgtccaggtc caaagtggag agcctgcctt 480
tgggtttagt tccgaatca gaatccatc tcaccagggc gaaattttaa tttaaaaact 540
taaaaagaga cttttctaaa aaaaaaaaaa aaaaaaaaaa a 581

```

&lt;210&gt; 66

&lt;211&gt; 67

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 66

```

Met Cys Val Arg Met Gly Leu Ala His Leu Gly Cys Ser Gly Val Gly
  1              5              10              15
Leu Pro Cys Cys Arg Ala Asp Gly Val Met Leu Trp Ala Gln Gly Ile
              20              25              30
Cys Gln Leu Pro Pro Leu Thr Ala Met Pro Ser Gly Ile Trp Lys Arg
              35              40              45
Thr Pro Leu Leu Gln Ser Leu Gly Ser His Ile Ala Ala Ala Gly Pro
              50              55              60
Arg Arg Ala
  65

```

&lt;210&gt; 67

&lt;211&gt; 1916

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 67

```

gaaggagaga cggctggcca ccatgcacgg ctctgcagt ttctgatgc ttctgctgcc 60
gctactgcta ctgctggtgg ccaccacagg ccccgttgga gccctcacag atgaggagaa 120
acgttttgatg gtggagctgc acaacctcta ccgggcccag gtatccccga cggcctcaga 180
catgctgcac atgagatggg acgaggagct ggccgccttc gccaaaggcct acgcacggca 240
gtgcgtgtgg ggccacaaca aggagcggcg gcgcccgggc gagaatctgt tcgccatcac 300
agatgaggag ccagttacct tcccctaatc gacctatgtt cctatcccaa aatcagcaga 360
caaagtgaca gacaaaacaa aagtgccctc taggagccca gagaactctc tggaccccaa 420
gatgtccctg acagggggcaa gggaaactcct accccatgcc caggaggagg ctgagggtga 480
ggctgagttg cctccttcca gtgaggtctt ggcctcagtt tttccagccc aggacaagcc 540
aggtagctgc caggccacac tggaccacac ggggcacacc tcctccaagt ccctgcccaa 600
tttccccaat acctctgcc cgcctaagtc cacgggtggg cgtgccctgg ctctgcagtc 660
gtccttgcca ggtaaggccc atagcatctg tcccactttc ctctggctc tggaatgtca 720
gtatcctgcc ccagcatagg tggtagagc atggtggggc atgcacctc tgagtaggag 780
cttccctcct ggctgctgcc ccacctcctt gcctgggtgg gtggcgggga catcaggcac 840
agctgtctct aggtgaggc aaagcctctt tcttcccaca ggtgcagagg gccctgacaa 900
gcctagcgtc gtgtcagggc tgaactcggg ccctggteat gtgtggggcc ctctcctggg 960
actactgctc ctgcccctc tgggtgtggc tggaaatttc tgaaggggat accactcaa 1020
ggcaaggcct ggtgaggggg gccctggcct catacccacc tggattgtct tcctccaagt 1080

```

```

gagagaccac agcttctctgg gcaggctcctg ctctgtggcc cagcagcccc ccttcacccc 1140
aacttctggc cagattccag gccagcactc ttgtcctcct gggaggcgctc tacagggcca 1200
gcccttgcca ctgccccagg agtgccttgg ctctgggtag gcccatcctt cagctggctg 1260
cagactgttc tgagcgggtat ttacatgtgc ccactctcag gttgtcctgt ggccatcagc 1320
ttctctccca gacagaggat ctcaggcttc ccaggaacct ccggggccctt cccagtcctc 1380
tggcctcttc cttgagccat ctgagtcag gactgttccc cagaagtgcc tcttgccctc 1440
tcagggtgaa gaggtcagct gtctcctgt catcttccc accctgtccc cagccccctaa 1500
acaagatact tcttggttaa ggccctccgg aagggaagg ctacggggca tgtgcctcat 1560
cacaccatcc atcctggagg cacaaggcct ggctggctgc gagctcagga ggccgcctga 1620
ggactgcaca cggggccac acctctcctg cccctccctc ctgagtcctg ggggtgggag 1680
gatttgaggg agctcactgc ctacctggcc tggggctgtc tgcccacaca gcatgtgcgc 1740
tctcctgag tgcctgtgta gctggggatg gggattccta ggggcagatg aaggacaagc 1800
cccactggag tggggttctt tgagtggggg aggcagggac gaggaagga aagtaactcc 1860
tgactctcca ataaaaacct gtccaacctg tgaaaaaaaa aaaaaaaaaa aaaaaa 1916

```

&lt;210&gt; 68

&lt;211&gt; 238

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 68

```

Met His Gly Ser Cys Ser Phe Leu Met Leu Leu Leu Pro Leu Leu Leu
  1              5              10              15

Leu Leu Val Ala Thr Thr Gly Pro Val Gly Ala Leu Thr Asp Glu Glu
      20              25              30

Lys Arg Leu Met Val Glu Leu His Asn Leu Tyr Arg Ala Gln Val Ser
  35              40              45

Pro Thr Ala Ser Asp Met Leu His Met Arg Trp Asp Glu Glu Leu Ala
  50              55              60

Ala Phe Ala Lys Ala Tyr Ala Arg Gln Cys Val Trp Gly His Asn Lys
  65              70              75              80

Glu Arg Gly Arg Arg Gly Glu Asn Leu Phe Ala Ile Thr Asp Glu Glu
      85              90              95

Pro Val Thr Phe Pro Lys Ser Thr His Val Pro Ile Pro Lys Ser Ala
  100              105              110

Asp Lys Val Thr Asp Lys Thr Lys Val Pro Ser Arg Ser Pro Glu Asn
  115              120              125

Ser Leu Asp Pro Lys Met Ser Leu Thr Gly Ala Arg Glu Leu Leu Pro
  130              135              140

His Ala Gln Glu Glu Ala Glu Ala Glu Leu Pro Pro Ser Ser
  145              150              155              160

Glu Val Leu Ala Ser Val Phe Pro Ala Gln Asp Lys Pro Gly Glu Leu
      165              170              175

Gln Ala Thr Leu Asp His Thr Gly His Thr Ser Ser Lys Ser Leu Pro
  180              185              190

Asn Phe Pro Asn Thr Ser Ala Thr Ala Asn Ala Thr Gly Gly Arg Ala
  195              200              205

```

Leu Ala Leu Gln Ser Ser Leu Pro Gly Lys Ala His Ser Ile Cys Pro  
 210 215 220

Thr Phe Leu Leu Ala Leu Glu Cys Gln Tyr Pro Ala Pro Ala  
 225 230 235

<210> 69  
 <211> 2051  
 <212> DNA  
 <213> Homo sapiens

<400> 69  
 gggctcgtcg gccgctcctg gaggcggcgg cgaggagcgca gggggcgcgcc ggccccgggga 60  
 ctcgatttcc ccggttcccc ctccacccca cgcggcctgg accatggacg ccagatgggtg 120  
 ggcagtggtg gtgctggctg cgttccccct cctaggggca ggtggggaga ctcccgaagc 180  
 ccctccggag tcatggaccc agctatggtt cttccgattt gtggtgaatg ctgctggcta 240  
 tgccagcttt atggtacctg gctacctcct ggtgcagtac ttcaggcgga agaactacct 300  
 ggagaccggg aggggcctct gctttccctt ggtgaaagct tgtgtgtttg gcaatgagcc 360  
 caaggcctct gatgagggtc ccctggcgcc ccgaacagag gcggcagaga ccaccccgat 420  
 gtggcaggcc ctgaagctgc tcttctgtgc cacagggtc cagggtgtctt atctgacttg 480  
 ggggtgtgctg caggaaagag tgatgacctg cagctatggg gccacagcca catcaccggg 540  
 tgagcgcttt acggactcgc agttcctggt gctaataaac cgagtgtctg cactgattgt 600  
 ggctggcctc tctgtgttc tctgcaagca gccccggcat ggggcaccca tgtaccgta 660  
 ctcttttgc agcctgtcca atgtgcttag cagctggtgc caatacgaag ctcttaagtt 720  
 cgtcagcttc cccacccagg tgctggccaa ggcctctaag gtgatccctg tcatgctgat 780  
 gggaaagctt gtgtctcggc gcagctacga acaactggag tacctgacag ccacactcat 840  
 ctccattggg gtcagcatgt ttctgctatc cagcggacca gagccccgca gctccccagc 900  
 caccacactc tcaggcctca tcttactggc aggttatatt gcttttgaca gcttcacctc 960  
 aaactggcag gatgccctgt ttgcctataa gatgtcatcg gtgcagatga tgtttggggg 1020  
 caattttctt tctgctctt tcacagtggg ctactgcta gaacaggggg ccctactgga 1080  
 gggaaacccg ttcattgggg gacacagtga gtttgctgcc catgccctgc tactctccat 1140  
 ctgtcccgca tgtggccagc tcttcatctt ttacaccatt gggcagtttg gggctgccgt 1200  
 cttcaccatc atcatgacct tccgccaggc ctttgccatc cttctttcct gccttctcta 1260  
 tggccacact gtcactgtgg tgggagggct ggggtgggt gtggtctttg ctgccctcct 1320  
 gctcagagtc tacgcgcggg gccgtctaaa gcaacgggga aagaaggctg tgcctgttga 1380  
 gtctcctgtg cagaagggtt gaggtggaa agggcctgag ggtgaaagt aaataggacc 1440  
 ctcccacat ccccttctgc tgtaacctct gagggagctg gctgaaagg caaaatgcag 1500  
 gtgttttctc agtatcacag accagctctg cagcagggga ttggggagcc caggaggcag 1560  
 ctttcccttt tgccttaagt caccatctt ccagtaagca gtttattctg agccccgggg 1620  
 gtagacagtc ctcaagtggg ggttttgggg agtttgggg caagagagca taggtaggtt 1680  
 ccacagttac tcttccaca agttccctta agtcttgccc tagctgtgct ctgccacctt 1740  
 ccagactcac tccctctgc aaataacctgc atttcttacc ctggtgagaa aagcacaagc 1800  
 ggtgtaggct ccaatgctgc tttcccagga ggggtgaagat ggtgctgtgc tgaggaaagg 1860  
 ggatgcagag ccctgcccag caccaccacc tcctatgctc ctggatccct aggtctctgtt 1920  
 ccatgagcct gttgcaggtt ttggtacttt agaaatgtaa ctttttgctc ttataatttt 1980  
 attttattaa attaaattac tgcaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040  
 aaaaaaaaaa a 2051

<210> 70  
 <211> 432  
 <212> PRT  
 <213> Homo sapiens

<400> 70  
 Met Asp Ala Arg Trp Trp Ala Val Val Val Leu Ala Ala Phe Pro Ser  
 1 5 10 15  
 Leu Gly Ala Gly Gly Glu Thr Pro Glu Ala Pro Pro Glu Ser Trp Thr  
 20 25 30

Gln Leu Trp Phe Phe Arg Phe Val Val Asn Ala Ala Gly Tyr Ala Ser  
 35 40 45  
 Phe Met Val Pro Gly Tyr Leu Leu Val Gln Tyr Phe Arg Arg Lys Asn  
 50 55 60  
 Tyr Leu Glu Thr Gly Arg Gly Leu Cys Phe Pro Leu Val Lys Ala Cys  
 65 70 75 80  
 Val Phe Gly Asn Glu Pro Lys Ala Ser Asp Glu Val Pro Leu Ala Pro  
 85 90 95  
 Arg Thr Glu Ala Ala Glu Thr Thr Pro Met Trp Gln Ala Leu Lys Leu  
 100 105 110  
 Leu Phe Cys Ala Thr Gly Leu Gln Val Ser Tyr Leu Thr Trp Gly Val  
 115 120 125  
 Leu Gln Glu Arg Val Met Thr Arg Ser Tyr Gly Ala Thr Ala Thr Ser  
 130 135 140  
 Pro Gly Glu Arg Phe Thr Asp Ser Gln Phe Leu Val Leu Met Asn Arg  
 145 150 155 160  
 Val Leu Ala Leu Ile Val Ala Gly Leu Ser Cys Val Leu Cys Lys Gln  
 165 170 175  
 Pro Arg His Gly Ala Pro Met Tyr Arg Tyr Ser Phe Ala Ser Leu Ser  
 180 185 190  
 Asn Val Leu Ser Ser Trp Cys Gln Tyr Glu Ala Leu Lys Phe Val Ser  
 195 200 205  
 Phe Pro Thr Gln Val Leu Ala Lys Ala Ser Lys Val Ile Pro Val Met  
 210 215 220  
 Leu Met Gly Lys Leu Val Ser Arg Arg Ser Tyr Glu His Trp Glu Tyr  
 225 230 235 240  
 Leu Thr Ala Thr Leu Ile Ser Ile Gly Val Ser Met Phe Leu Leu Ser  
 245 250 255  
 Ser Gly Pro Glu Pro Arg Ser Ser Pro Ala Thr Thr Leu Ser Gly Leu  
 260 265 270  
 Ile Leu Leu Ala Gly Tyr Ile Ala Phe Asp Ser Phe Thr Ser Asn Trp  
 275 280 285  
 Gln Asp Ala Leu Phe Ala Tyr Lys Met Ser Ser Val Gln Met Met Phe  
 290 295 300  
 Gly Val Asn Phe Phe Ser Cys Leu Phe Thr Val Gly Ser Leu Leu Glu  
 305 310 315 320  
 Gln Gly Ala Leu Leu Glu Gly Thr Arg Phe Met Gly Arg His Ser Glu  
 325 330 335  
 Phe Ala Ala His Ala Leu Leu Leu Ser Ile Cys Ser Ala Cys Gly Gln  
 340 345 350

Leu Phe Ile Phe Tyr Thr Ile Gly Gln Phe Gly Ala Ala Val Phe Thr  
 355 360 365  
 Ile Ile Met Thr Leu Arg Gln Ala Phe Ala Ile Leu Leu Ser Cys Leu  
 370 375 380  
 Leu Tyr Gly His Thr Val Thr Val Val Gly Gly Leu Gly Val Ala Val  
 385 390 395 400  
 Val Phe Ala Ala Leu Leu Leu Arg Val Tyr Ala Arg Gly Arg Leu Lys  
 405 410 415  
 Gln Arg Gly Lys Lys Ala Val Pro Val Glu Ser Pro Val Gln Lys Val  
 420 425 430

&lt;210&gt; 71

&lt;211&gt; 2557

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

tatatgacat tgtccacaga aacaaatata aagattggat caaaggatgg agaaatttta 60  
 atgcagggca tgaacatcct gatcaaaact catttacttt tgctcccaat ggtgtgcctt 120  
 tcattactga ggctctgtac gggccaaagt acaccttctt caacaatgtt ttgatgtttt 180  
 cccagctgt gtcaaagagc tgcttttctc cctgggtggg tcaggtcaca gaagactgct 240  
 catcaaaatg gtctaataac aagcatgacc tggcagctag ttgtcagggg aggggtggtg 300  
 cagcagagga gaaaaatggg gtggttttca tccgaggaga aggtgtggga gcttataacc 360  
 cccagctcaa cctgaagaat gttcagagga atctcactct cctacatcca cagctgcttc 420  
 tccttgtaga ccaatacac ctgggagagg agagtccctt ggagacagca gcgagcttct 480  
 tccataatgt ggatgttcct tttgaggaga ctgtggtaga tgggtgccat ggggctttca 540  
 tcaggcagag agatgggtctc tataaaatgt actggatgga cgatactggc tacagcgaga 600  
 aagcaacctt tgctcagtg acatatactc ggggctatcc ctacaacggg acaaactatg 660  
 tgaatgtcac catgcacctc cgaagtccca tcaccagggc agcttacctc ttcatagggc 720  
 catctataga tgttcagagc ttcactgtcc acggagactc tcagcaactg gatgtgttca 780  
 tagccaccag caaacatgcc tacgccacat acctgtggac aggtgaggcc acaggacagt 840  
 ctgcctttgc acaggtcatt gctgategtc acaaaattct gtttgaccgg aattcagcca 900  
 tcaagagcag cattgtccct gaggtgaagg actatgctgc tattgtggaa cagaacttgc 960  
 agcattttta accagtgttt cagctgctgg agaagcagat actgtcccga gtccggaaca 1020  
 cagctagctt taggaagact gctgaacgcc tgctgagatt ttcagataag agacagactg 1080  
 aggaggccat tgacaggatt tttgccatat cacagcaaca gcagcagcaa agcaagtcaa 1140  
 agaaaaaccg aagggcaggc aaacgtata aatttgtgga tgctgtccct gatatttttg 1200  
 cacagattga agtcaatgag aaaaagatta gacagaaagc tcagattttg gcacagaaag 1260  
 aactacccat agatgaagat gaagaaatga aagacctttt agattttgca gatgtaacat 1320  
 acgagaaaaca taaaaatggg ggcttgatta aaggccggtt tggacaggca cggatggtga 1380  
 caactacaca cagcagggcc ccatcactgt ctgcttccta taccagggtt ttcctgattc 1440  
 tgaacattgc tattttcttt gtcattgttg caatgcaact gacttatttc cagagggcc 1500  
 agagcctaca tggccaaaga tgtctttatg cagttcttct catagatagc tgtattttat 1560  
 tatggttgta ctcttcttgt tcccaatcac agtgttagca ctgaagctat aaattacctg 1620  
 gtcattttgt gatcacaaga gtctatgcaa aaaaaaaat ttctttacc cagattatca 1680  
 gattttttct cctcagattc attttaacaa attaaggga gatattttga cacaagaaag 1740  
 cagggaacgtg gagaaattgg agcaggaaaa gaaattatca aagcaataga aatagcttgg 1800  
 tggctctatg gtgttttttg aagtatttgg cattgcta atagcagtc atatatgtact 1860  
 acttttagaa gaaacaaaa gtctattttt taaagtaat ttttttctta tgagaaaaag 1920  
 gtttagatag aattgggttt tattaatatt aatttaatgc tattagcaat ttccatatac 1980  
 tatattgttg aaaagactga agaatacaat tctgagaaat ataaaaaaat tttaatggta 2040  
 tactcatgtt gaaagataaa tgttgctaag tcctggtatg atgggtgtgag cttccttggg 2100  
 gaagtacttc ttgagttatg taactaacag gatgttttac tacagatctg gatggctatt 2160  
 cagataacat ggcaaaaaat gatagcagaa gatcattaaa aacttaaaat atattttatt 2220

agaaaacatt	tatctatgaa	tgaatatctt	cttgatgctg	gtctctgcac	acatatgctt	2280
ggttactctg	atgcatcctt	tgggtgtctc	ataagtgcga	tgattacaga	taatactgta	2340
ttttctctat	atggaaaacc	gttatagacc	caataacaac	taaacctctt	aaaagaaaat	2400
attttctatt	atgaatgttg	attttcatac	caaagaagat	ggagagtcta	aaatttggtg	2460
atgattctta	tgttttttta	atagaaaaac	ttcttcaagt	ttattttcct	aaataaacat	2520
cataattgtg	aaaaaaaaaa	aaaaaaaaaa	aaaaaaa			2557

```
<210> 72
<211> 474
<212> PRT
<213> Homo sapiens
```

<400> 72																
Met	Phe	Ser	Pro	Ala	Val	Ser	Lys	Ser	Cys	Phe	Ser	Pro	Trp	Val	Gly	
1				5					10					15		
Gln	Val	Thr	Glu	Asp	Cys	Ser	Ser	Lys	Trp	Ser	Lys	Tyr	Lys	His	Asp	
			20					25					30			
Leu	Ala	Ala	Ser	Cys	Gln	Gly	Arg	Val	Val	Ala	Ala	Glu	Glu	Lys	Asn	
	35						40					45				
Gly	Val	Val	Phe	Ile	Arg	Gly	Glu	Gly	Val	Gly	Ala	Tyr	Asn	Pro	Gln	
	50					55					60					
Leu	Asn	Leu	Lys	Asn	Val	Gln	Arg	Asn	Leu	Ile	Leu	Leu	His	Pro	Gln	
65					70					75					80	
Leu	Leu	Leu	Leu	Val	Asp	Gln	Ile	His	Leu	Gly	Glu	Glu	Ser	Pro	Leu	
				85					90					95		
Glu	Thr	Ala	Ala	Ser	Phe	Phe	His	Asn	Val	Asp	Val	Pro	Phe	Glu	Glu	
			100					105					110			
Thr	Val	Val	Asp	Gly	Val	His	Gly	Ala	Phe	Ile	Arg	Gln	Arg	Asp	Gly	
		115					120					125				
Leu	Tyr	Lys	Met	Tyr	Trp	Met	Asp	Asp	Thr	Gly	Tyr	Ser	Glu	Lys	Ala	
	130					135					140					
Thr	Phe	Ala	Ser	Val	Thr	Tyr	Pro	Arg	Gly	Tyr	Pro	Tyr	Asn	Gly	Thr	
145					150					155					160	
Asn	Tyr	Val	Asn	Val	Thr	Met	His	Leu	Arg	Ser	Pro	Ile	Thr	Arg	Ala	
			165						170					175		
Ala	Tyr	Leu	Phe	Ile	Gly	Pro	Ser	Ile	Asp	Val	Gln	Ser	Phe	Thr	Val	
		180						185					190			
His	Gly	Asp	Ser	Gln	Gln	Leu	Asp	Val	Phe	Ile	Ala	Thr	Ser	Lys	His	
	195						200					205				
Ala	Tyr	Ala	Thr	Tyr	Leu	Trp	Thr	Gly	Glu	Ala	Thr	Gly	Gln	Ser	Ala	
	210					215					220					
Phe	Ala	Gln	Val	Ile	Ala	Asp	Arg	His	Lys	Ile	Leu	Phe	Asp	Arg	Asn	
225				230						235					240	
Ser	Ala	Ile	Lys	Ser	Ser	Ile	Val	Pro	Glu	Val	Lys	Asp	Tyr	Ala	Ala	
				245					250					255		



Ile Val Glu Gln Asn Leu Gln His Phe Lys Pro Val Phe Gln Leu Leu  
 260 265 270  
 Glu Lys Gln Ile Leu Ser Arg Val Arg Asn Thr Ala Ser Phe Arg Lys  
 275 280 285  
 Thr Ala Glu Arg Leu Leu Arg Phe Ser Asp Lys Arg Gln Thr Glu Glu  
 290 295 300  
 Ala Ile Asp Arg Ile Phe Ala Ile Ser Gln Gln Gln Gln Gln Ser  
 305 310 315 320  
 Lys Ser Lys Lys Asn Arg Arg Ala Gly Lys Arg Tyr Lys Phe Val Asp  
 325 330 335  
 Ala Val Pro Asp Ile Phe Ala Gln Ile Glu Val Asn Glu Lys Lys Ile  
 340 345 350  
 Arg Gln Lys Ala Gln Ile Leu Ala Gln Lys Glu Leu Pro Ile Asp Glu  
 355 360 365  
 Asp Glu Glu Met Lys Asp Leu Leu Asp Phe Ala Asp Val Thr Tyr Glu  
 370 375 380  
 Lys His Lys Asn Gly Gly Leu Ile Lys Gly Arg Phe Gly Gln Ala Arg  
 385 390 395 400  
 Met Val Thr Thr Thr His Ser Arg Ala Pro Ser Leu Ser Ala Ser Tyr  
 405 410 415  
 Thr Arg Leu Phe Leu Ile Leu Asn Ile Ala Ile Phe Phe Val Met Leu  
 420 425 430  
 Ala Met Gln Leu Thr Tyr Phe Gln Arg Ala Gln Ser Leu His Gly Gln  
 435 440 445  
 Arg Cys Leu Tyr Ala Val Leu Leu Ile Asp Ser Cys Ile Leu Leu Trp  
 450 455 460  
 Leu Tyr Ser Ser Cys Ser Gln Ser Gln Cys  
 465 470

&lt;210&gt; 73

&lt;211&gt; 3442

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 73

agcgccatag ccttaggact atcggtcaca tcctcgcgct cctgctccgg ctctccatc 60  
 ttggcctcgg cagtggcggc tgccgggagg atgtgccgcc ttctggcagg gggaagaagg 120  
 aggagaagat gaagaagcac cggcgggcct tggccctggt ctctgcctc tttctgtgct 180  
 ctctggtctg gcttcccagc tggcgtgtat gttgtaaaga gagttcttca gcttcagcgt 240  
 catcatatta ctctcaagat gacaactgcg cactagaaaa tgaagatgta caattccaga 300  
 aaaagggtgcc ttaaataaaag ttaacattat aatttgtgtg tcagctttct gagagtgtct 360  
 gaaaacattc agaataaaag taatttcaaa aatgccatgc cataattttt ggtggagatt 420  
 cgtatttttc tttgtatta ctgaggtaat tcttaaatga tgacggctca gagatattta 480  
 tatgttttatt cttgttgctt gtgattggaa ttttgtaaca tccttagtcc tgtagtatag 540  
 ggtagtgcat gttcagagag ctgaaaatga gtgatttgtt tttaatgata tataatagag 600

```

tttttaaaga aaccccaatt ttgcaaatga gaactattag aagtgagtat tccagggaga 660
atctacagtc agtggctttg tagtttgttt tatgtaatta ggtgcagtaa atataaagac 720
tccaaaaaac ctcatggctt taaaacatgt attattttac cattctaaat gataaaatgga 780
gagtattgac ttgatttggc ttttgccctt aacttcagtg tattttattt aaattactga 840
agtaattttat attgggtataa aatattttatt atttatgtct cattcccttc cccaaaggaa 900
caacagtccta ctatttcttg tgaaaacggt tggacctgtg tctgtgggta tagataaacac 960
ctttatattt gccacaaaat taatattttt atttccatga aaatatcata taaatatatt 1020
acaaattgcc atcccccttc cctccagtg atagctcata tatcatcttt tcatgttgat 1080
atattctgaa tggctgcatt ttaatcttgt tatatagaca ttatttatca gttgttcttt 1140
attgacaaac ttttgcattt tgttccctgc catgtgtgca cacatgtggg tgtgtgtatc 1200
aacacatact tacaattaca tacttgaggg ttttttgggc atatttagag tggattattt 1260
aaatttaagt tatgttggtt atatttctgg gatggagcat ttaaaatttt gaagacttgc 1320
cgcatacttc aaaaaaact ataattttat attttcacc acacagtata taagattttg 1380
atttaatccc ctttttgatg aagcattttg caaacaaatg tcataaaagt gaatttaagc 1440
aaagcattaa gggggaaagc atgttctttc agtaaacatt tagataccta gtatgagcta 1500
gataggctca ggtaaacaaa gattgaatat gaacttttat gaaaattaca ctctagaagg 1560
cataaaggta ttagaaattc taacttgaat gtatgtcaaa aaagagatga aagaaagagc 1620
tttatgtatc ttcatattgc aggtattcat tggagtttta ctagggtgctt acaacgggtgc 1680
tggatggtaa gactaaaatg cttattaaga aatcaaatga agttcttgaa ctgggtgtatt 1740
gcacatttta aatgtttagt caatgagggg attcaaggaa atagacagta atcttgtata 1800
caaatgaaat taattagagt caaataaaca aggcagtttt ttggttttta agtatggaaa 1860
aaaaccatct tctgtattgt ttataactgt tttgtaaaac agttttatca ggtgattttt 1920
gttgttgttg ttatatagga tgaagagag ggacctatca atgccgaatc attgggaaaa 1980
tcaggttcaa atttacctat ttctccaaa gaacataaat taaaagatga ttctattgtg 2040
gatgtacaag taagctatgt cgctttgatt ttcaataata tgtcatttca aactacttta 2100
caagattgaa aaccttttgt caccatattg tgtgtgtatt gttagttttt tcaactttgag 2160
gtactctgta actggactta agattactta cctgctaata gtactacttt tgagaacatg 2220
taaaattaca gataataata aatgtgacta gtctcttggt agtaaaagtt tgagtataaa 2280
tcctcatttc ttctcgggtt ctatttttgt tcattatgat gtatcttgtc tcttcagatt 2340
ttcagttgtt aagaaatttt ttcttaacct gaatcacttg gaacaaaagg gagtattttt 2400
atgtagaaca cagagtattt tgataaccag gatttccact tatttggggt atgaatagat 2460
aatagatcaa aatatgtttt ttcccttttt ctctcatatt acattctgaa tattgtttat 2520
tttgtgcacc agactttatg aaaaatacta aaaactagaa tatatacacc ataaactgat 2580
cagaatgctg aggtatctgg aaaccatgta ataccagtaa tggctgaagg tactggggat 2640
attttatctg taggcagaag gactaaggaa aaactaaaat ctgcaatatt aaaaaagaa 2700
cacacatttc aactcagaat gagggagaat tctttctttt taaaaaata cagccttggc 2760
taattttttt ttcatatttt attttttaac cttgtacttt ttgcaagagc cagggtattt 2820
ttataactta ctgagttatc taaagcaata gaattagtgt cattttattg gattctggaa 2880
gacatttgtc aagggtttat gcataggagt aagagaaaaa tgatacttgt agcccttaca 2940
attgtatgac tatcctgaac actcctaccg ttattcataa aacttttatt atgatgtagt 3000
tgtgtcctaaa tacttccgag tgcttaccat gtcttgata cctatgttgt tcattacagt 3060
agccactagc cacatgtggc cattaaagcac ttgaaatgcy gctagtccaa attgaaattt 3120
actgtaagtg taaaatacac attggatttt gaagacttaa ttaaaaaata gcatgtaggc 3180
cgggcacagt ggatcacctg aggtcaggag ttcgagacca gcctggccaa catggtgaaa 3240
ccctgtctct accaaaaata caaaaattag ccaggtgtgg tgggtgcacgc ctatagtctc 3300
agctactcag gaggtgagg caggagaatt gcttgaaccc agggaggcga gggtgcagt 3360
agccgagatc atgccactgc actccagcct ggggcaacag agtgagactc tgtctcaaaa 3420
aaaaaaaaa aaaaaaaaaa aa 3442

```

&lt;210&gt; 74

&lt;211&gt; 61

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 74

Met Lys Lys His Arg Arg Ala Leu Ala Leu Val Ser Cys Leu Phe Leu  
1 5 10 15

Cys Ser Leu Val Trp Leu Pro Ser Trp Arg Val Cys Cys Lys Glu Ser  
20 25 30

Ser Ser Ala Ser Ala Ser Ser Tyr Tyr Ser Gln Asp Asp Asn Cys Ala  
 35 40 45

Leu Glu Asn Glu Asp Val Gln Phe Gln Lys Lys Val Pro  
 50 55 60

<210> 75

<211> 1159

<212> DNA

<213> Homo sapiens

<400> 75

gagcggcgcc ggtggcgcc gggagactgg gagggcttcc ggggctgccg gtctgagtgc 60  
 agagctgctg tcatggcgcc cgetctgtgg ggcttcttcc cegtcctgct gctgctgctg 120  
 ctatcggggg atgtccagag ctccggaggtg cccggggctg ctgctgaggg atcgggaggg 180  
 agtggggctg gcataggaga tcgcttcaag attgaggggc gtgcagttgt tccaggggtg 240  
 aagcctcagg actggatctc ggcggcccga gtgctggtag acggagaaga gcacgtcggg 300  
 ttctttaaga cagatgggag ttttgtggtt catgatatac cttctggatc ttatgtagt 360  
 gaagtgttat ctccagctta cagatttgat cccgttcgag tggatatcac ttcgaaagga 420  
 aaaaatgagag caagatatgt gaattacatc aaacatcag aggttgtcag actgccctat 480  
 cctctccaaa tgaatcttc aggtccacct tctacttta ttaaaagga atcgtggggc 540  
 tggacagact ttctaataa cccaatggtt atgatgatgg ttcttcttctt attgatattt 600  
 gtgcttctgc ctaaagtgt caacacaagt gatcctgaca tgagacggga aatggagcag 660  
 tcaatgaata tgctgaattc caaccatgag ttgcctgatg tttctgagtt catgacaaga 720  
 ctcttctctt caaaatcatc tggcaaatct agcagcggca gcagtaaac aggcaaaagt 780  
 ggggctggca aaaggaggtg gtcaggccgt ccagagctgg catttgaca aacacggcaa 840  
 cactgggtgg catccaagtc ttggaaaacc gtgtgaagca actactataa acttgagtca 900  
 tcccgacgtt gatctcttac aactgtgtat gttaactttt tagcacatgt tttgtacttg 960  
 gtacacgaga aaaccagct ttcacttttt gtctgtatga ggtcaatatt gatgtcactg 1020  
 aattaattac agtgcctat agaaaatgcc attaataaat tatatgaact actatacatt 1080  
 atgtatatta attaaaacat cttaatccag aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140  
 aaaaaaaaaa aaaaaaaaaa 1159

<210> 76

<211> 242

<212> PRT

<213> Homo sapiens

<400> 76

Met Ala Ala Ala Leu Trp Gly Phe Phe Pro Val Leu Leu Leu Leu Leu  
 1 5 10 15  
 Leu Ser Gly Asp Val Gln Ser Ser Glu Val Pro Gly Ala Ala Ala Glu  
 20 25 30  
 Gly Ser Gly Gly Ser Gly Val Gly Ile Gly Asp Arg Phe Lys Ile Glu  
 35 40 45  
 Gly Arg Ala Val Val Pro Gly Val Lys Pro Gln Asp Trp Ile Ser Ala  
 50 55 60  
 Ala Arg Val Leu Val Asp Gly Glu Glu His Val Gly Phe Leu Lys Thr  
 65 70 75 80  
 Asp Gly Ser Phe Val Val His Asp Ile Pro Ser Gly Ser Tyr Val Val  
 85 90 95  
 Glu Val Val Ser Pro Ala Tyr Arg Phe Asp Pro Val Arg Val Asp Ile

100	105	110
Thr Ser Lys Gly Lys Met Arg Ala Arg Tyr Val Asn Tyr Ile Lys Thr		
115	120	125
Ser Glu Val Val Arg Leu Pro Tyr Pro Leu Gln Met Lys Ser Ser Gly		
130	135	140
Pro Pro Ser Tyr Phe Ile Lys Arg Glu Ser Trp Gly Trp Thr Asp Phe		
145	150	155
Leu Met Asn Pro Met Val Met Met Met Val Leu Pro Leu Leu Ile Phe		
165	170	175
Val Leu Leu Pro Lys Val Val Asn Thr Ser Asp Pro Asp Met Arg Arg		
180	185	190
Glu Met Glu Gln Ser Met Asn Met Leu Asn Ser Asn His Glu Leu Pro		
195	200	205
Asp Val Ser Glu Phe Met Thr Arg Leu Phe Ser Ser Lys Ser Ser Gly		
210	215	220
Lys Ser Ser Ser Gly Ser Ser Lys Thr Gly Lys Ser Gly Ala Gly Lys		
225	230	235
		240

Arg Arg

<210> 77  
 <211> 2462  
 <212> DNA  
 <213> Homo sapiens

<400> 77  
 agaccggaag tgagtgatcg aaagcatggc gtcggtggtg ttggcgctga ggaccgggac 60  
 agccgtttaca tccttgctaa gcccactcc ggctacagct cttgctgtca gatacgcatc 120  
 caagaagtcg ggtggtagct ccaaaaacct cgggtggaaag tcatcaggca gacgccaagg 180  
 cattaagaaa atggaaggtc actatgttca tgctgggaac atcattgcaa cacagcgcca 240  
 tttccgctgg caccaggtg cccatgtgag ttgctcgtt gctgcccccc ttttccctt 300  
 tctagggtga cctctccttg cccctaagca tggtaataac agttgcatgt attgagtgt 360  
 taccaaatgg caagcattgt gctgattccc atgcctacac gatctcattt cttccttacc 420  
 acatccctgt aagtaagggt aaatgccaga gacctagacg ggggtggaagg agcaagtgc 480  
 tgctgggatt tgcaccaggt ctgcctaact ccagatcac tatgatttgc cctggtgttg 540  
 cattggcctg gtattcttgg ttctccttcc taacctcag ttcttggagg acaagatacc 600  
 tgggtgattta aaatattatt ttagtcttgg gaaactaatt tcaatttatt agtttttcat 660  
 tattctaaga cttcttcctt tgtataaact tacttgcaga tgggtgaaag atagcttgaa 720  
 tttaatgaaa tagaattcag tgtgccagga gttagggttca taccagggtg ggatttcagt 780  
 ttcagcaagt tggcatcatt tttatagggt aggacagttc tttgttacag agacataacg 840  
 atactgtact ctccacatta ttttaaata tagttgcctt tgtaaacagt tccccctccc 900  
 ccaccctttt ttcgataatt ttagtaggta caaaaagggt gatagtaacc tgtatttatt 960  
 tcaaagatgc taatttgtat gtgaattgat tatgtatgta gtactttttt accttttaga 1020  
 acttaagaca ttagatgtgt ggctatgtca acaataataa tcaaagctaa cacattttga 1080  
 gctagacagt attctaggcc ctttatgatt gttgagcatt aatcctcaa atgaccctgt 1140  
 aaaatagtta atattatccc catttgacag acgaaaaaat agaggcacag tgaagtaact 1200  
 tgcctaagac cacacaggag taagtagcag tgctggtatt tgaaccaggc atcctggctc 1260  
 cagagcccat tcttttaate agagagacag agaataatgc tggactacct gggtaaaatt 1320  
 tgccacacac ccagcctgaa gttctaaaag aaaaagctaa aaaatatggt taaaaactgt 1380  
 atatacaaca cagtttaate tctctgcttc atatttcaca tctgaaaagt ggcgttgga 1440

```

ataatacctc tttcactagg tgagaattaa atgacttaaat gtgtattaaa gggcttacat 1500
gggtattagc taatgcagta gtatgattgt tttattagca tatgcattat ccatatagct 1560
ttggtttata ttttccatag agcctaaact ttatagatca ctatctaaat gaaaatttac 1620
atcaaaactag tgttttaata tgacagttct agtggtgttt aagtctaat gaagatttag 1680
ggcatcttcc tgggttttgg gatttgtgca gtgtggcttc tgaccagctg catagacatg 1740
gtccaggta acagtccttg ggactgagca gtggcccatg ttggctgttg gagctttctt 1800
ttccctcatt tgttatgagt caataaagaa gcagaagggt tggacgtgct agccagtggg 1860
aaagggaacc tgagatggag ctctctgac accctttcag cccagggcc cggtctgtg 1920
tgagaagtct gtcccaaaat tctgggacgg accacctcac tgaccagctg accagctgac 1980
caggctctgt tgttgagggt ggggtgttgg aagaataaat gtctgtatgc cctggaagag 2040
gggatagctc gctacactaa ggaggtctac gtgcctcatc ccagaaacac ggaggctgtg 2100
gatctgatca ccaggctgcc caagggtgct gtgctctaca agacttttgt ccacgtggtt 2160
cctgccaagc ctgagggcac cttcaaaact gtagctatgc tttgatgtcc tgttgaggcc 2220
atcggacaga gactggagcc cagggtgacag gagatggtga taccagaagt caagggttgg 2280
ggtggcgaca cggcctccc aggaagagggt ctgcttgatg gtgactctgc aggagactct 2340
gaagtgcagt ctgggaaacc ctttgggaga cctgacctgg ggccaaaaat aaagtgcagc 2400
agcgtcatga acgcatgcta tttagggaca aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2460
aa

```

&lt;210&gt; 78

&lt;211&gt; 94

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 78

```

Met Ala Ser Val Val Leu Ala Leu Arg Thr Arg Thr Ala Val Thr Ser
  1              5              10             15

Leu Leu Ser Pro Thr Pro Ala Thr Ala Leu Ala Val Arg Tyr Ala Ser
      20              25             30

Lys Lys Ser Gly Gly Ser Ser Lys Asn Leu Gly Gly Lys Ser Ser Gly
      35              40             45

Arg Arg Gln Gly Ile Lys Lys Met Glu Gly His Tyr Val His Ala Gly
      50              55             60

Asn Ile Ile Ala Thr Gln Arg His Phe Arg Trp His Pro Gly Ala His
      65              70             75             80

Val Ser Cys Ser Val Ala Ala Pro Leu Phe Pro Phe Leu Gly
      85              90

```

&lt;210&gt; 79

&lt;211&gt; 1178

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 79

```

taacttattt ttaaaggata ttttacaata atttatgata atatgcttaa agatatacta 60
ttctttcatt tttttctttt aggtttgttt tctgccctag acagtcattc aaatttgggt 120
gtcatttctt atggtgtttc catgacgact ttggaagacg tatttttttaa gctagaagtt 180
gaagcagaaa ttgaccaagc aggtaaaaac agaactaaca aaacatttta ggcaatgaat 240
cagacagatg gtgaataaga caaaaattat ataaatgtaa ttttatattt tttttagaat 300
aaagggtaga aataagtaaa taagtaatga gatacttact gatagtgaata ggagtaatgc 360
aggaaataat ggggtggtct tagaaggcag cgaacttaga ttatatcatc aggaaggact 420
ttgttgagat gatacttgac ctgaggcctg tataatgaga aaaatcaagt aggtgaagat 480
tttagggcag aaattttcag ataaagaact gcaactgcaa aggtattaac tgaggaatga 540
gttgatgtc tttgagggat aaaaaaaggc taccatggcc ggagccatag tgaagctatt 600

```

```

acacagcaaa gtcaggagga tactcaaagg gatagatcat gttctgcctt tgaacgtctg 660
gaaaacaatt tcatttggtt tectattgta gtgtggaagc tatcagagag ttttaaagca 720
aataggtacc atgatctgat ttgaattttt aaaaaaacac gttctggttg ttgaatgata 780
agagtattat aaaggaagtg tggaagtagg aagaccagtt aggaaactgc tatagttaat 840
actatcttga actagggtgt cagtgaatgt agggaggcgg ggcgcagtgg ttcatgcctg 900
taatcccagc actttgggag gccaaaggcag ggagatcgct tgagcccagg agttccagac 960
cagtcttggc aacatggcaa aaccttgtca caaaaaacac aaaaattagc cgggtatggt 1020
agtgtgtgcc tatagtccca actactctag aggctgcggt gggagaatca cctgaggcca 1080
ggaggtcaag gctgcagtga gccatgattg cgccactgca ctctagcctg ggcatacaga 1140
ggagaccctg tttaaaaaaa aaaaaaaaaa aaaaaaaa 1178

```

&lt;210&gt; 80

&lt;211&gt; 62

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 80

```

Met Leu Lys Asp Ile Leu Phe Phe His Phe Phe Leu Leu Gly Leu Phe
  1             5             10            15

```

```

Ser Ala Leu Asp Ser His Ser Asn Leu Gly Val Ile Ser Tyr Gly Val
      20             25            30

```

```

Ser Met Thr Thr Leu Glu Asp Val Phe Leu Lys Leu Glu Val Glu Ala
      35             40            45

```

```

Glu Ile Asp Gln Ala Gly Lys Asn Arg Thr Asn Lys Thr Phe
      50             55            60

```

&lt;210&gt; 81

&lt;211&gt; 1285

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 81

```

aaaggttcca gaaacagatg aggtagagct cacctgccag gctacagggt atcctctggc 60
agaagtatcc tggccaaacg tcagcgttcc tgccaacacc agccactcca ggaccctga 120
aggcctctac caggtcacca gtgttctgcg cctaaagcca ccccttgcca gaaacttcag 180
ctgtgtgttc tggaatactc acgtgaggga acttactttg gccagcattg accttcaaag 240
tcagatggaa cccaggaccc atccaacttg gctgcttcac attttcatcc cctcctgcat 300
cattgctttc attttcatag ccacagtgat agccctaaga aaacaactct gtcaaaagct 360
gtattcttca aaagacacaa caaaaagacc tgtcaccaca acaaagaggg aagtgaacag 420
tgctatctga acctgtggtc ttgggagcca gggtgacctg atatgacatc taaagaagct 480
tctggactct gaacaagaat tcggtggcct gcagagcttg ccatttgcac ttttcaaatg 540
cctttggatg acccagcact ttaatctgaa acctgcaaca agactagcca acacctggcc 600
atgaaaacttg ccccttcact gatctggact cacctctgga gcctatggct ttaagcaagc 660
actactgcac tttacagaat taccctactg gatcctggac ccacagaatt ccttcaggat 720
ccttcttgct gccagactga aagcaaaagg aattatttcc cctcaagttt tctaagtgat 780
ttccaaaagc agaggtgtgt ggaaatttcc agtaacagaa acagatgggt tgccaataga 840
gttatttttt atctatagct tcctctgggt actagaagag gctattgaga ctatgagctc 900
acagacaggg cttcgacaaa actcaaatca taattgacat gttttatgga ttactggaat 960
cttgatagca taatgaagt gttctaatta acagagagca tttaaatata cactaagtgc 1020
acaaattgtg gagtaaatgc atcaagctct gtttttgagg tctaagtcac aaagcatttg 1080
ttttaacctg taatggcacc atgtttaatg gtggtttttt tttttgaact acatctttcc 1140
tttaaaaatt attggtttct ttttatttgt ttttacctta gaaatcaatt atatacagtc 1200
aaaaataaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1260
aaaaaaaaaa aaaaaaaaaa aaaaaa 1285

```

&lt;210&gt; 82

<211> 61  
 <212> PRT  
 <213> Homo sapiens

<400> 82  
 Met Glu Pro Arg Thr His Pro Thr Trp Leu Leu His Ile Phe Ile Pro  
 1 5 10 15  
 Ser Cys Ile Ile Ala Phe Ile Phe Ile Ala Thr Val Ile Ala Leu Arg  
 20 25 30  
 Lys Gln Leu Cys Gln Lys Leu Tyr Ser Ser Lys Asp Thr Thr Lys Arg  
 35 40 45  
 Pro Val Thr Thr Thr Lys Arg Glu Val Asn Ser Ala Ile  
 50 55 60

<210> 83  
 <211> 654  
 <212> DNA  
 <213> Homo sapiens

<400> 83  
 gagcatagac caccaggctg agtatcctga cctgagtcac cccagggat caggagcctc 60  
 cagcagggaa ccttccatta tattcttcaa gcaacttaca gctgcaccga cagttgcat 120  
 gaaagtctta atctcttccc tctctctgtt gctgccacta atgctgatgt ccatggctctc 180  
 tagcagcctg aatccagggg tcgccagagg ccacagggac cgaggccagg cttctaggag 240  
 atggctccag gaaggcggcc aagaatgtga gtgcaaagat tggttcctga gagccccgag 300  
 aagaaaattc atgacagtgt ctgggctgcc aaagaagcag tgccccctgtg atcatttcaa 360  
 gggcaatgtg aagaaaacaa gacaccaaag gcaccacaga aagccaaaca agcattccag 420  
 agcctgccag caatttctca aacaatgtca gctaagaagc ttgtctctgc cttttagtagga 480  
 gctctgagcg cccactcttc caattaaaca ttctcagcca agaagacagt gagcacacct 540  
 accagacact cttcttctcc cactcactc tcccactgta cccacccta aatcattcca 600  
 gtgctctcaa aaagcatggt tttcaagatc aaaaaaaaaa aaaaaaaaaa aaaa 654

<210> 84  
 <211> 119  
 <212> PRT  
 <213> Homo sapiens

<400> 84  
 Met Lys Val Leu Ile Ser Ser Leu Leu Leu Leu Leu Pro Leu Met Leu  
 1 5 10 15  
 Met Ser Met Val Ser Ser Ser Leu Asn Pro Gly Val Ala Arg Gly His  
 20 25 30  
 Arg Asp Arg Gly Gln Ala Ser Arg Arg Trp Leu Gln Glu Gly Gly Gln  
 35 40 45  
 Glu Cys Glu Cys Lys Asp Trp Phe Leu Arg Ala Pro Arg Arg Lys Phe  
 50 55 60  
 Met Thr Val Ser Gly Leu Pro Lys Lys Gln Cys Pro Cys Asp His Phe  
 65 70 75 80  
 Lys Gly Asn Val Lys Lys Thr Arg His Gln Arg His His Arg Lys Pro  
 85 90 95

Asn Lys His Ser Arg Ala Cys Gln Gln Phe Leu Lys Gln Cys Gln Leu  
 100 105 110

Arg Ser Phe Ala Leu Pro Leu  
 115

<210> 85  
 <211> 1176  
 <212> DNA  
 <213> Homo sapiens

<400> 85  
 aatctttttt taaataagta ttttggtgag gttgaagaat tgctggcaat taaaagaata 60  
 gagctaatta tggctttcat cattcattca tgtatttatt gagcacctac ttattatggg 120  
 gctcaacact tgttactgca agctacctta atttccaag agtgggtgct tactctgttt 180  
 tttctgatat ggtcttccaa tcagtgtgtg taacatacct gttgtttatc agccattgta 240  
 ggtggctgtg tctgttgcac catcataaga agtttaagct ttgtgctctg ataaattgtg 300  
 ttctgttaaa gaggttagta ggatgaaaac agcaaaacaa taattttttc aacaaattgt 360  
 aaattataag aaaaagagtt ggtttgtgta caacaatttt aatgattccc ttgttcattt 420  
 ttgctgtgaa atgcactgaa aaaaatcctc aaaatgagtt atagttcctg tgttgggaaa 480  
 attgacaaat aataaaacta gagaacaaac aataatgctt ctgtctcttt tacgaatgga 540  
 gagagaaagt ttatattcag tagagttatt gccctgttca ttgagaggg gcattggattt 600  
 tctgtttaag tccttcagga atcttcagct aggtggtaaa ttaataaga gtttctaaaa 660  
 attgaaatgt ttaactttta aatattctgg agatagaaga agaataataa atgaaaccag 720  
 gctgatctgc atgcagtggc atttacaact aactgatcac aaccaattat agattcctta 780  
 ttttgtttat tgtgaggcag agtctgactc tgtcaccag gatggagtac agtgcatagc 840  
 tcactgcagt cttgacctcc caggctaaac caatcattcc acttcacct cccaagtagc 900  
 tgagaccaca ggcacacaac accacaacca gctgattggt gtactgtttg tatagactgg 960  
 atctcactat gttgccaga ctgggtctga attcctgagc tcaagcagtc ctcccacctc 1020  
 agcctcccaa agtactggga ttacaggcgt gaggtaccct gccagagccc agttacagat 1080  
 ttctttgttc cttctctctc ccactgctta acttgattag cctttaaaaa agaaataaat 1140  
 aaaaattttt aaaaaaaaaa aaaaaaaaaa aaaaaa 1176

<210> 86  
 <211> 78  
 <212> PRT  
 <213> Homo sapiens

<400> 86  
 Met Tyr Leu Leu Ser Thr Tyr Leu Leu Trp Cys Ser Thr Leu Val Thr  
 1 5 10 15  
 Ala Ser Tyr Leu Asn Phe Pro Arg Val Val Pro Tyr Ser Val Phe Ser  
 20 25 30  
 Asp Met Val Phe Gln Ser Val Cys Val Thr Tyr Leu Leu Phe Ile Ser  
 35 40 45  
 His Cys Arg Trp Leu Cys Leu Leu His His His Lys Lys Phe Lys Leu  
 50 55 60  
 Cys Ala Leu Ile Asn Cys Val Leu Leu Lys Arg Leu Val Gly  
 65 70 75

<210> 87  
 <211> 1476  
 <212> DNA  
 <213> Homo sapiens



&lt;400&gt; 87

gaggctgcag aggtgccatg gctgagtcac acctgctgca gtggctgctg ctgctgctgc 60  
 ccacgctctg tggcccaggc actgctgcct ggaccacctc atccttggcc tgtgccagg 120  
 gccctgagtt ctggtgccaa agcctggagc aagcattgca gtgcagagcc ctagggcatt 180  
 gcttacagga agtctgggga catgtgggag ccgatgacct atgccaagag tgtgaggaca 240  
 tcgtccacat ccttaacaag atggccaagg aggccatttt ccaggatctc tccgagcagc 300  
 aattccccat tcctctcccc tattgctggc ctctgcaggg ctctgatcaa gcggatccaa 360  
 gccatgattc ccaaggggtg gctagctgtg gcagtggccc aggtgtgccc cgtggtagct 420  
 ctgggtggcgg gcgcacatctg ccagtgcctg gctgagcgct actccgtcat cctgctcgac 480  
 acgctgctgg gccgcatgct gcccagctg gtctgccgcc tcgtcctccg gtgctccatg 540  
 gatgacagcg ctggcccaag agaattggctg ccgcgagact ctgagtgcc cctctgcatg 600  
 tccgtgacca cccaggccgg gaacagcagc gagcaggcca taccacaggc aatgctccag 660  
 gcctgtgttg gctcctggct ggacagggaa aagtgcgaagc aatttggtga gcagcacacg 720  
 ccccagctgc tgacctggt gcccaggggc tgggatgccc acaccacctg ccaggccctc 780  
 ggggtgtgtg ggaccatgtc cagccctctc cagtgtatcc acagccccga cctttgatga 840  
 gaactcagct gtccagaaaa agacacgctc ctttaaagtg ctgcagtatg gccagacgtg 900  
 gtggctcaca cctgcaatcc cagcacctta ggaggccgag gcaggaggat ccttgaggtc 960  
 aggagtccga gaccagcctc gccaacatgg tgaaacccca tttctactaa aaatacaaaa 1020  
 aattagccaa gtgtgggtgc atatgcctgt aatcccaact actcagaagg ccgaggcagg 1080  
 agaattactt gaacgcagga gaatcactgc agcccaggag gcagagggtt cagtgcagcc 1140  
 agattgcacc actgcactcc agcctgggtg acagagcaag actccatctc agtaaataaa 1200  
 taaataaata aaaagcgtg cagtgcctgt ggccctaccc tgaagtgcagc gggcccaggc 1260  
 ctacctcact ctctcccttg gcagagaagc agacgtccat agctcctctc cctcacaagc 1320  
 gctcccagcc tgccctccag ctgctgctct cccctcccag tctctactca ctgggatgag 1380  
 gttaggtcat gaggacacca aaaaccta aaataacaaa aagccaaaca agccttagct 1440  
 tttcttaaag acaaaaaaaaa aaaaaaaaa aaaaaa 1476

&lt;210&gt; 88

&lt;211&gt; 145

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 88

Met Ala Glu Ser His Leu Leu Gln Trp Leu Leu Leu Leu Leu Pro Thr  
 1 5 10 15  
 Leu Cys Gly Pro Gly Thr Ala Ala Trp Thr Thr Ser Ser Leu Ala Cys  
 20 25 30  
 Ala Gln Gly Pro Glu Phe Trp Cys Gln Ser Leu Glu Gln Ala Leu Gln  
 35 40 45  
 Cys Arg Ala Leu Gly His Cys Leu Gln Glu Val Trp Gly His Val Gly  
 50 55 60  
 Ala Asp Asp Leu Cys Gln Glu Cys Glu Asp Ile Val His Ile Leu Asn  
 65 70 75 80  
 Lys Met Ala Lys Glu Ala Ile Phe Gln Asp Leu Ser Glu Gln Gln Phe  
 85 90 95  
 Pro Ile Pro Leu Pro Tyr Cys Trp Pro Leu Gln Gly Ser Asp Gln Ala  
 100 105 110  
 Asp Pro Ser His Asp Ser Gln Gly Cys Ala Ser Cys Gly Ser Gly Pro  
 115 120 125  
 Gly Val Pro Arg Gly Thr Ser Gly Gly Gly Arg His Leu Pro Val Pro  
 130 135 140

Gly  
145

<210> 89  
<211> 2243  
<212> DNA  
<213> Homo sapiens

<400> 89  
agttactttgt aatcaagtga aaatataact ttatttttta actctattac attttatttt 60  
gtcatgtact aaaattatatt ctgtattgct ttataaaaa acagtggcat ttagcactgg 120  
cattgagact atagcacatc atttttgcca ttttcagtgc ttatattgtt aggtagaggc 180  
tggcacttta ttagaatgca agccacaaaa atatcaattt tgtttttttt gttaggggtg 240  
gtctcttttt ttctctttcc tctctctttt tttaacaaat gccttcttat agaaaaactt 300  
tctaagaggc aacaatttag aatggatatt ttgacgaatc ggcagtgtg taacagtgtat 360  
aacctgatct gtttgtttta aagattatta ccaagtgaat aattcagaat gaatagaatt 420  
tacactaaca tgctatataa aatgttaaag tctgatgctg tgaaagcaat ctagtgtctat 480  
atttctacct cctcatttgt cttaattatt tggtaagtgg gattatgatg agtaactgga 540  
ggggccttaga aacaaaaact ggatgaaaga gtatgcatga agaaaagctt ctttgataaa 600  
tgtggagttc ttcatataaa atatatattc atgaattcac agataagtac ttaaagaaca 660  
gacagtttac ttggcctaaa aatattttga tgtttactca aaaagtacct cttcaggtct 720  
tgagaacatg gaaaagaatt gagtgtttt aaatactttt tagaaagtaa tcataaaagt 780  
aaattgaatt tcaaacctat ttggcttctg ttttgtgaac ctttgaacta tatgtatgtg 840  
tataagggtg tacacatata tatatggcat ataacaagtg tacacatata cacataacaa 900  
gtgtagaagt atatatataa tacatacact cactctgtct ggtataggct aattttgaa 960  
aactcccata agtttctgct gcttctccca taactgtctg caccaccatc agaattcata 1020  
atcaaaccta acctttttgt ttggggcacc aaatctgaag acaaaattaa tttgcaccag 1080  
taaaactcaa gctgtcttct ttcttgaaaa ctaaacgttt aacgtataat gtctgttttg 1140  
atactgttcc aaattgttga ttgcatgtgg ttaatgttgc atttaggcac tttgcaattg 1200  
cataattcat taatgttttg tgagcttgca tttgtgagtt attggatgat cagactgaat 1260  
tttgtcaagt atcacattgt acatcttgcc tagatgtcga tgactgcaag taataatata 1320  
gtttataatg aaactatcta caattcttgt ttttagcacat ctgttatccg taaaacacct 1380  
gtaactagct tttttaattt attatttgaa ttttaggata gcgaatcact aatttttagt 1440  
tgctgagggt ggcatttttag tgattattaa gcacttctgt cagtctttga aaaaagaacg 1500  
tattttttgt gctttgaaga tctctgaaga atttctttta taatagaatg ggcagtgtat 1560  
gtaacagttt tatgtcaaat gatctgtgct gtagaaaaac attaacctt gttcaaaaaa 1620  
gaaatggata aacttggcct ttctaagtgg taagaatgac ctgtcactat aatatactgt 1680  
atgtttacat tttatttaaa ttttatctct tatgtatagg gtgataacct tcccagaaa 1740  
caacagtgtat tgcgattgtt ttctagaaac ttctttaaag tgccacattt ggcagtacaa 1800  
atgagtctga gtgtaatagc ccagagattt atatatagtt gaatgtctaa aatggtaaaa 1860  
tgtgccactg tgtcaagtta cagtggctta tgtttttcat agtaattcaa atgaacttcc 1920  
tatttttgat agtaaatgtc atttaatagt atacttgcca tttgagcctc actgcaaaa 1980  
tagtgcagag gagaaaacaa tttttaatgt aatcttgatt ttacctcata tactgtacat 2040  
tccaaaaact ctaaaacttt taaagattat agatacacta ccaaacatat caccttaaaa 2100  
ttgtataagg ctgaatgaac ttcatacaaa tgaaaaaaat ctcataaaaa tacataaact 2160  
atgtagcaaa agtatctgta aaatccatgg aaaataaaag ttgtatcatt ctttttgaaa 2220  
aaaaaaaaa aaaaaaaaaa aaa 2243

<210> 90  
<211> 61  
<212> PRT  
<213> Homo sapiens

<400> 90  
Met Gln Ala Thr Lys Ile Ser Ile Leu Phe Phe Leu Leu Gly Trp Val  
1 5 10 15  
Phe Phe Phe Ser Phe Pro Leu Ser Phe Phe Asn Lys Cys Leu Leu Ile

20 25 30

Glu Lys Leu Ser Lys Arg Gln Gln Phe Arg Met Asp Ile Leu Thr Asn  
35 40 45

Arg His Glu Cys Asn Ser Asp Asn Leu Ile Cys Leu Phe  
50 55 60

<210> 91  
<211> 1041  
<212> DNA  
<213> Homo sapiens

<400> 91  
aaatgggctg tcttcacggtg tggaatacaa cataatggag ttggaacaag aacttgaaaa 60  
tgtaaagact cttaaagaaa aattagagag gcgaaaaaag gcttcagcat gggaaaagaaa 120  
tttggtgtat cccgctgtta tgggtctcct tcttattgag acatccatct cggtcctctt 180  
ggtggcttgt aatattcttt gcctattggt tgatgaaaca gcaatgccaa aaggaacaag 240  
ggggcctgga ataggaaatg cctctctttc tacgtttggt tttgtgggag ctgcgcttga 300  
aatcattttg attttctatc ttatggtgtc ctctgtgtgc ggcttctata gccttcgatt 360  
ttttggaaac ttactccca agaaagatga cacaactatg acaaagatca ttggaaattg 420  
tgtgtccatc ttggttttga gctctgctct gcctgtgatg tcgagaacac tgggaatcac 480  
tagatttgat ctacttggtg actttggaag gtttaattgg ctgggaaatt tctatattgt 540  
attatcctac aatttgcttt ttgctattgt gacaacattg tgtctggtcc gaaaattcac 600  
ctctgcagtt cgagaagaac ttttcaaggc cctagggcct cataaacttc acttaccaaa 660  
tacttcaagg gattcagaaa cagccaagcc ttctgtaaat gggcatcaga aagcactgtg 720  
agacgcacag acggcgtctt ctgccaccaa gagacccgag aactccagat tcacgacatt 780  
cctgtcccat gtagaagcat ttccattcaa ccgtggcccc tcttcagaac cttagacctat 840  
cagtgcatt ttttttcat aatctacgaa gaacttggt atggctgac ttttttaaat 900  
ttaactttct gatggacct gtagtttcca gttaagtga gattccttac agacatatag 960  
aacagcgcat tcttctgtag acatttgctc atgttggtta atacaatcac ccatatgaaa 1020  
aaaaaaaaa aaaaaaaaaa a 1041

<210> 92  
<211> 228  
<212> PRT  
<213> Homo sapiens

<400> 92  
Met Glu Leu Glu Gln Glu Leu Glu Asn Val Lys Thr Leu Lys Thr Lys  
1 5 10 15  
Leu Glu Arg Arg Lys Lys Ala Ser Ala Trp Glu Arg Asn Leu Val Tyr  
20 25 30  
Pro Ala Val Met Val Leu Leu Leu Ile Glu Thr Ser Ile Ser Val Leu  
35 40 45  
Leu Val Ala Cys Asn Ile Leu Cys Leu Leu Val Asp Glu Thr Ala Met  
50 55 60  
Pro Lys Gly Thr Arg Gly Pro Gly Ile Gly Asn Ala Ser Leu Ser Thr  
65 70 75 80  
Phe Gly Phe Val Gly Ala Ala Leu Glu Ile Ile Leu Ile Phe Tyr Leu  
85 90 95  
Met Val Ser Ser Val Val Gly Phe Tyr Ser Leu Arg Phe Phe Gly Asn  
100 105 110

Phe Thr Pro Lys Lys Asp Asp Thr Thr Met Thr Lys Ile Ile Gly Asn  
 115 120 125  
 Cys Val Ser Ile Leu Val Leu Ser Ser Ala Leu Pro Val Met Ser Arg  
 130 135 140  
 Thr Leu Gly Ile Thr Arg Phe Asp Leu Leu Gly Asp Phe Gly Arg Phe  
 145 150 155 160  
 Asn Trp Leu Gly Asn Phe Tyr Ile Val Leu Ser Tyr Asn Leu Leu Phe  
 165 170 175  
 Ala Ile Val Thr Thr Leu Cys Leu Val Arg Lys Phe Thr Ser Ala Val  
 180 185 190  
 Arg Glu Glu Leu Phe Lys Ala Leu Gly Leu His Lys Leu His Leu Pro  
 195 200 205  
 Asn Thr Ser Arg Asp Ser Glu Thr Ala Lys Pro Ser Val Asn Gly His  
 210 215 220  
 Gln Lys Ala Leu  
 225

<210> 93  
 <211> 1792  
 <212> DNA  
 <213> Homo sapiens

<400> 93  
 atgggacggg acttgcgcc cggtgccgc gtgctcctgc tcctgcttct gctcctgctg 60  
 gtgtacctga ctcagccagg caatggcaac gagggcagcg tcaactggaag ttgttattgt 120  
 ggtaaaagaa tttcttcga ctcgccgcca tcggttcagt tcatgaatcg tctccgaaa 180  
 cacctgagag cttaccatcg gtgtctatac tacacgaggt tccagctcct ttcctggagc 240  
 gtgtgtggag gcaacaagga cccatgggtt caggaattga tgagctgtct tgatctcaa 300  
 gaatgtggac atgcttactc ggggattgtg gccaccaga agcatttact tctaccagc 360  
 ccccaactt ctcaggcctc agagggggca tcttcagata tccacacccc tgcccagatg 420  
 ctctgtcca cctgcagtc cactcagcgc cccaccctcc cagtaggagc actgtcctcg 480  
 gacaagagc tcaactcgtc caatgaaacc accattcaca ctgcccggcca cagtctggca 540  
 gttgggcctg aggtcgggga gaaccagaag cagccggaaa aaaatgctgg tcccacagcc 600  
 aggacatcag ccacagtgc ggtcctgtgc ctcctggcca tcatcttcat cctcacgca 660  
 gccctttcct atgtgctgtg caagaggagg agggggcagt caccgcagtc ctctccagat 720  
 ctgcccgttc attatatacc tgtggcacct gactctaata cctgagccaa gaatggaagt 780  
 ttgtgaggag acggactcta tgttgcccag gctgttatgg aactcctgag tcaagtgatc 840  
 ctcccacctt ggcctctgaa ggtgcgagga ttataggcgt cactaccac atccagccta 900  
 cacgtatttg ttaatatcta acataggact aaccagccac tgccctctct taggcccctc 960  
 atttaaaaac ggttatacta taaaatctgc ttttcacact gggtgataat aacttggaca 1020  
 aattctatgt gtattttgtt ttgttttgc ttgctttgtt ttgagacgga gtctcgctct 1080  
 gtcatccagg ctggagtga gtggcatgat ctggctcac tgcaaccccc atctcccagg 1140  
 ttcaagcgat tctcctgcct cctcctaagt agctgggact acaggtgctc accaccacac 1200  
 ccggctaatt tttgtattt ttagtagaga cggggtttca ccatgttgac caggctggtc 1260  
 tcgaactcct gacctggtga tctgccacc caggcctccc aaagtgctgg gattaaagg 1320  
 gtgagccacc atgctggcc ctatgtgtgt ttttaacta ctaaaaatta tttttgta 1380  
 gattgagctc tctttatgga acaactggc ctcagccctt gcgcccttac tgtgattcct 1440  
 ggcttcattt tttgctgatg gttccccctc gtcccaaatc tctctcccag tacaccagtt 1500  
 gttcctcccc cactcagcc ctctcctgca cctcctgta cccgcaacga aggcctgggc 1560  
 tttcccacc tccctcttag caggtgccgt gctgggacac catacgggtt ggtttcacct 1620  
 ctcagtcctc ttgcctaccc cagtgaagat ctgatcttgt ttttattgtt attgctttta 1680

ttattattgc ttttattatc attaaaactc tagttcttgt tttgtctctc cgaaaaaaaa 1740  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 1792

<210> 94  
 <211> 254  
 <212> PRT  
 <213> Homo sapiens

<400> 94  
 Met Gly Arg Asp Leu Arg Pro Gly Ser Arg Val Leu Leu Leu Leu Leu  
           1                  5                  10                  15  
  
 Leu Leu Leu Leu Val Tyr Leu Thr Gln Pro Gly Asn Gly Asn Glu Gly  
                   20                  25                  30  
  
 Ser Val Thr Gly Ser Cys Tyr Cys Gly Lys Arg Ile Ser Ser Asp Ser  
           35                  40                  45  
  
 Pro Pro Ser Val Gln Phe Met Asn Arg Leu Arg Lys His Leu Arg Ala  
           50                  55                  60  
  
 Tyr His Arg Cys Leu Tyr Tyr Thr Arg Phe Gln Leu Leu Ser Trp Ser  
           65                  70                  75                  80  
  
 Val Cys Gly Gly Asn Lys Asp Pro Trp Val Gln Glu Leu Met Ser Cys  
                   85                  90                  95  
  
 Leu Asp Leu Lys Glu Cys Gly His Ala Tyr Ser Gly Ile Val Ala His  
                   100                  105                  110  
  
 Gln Lys His Leu Leu Pro Thr Ser Pro Pro Thr Ser Gln Ala Ser Glu  
           115                  120                  125  
  
 Gly Ala Ser Ser Asp Ile His Thr Pro Ala Gln Met Leu Leu Ser Thr  
           130                  135                  140  
  
 Leu Gln Ser Thr Gln Arg Pro Thr Leu Pro Val Gly Ser Leu Ser Ser  
           145                  150                  155                  160  
  
 Asp Lys Glu Leu Thr Arg Pro Asn Glu Thr Thr Ile His Thr Ala Gly  
                   165                  170                  175  
  
 His Ser Leu Ala Val Gly Pro Glu Ala Gly Glu Asn Gln Lys Gln Pro  
                   180                  185                  190  
  
 Glu Lys Asn Ala Gly Pro Thr Ala Arg Thr Ser Ala Thr Val Pro Val  
           195                  200                  205  
  
 Leu Cys Leu Leu Ala Ile Ile Phe Ile Leu Thr Ala Ala Leu Ser Tyr  
           210                  215                  220  
  
 Val Leu Cys Lys Arg Arg Arg Gly Gln Ser Pro Gln Ser Ser Pro Asp  
           225                  230                  235                  240  
  
 Leu Pro Val His Tyr Ile Pro Val Ala Pro Asp Ser Asn Thr  
                   245                  250

<210> 95  
 <211> 1234

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 95

```

gagactccag atttccctgt caaccacgag gagtccagag aggaaacgcg gagcggagac 60
aacagtacct gacgcctctt tcagcccggy atcgccccag cagggatggg cgacaagatc 120
tggtgcctt tccccgtgct ccttctggcc gctctgcctc cgggtgctgt gcctggggcg 180
gccggcttca caccttcctt cgatagcgac ttcaccttta cccttccgc cggccagaag 240
gagtgcttct accagcccat gccctgaag gcctcgctgg agatcgagta ccaagtttta 300
gatggagcag gattagatat tgatttccat cttgcctctc cagaaggcaa aacctagtt 360
tttgaacaaa gaaaatcaga tggagtccac actgtagaga ctgaagttgg tgattacatg 420
ttctgctttg acaatacatt cagcaccatt tctgagaagg tgattttctt tgaattaatc 480
ctggataata tgggagaaca ggcacaagaa caagaagatt ggaagaaata tattactggc 540
acagatatat tggatatgaa actggaagac atcctggaat ccatcaacag catcaagtcc 600
agactaagca aaagtgggca catacaaatt ctgcttagag catttgaagc tcgtgatcga 660
aacatacaag aaagcaactt tgatagagtc aatttctggt ctatggttaa tttagtggtc 720
atggtggtgg tgtcagccat tcaagtttat atgtggaaga gtctgtttga agataagagg 780
aaaagtagaa cttaaaactc caaactagag tacgtaacat tgaaaaatga ggcataaaaa 840
tgcaataaac tgttacagtc aagaccatta atggtcttct ccaaaatatt ttgagatata 900
aaagtaggaa acaggtataa ttttaatgtg aaaattaagt cttcactttc tgtgcaagta 960
atcctgctga tccagttgta cttaagtgtg taacaggaat attttgcaga atataggttt 1020
aactgaatga agccatatta ataactgcat tttcctaact ttgaaaaatt ttgcaaatgt 1080
cttaggtgat ttaataaaat gagtattggg cctaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa                                     1234

```

&lt;210&gt; 96

&lt;211&gt; 229

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 96

```

Met Gly Asp Lys Ile Trp Leu Pro Phe Pro Val Leu Leu Leu Ala Ala
  1              5              10              15

Leu Pro Pro Val Leu Leu Pro Gly Ala Ala Gly Phe Thr Pro Ser Leu
      20              25              30

Asp Ser Asp Phe Thr Phe Thr Leu Pro Ala Gly Gln Lys Glu Cys Phe
      35              40              45

Tyr Gln Pro Met Pro Leu Lys Ala Ser Leu Glu Ile Glu Tyr Gln Val
      50              55              60

Leu Asp Gly Ala Gly Leu Asp Ile Asp Phe His Leu Ala Ser Pro Glu
      65              70              75              80

Gly Lys Thr Leu Val Phe Glu Gln Arg Lys Ser Asp Gly Val His Thr
      85              90              95

Val Glu Thr Glu Val Gly Asp Tyr Met Phe Cys Phe Asp Asn Thr Phe
      100             105             110

Ser Thr Ile Ser Glu Lys Val Ile Phe Phe Glu Leu Ile Leu Asp Asn
      115             120             125

Met Gly Glu Gln Ala Gln Glu Gln Glu Asp Trp Lys Lys Tyr Ile Thr
      130             135             140

Gly Thr Asp Ile Leu Asp Met Lys Leu Glu Asp Ile Leu Glu Ser Ile

```

145                      150                      155                      160

Asn Ser Ile Lys Ser Arg Leu Ser Lys Ser Gly His Ile Gln Ile Leu  
                                  165                                   170                                   175

Leu Arg Ala Phe Glu Ala Arg Asp Arg Asn Ile Gln Glu Ser Asn Phe  
                                  180                                   185                                   190

Asp Arg Val Asn Phe Trp Ser Met Val Asn Leu Val Val Met Val Val  
                                  195                                   200                                   205

Val Ser Ala Ile Gln Val Tyr Met Leu Lys Ser Leu Phe Glu Asp Lys  
                                  210                                   215                                   220

Arg Lys Ser Arg Thr  
 225

&lt;210&gt; 97

&lt;211&gt; 1204

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 97

```

gtgcttcctg gcaggagagc agatgtgatt ctgagcgaca tggcgcccaa tgccacaggg 60
ttccgggacc tcgatcatga caggctcatc agcctgtgcc tgacccttct cagcgtgacc 120
ccagacatcc tgcaacctgg ggggacattc ctttgtaaaa cctgggctgg aagtcaaagc 180
cgtcgggttac agaggagact gacagaggaa ttccagaatg taaggatcat caaacctgaa 240
gccagcagga aagagtcac agaatgttac ttcttgccca cacagtacca cggaaggaag 300
ggcactgtga agcagtgagg atttcttgtg ccattttcat aatggtcatt agctcctttt 360
aagctagaaa cgtagcctga gctcctgaag agttcctggg agatttgagc tgatttttga 420
gatggagcag gacaagtggg gagtctctct ctctcttctc ctctctctct ttttaaccaa 480
aaagagatga caaaactaag ttcagggggc atggaaaatg aaaaagtccg ctatatgttg 540
atttgggaag agaagttat caagagaaag aggtgaggat ggaaggatgg agaaaaacag 600
actgtgggaa ggatcagaag gaatccgccg aggcagggat ggggtgtgcc atgtgtgcct 660
tgacgggact tcattctata gactgttaaa ctgtcacaca caaacaggct ttccaccctt 720
gctctgagag caccacgcac agatttccag ttcttagtgt ggctgtttta agtagaaaat 780
ctgggggctg ggtgaggcca ctcatgcctg taaaccagg gctttagaag gctgaggctg 840
ggggattgct tgaagtcagg agttcaagac caacctgggc aacatagcaa cccccccat 900
gtctacaaaa atgaaaaacc aaaaagcaaa ccaaaagaaa aatctgaaat ttccatctgg 960
ggattaaact ctgtctttct ggtgaacaat atagcaattc acgcattctt caagcagcaa 1020
aagttcccgg aacaattagg gaagacgtat ggtctgaatt tatccaggca gtgggtctgc 1080
tttggttttt gctggaaatt tatatcagtg tctgggctcc caagaacata aatgtaattg 1140
ccaaagcaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200
aaaaa                                           1204

```

&lt;210&gt; 98

&lt;211&gt; 92

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 98

Met Ala Pro Asn Ala Thr Gly Phe Arg Asp Leu Asp His Asp Arg Leu  
   1                                  5                                  10                                  15

Ile Ser Leu Cys Leu Thr Leu Leu Ser Val Thr Pro Asp Ile Leu Gln  
                                   20                                  25                                  30

Pro Gly Gly Thr Phe Leu Cys Lys Thr Trp Ala Gly Ser Gln Ser Arg  
                                   35                                  40                                  45

Arg Leu Gln Arg Arg Leu Thr Glu Glu Phe Gln Asn Val Arg Ile Ile  
50 55 60

Lys Pro Glu Ala Ser Arg Lys Glu Ser Ser Glu Val Tyr Phe Leu Ala  
65 70 75 80

Thr Gln Tyr His Gly Arg Lys Gly Thr Val Lys Gln  
85 90

<210> 99

<211> 1343

<212> DNA

<213> Homo sapiens

<400> 99

tttttttttt ttttttttga cggagtctcg ctctgttgcc caggctggag 60  
tgcagtggca tgcctcgcac tctactgcaat ctatgcctcc caggttcaag cgattctcct 120  
gccctaggct cccgcatagc tgggattaca ggcaagcacc accacctctg gctgattttt 180  
atatttttag tagagacggg gtttcacat gttggccacg ctatgcctcc aagcttcttg 240  
ataaccgaca agtcaaggcc aaagatttca ggcaccagat accaccaagt gcgtttacca 300  
acattttgat gttttccttt atttatgtcc tgcttttttg catggaaatt gacctctaaa 360  
ctgtacaact cagatcttaa gacagggaaa tacagtgaac actccatatt caccgggttc 420  
acattttgtg attcaaccaa ctacagattg aaaatatttg gaaaaataaa aaggatagtt 480  
gtgtttgtac tgaacatgaa cagatttctt tttgtcatc atttctctaa caatacaaca 540  
gcgatgtaca cacagatcct tagctcattc accaaaatta cctgtgcac cagagcccca 600  
ttttcttacc aatcacatct gcatgtggtg gtgcttttcc agttagaaaa caaaactggt 660  
gtcctgtgtg ctgtcaacca aacaaaactt ttcattgcag aagcgttgtt ccattttatg 720  
tggccgacag agttgtgtta acaaagtatt tggtcagcgt agaaaatctc aagggctcta 780  
gaatttcacag tggaaacttg aagtactgta tcttgaactt gtaaaaagga gggggaacct 840  
caactgtact caggaaatag aatctcaagt gagtttataa aaagctgtga tctcgggtga 900  
ctttctgggc cagcagacag tgggagcagc ccagggggctc tagggagagc cccctgctga 960  
gtgtccctgg ccccatggg aaactttcca ttaaaggcct ggtggatggg gtgggggtag 1020  
tggggatgct aaagacgcat ttcctgagct tgtttctggt tgatacattg atgactcaga 1080  
caagcgtctt ctactcatc tagttcgagc tgctagggaa gaaagcattc aagacctgat 1140  
acttatctga ggagtggcag gaagtctggg gtgacttttc tcaagggtggg gggacaaaga 1200  
gaatgagagg aaaatgaaac agagtgtaaa ccaaaaataa aattccaagt ccgcctacca 1260  
tcggaatgga cccctcctct cagccgaggg cattccaaag tcaacctatg aaactacttc 1320  
aggccatgag aggaagggga gga 1343

<210> 100

<211> 210

<212> PRT

<213> Homo sapiens

<400> 100

Met Ile Ser Thr His Cys Asn Leu Cys Leu Pro Gly Ser Ser Asp Ser  
1 5 10 15

Pro Ala Leu Gly Ser Arg Ile Ala Gly Ile Thr Gly Lys His His His  
20 25 30

Leu Trp Leu Ile Phe Ile Phe Leu Val Glu Thr Gly Phe His His Val  
35 40 45

Gly His Ala Ser Ile Ser Ser Phe Leu Ile Thr Asp Lys Ser Arg Pro  
50 55 60

Lys Ile Ser Gly Thr Arg Tyr His Gln Val Arg Leu Pro Thr Phe Val



```
<210> 101
<211> 1529
<212> DNA
<213> Homo sapiens
```

<400>	101						
gccccggggcc	cccacgtect	ttgagccgcg	gagggccccc	cccttgccgc	ccggcggtgac	60	
ctcagccccc	ggcttcccc	atctgcccac	agccaacccc	acagggcctg	gggagcgggg	120	
cccgcggggc	gcagtgagg	tgatccggga	gtccagcagc	accagggga	tggtgggtgg	180	
cattgtggc	gcgcgcgcg	tctgcattct	catctctctc	tcagccatgt	ataagtaccg	240	
caatcgtyat	gagggctcct	accaggttga	ccagagccga	aactacatca	gtaactcggc	300	
ccagagcaat	ggggcggttg	tgaaagagaa	ggccccggct	gcccccaaga	cgcccagcaa	360	
ggccaagaag	aacaaagaca	aggagtatta	tgtctgagcc	cccggcactg	cgccccactg	420	
ccagctgccc	ctctctgggag	ggcccgggag	gagggtgcca	ccctctccct	gccagggggc	480	
tggggaccct	ctccctggct	gcctcaggct	tctcttacga	agagggagcg	caaaaaaaga	540	
aaaggaaaaa	accgtgtctc	gcccccttcc	tctctgcgtc	cactgcgcgc	ctctcgtcagt	600	
ccccgggctg	actgtccctc	tcagctctgc	gcctgccagg	caggggcactg	gctcacagcc	660	
ctggggtgat	ttattttttt	aagggggtag	ttttattttg	gtggggttgg	gtgggaagga	720	
aggctggggg	ttttgtaaag	tgtccactgc	tcgtctgtgt	aattttctct	aatttttctt	780	
cttcttccct	ctgtccctcc	tgccttccct	ctcttcccaa	gccttccaat	ccccatccca	840	
ggcttgctgt	gtctcactgt	ccccaccctc	cttccctact	tctttttttg	tgtgtcttgt	900	
ttctcccttc	ctttctctcc	tttgggtttc	cagagtctgt	gggagaaggg	cgggaggggt	960	
ggcccagagt	gccccagtgg	tgggtggggg	ggggtggggc	aagtgtccca	actccctca	1020	
caggagaggg	cacctgcttg	gtgccgccca	gggaaggggc	tcaggcctga	cgggaaggcct	1080	
gttctgtgtg	tgccgccggg	cgacgtgcat	tgatggggaa	gctgctggag	gagcaggggt	1140	
gggggggtgg	agggagggga	aaggcaaatg	cagatatata	ttacagacaa	atactctaga	1200	
ttccacgagc	agcagcctgt	ggcacccgct	gggcgcgggc	agcaggggaag	agggcagcag	1260	
gcattgtcca	cagactgtct	gggtcacttc	tttgcccacg	ggctccctgc	tccccagatt	1320	
ttttttctct	ctttgttaac	aaatgtgtct	gagttcttga	aaacacccca	accccgaaaa	1380	
tgtgtgggaa	aaagaaaaaca	aaaactttcc	aaattccaaa	aaaaaaaaaa	aaaaaaaaaa	1440	

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1500  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1529

<210> 102  
 <211> 75  
 <212> PRT  
 <213> Homo sapiens

<400> 102  
 Met Val Val Gly Ile Val Ala Ala Ala Leu Cys Ile Leu Ile Leu  
 1 5 10 15  
 Leu Tyr Ala Met Tyr Lys Tyr Arg Asn Arg Asp Glu Gly Ser Tyr Gln  
 20 25 30  
 Val Asp Gln Ser Arg Asn Tyr Ile Ser Asn Ser Ala Gln Ser Asn Gly  
 35 40 45  
 Ala Val Val Lys Glu Lys Ala Pro Ala Ala Pro Lys Thr Pro Ser Lys  
 50 55 60  
 Ala Lys Lys Asn Lys Asp Lys Glu Tyr Tyr Val  
 65 70 75

<210> 103  
 <211> 733  
 <212> DNA  
 <213> Homo sapiens

<400> 103  
 ttgtgtcacc attcttgttg ctttttaaaa atcaggctaa tcatgtgggc catgtctctt 60  
 caaagcttga cctgcacaaa tgccatattt ctatttggac cacatattct ccattttgca 120  
 ttgagcagta gagtacagtg gaaagggaaat aagaatactg attattctga acagtttagt 180  
 cccaagagaa tagcgtttta aaaaagaaaa acaagatttg gagtcattgt gggttatttt 240  
 tgggtgggatg gaggatctta aaaatgccta attgtgagag aatcaattgc tgaaagtgtt 300  
 aaaatttctg aaaataaatg cttaattaca tatacaggaa ttaaatagtt tggaagaggg 360  
 ttggattatc attaccttta caatactgta taatcagaag ttctctgaac ctcaattgta 420  
 tatctagaca taataattgt tttctgtata ggatgttggt tggtttggtt ctgagtgttt 480  
 aaattttgca aaacacaaatg ttaaatttgt gcttcagtac ctagataaat tggaagaggt 540  
 aatgttctag tttctggaag gtaagcctgg gagacacata agcaattcac tgctataatt 600  
 tagttgatgt aaaatgacgg aaactgactc aatatgtcag gtttaactct gcccaaaagc 660  
 agcagacatg taagcagatg tgcaataaaa aatgatcttg atccatttca aaaaaaaaaa 720  
 aaaaaaaaaa aaa 733

<210> 104  
 <211> 52  
 <212> PRT  
 <213> Homo sapiens

<400> 104  
 Met Trp Ser Met Ser Leu Gln Ser Leu Thr Cys Thr Asn Ala Ile Phe  
 1 5 10 15  
 Leu Phe Gly Pro His Ile Leu His Phe Ala Leu Ser Ser Arg Val Gln  
 20 25 30  
 Trp Lys Gly Asn Lys Asn Thr Asp Tyr Ser Glu Gln Phe Ser Pro Lys  
 35 40 45

Arg Ile Ala Phe

50

&lt;210&gt; 105

&lt;211&gt; 2342

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 105

```

gaaccacaaa acccgccagg ccggtgcggg agctgaggag catccgctgc ggtcctcgcc 60
gagacccccg cgcggattcg ccggtccttc ccgcgggcgc gacagagctg tcctcgacc 120
tggaatggcag caggggcgcc ggggtcctct cgacgccaga gagaaatctc atcatctgtg 180
cagccttctt aaagcaaact aagaccagag ggaggattat ccttgacctt tgaagaccaa 240
aactaaactg aaatttaaaa tgttcttcgg gggagaaggg agcttgactt acacttttgt 300
aataatttgc ttcctgacac taaggctgtc tgctagtcag aattgcctca aaaagagtct 360
agaagatggt gtcattgaca tccagtcac tctttctaag ggaatcagag gcaatgagcc 420
cgtatatact tcaactcaag aagactgcat taattcttgc tgttcaacaa aaaacatata 480
aggggacaaa gcatgtaact tgatgatctt cgacactcga aaaacagcta gacaacccaa 540
ctgctacctt tttttctgtc ccaacgagga agcctgtcca ttgaaaccag caaaaggact 600
tatgagttac aggataatta cagattttcc atctttgacc agaaatttgc caagccaaaga 660
gttacccccag gaagattctc tcttacatgg ccaattttca caagcagtc ctcccctagc 720
ccatcatcac acagattatt caaagccac cgatatctca tggagagaca cactttctca 780
gaagtttggg tccctcagatc acctggagaa actatttaag atggatgaag caagtgccca 840
gtccttctgt tataaggaaa aaggccattc tcagagttca caattttcct ctgatcaaga 900
aatagctcat ctgctgcctg aaaatgtgag tgcgctccca gctacggtgg cagttgcttc 960
tccacatacc acctcggtc ctccaaagcc cgcacacctt ctaccacca atgcttcagt 1020
gacaccttct gggacttccc agccacagct ggccaccaca gctccacctg taaccactgt 1080
cacttctcag cctcccacga ccttcatttc tacagttttt acacgggctg cggctacact 1140
ccaagcaatg gctacaacag cagttctgac taccacctt caggcaccta cggactcgaa 1200
aggcagctta gaaaccatac cgtttacaga aatctccaac ttaactttga acacagggaa 1260
tgtgtataac cctactgcac tttctatgtc aaatgtggag tcttccacta tgaataaaac 1320
tgcttctctg gaaggtaggg aggccagtc aggcagttcc tcccagggca gtgttcaga 1380
aaatcagtac ggccttccat ttgaaaaatg gcttcttata ggtccctgc tctttggtgt 1440
cctgttctct gtgataggcc tcgtcctcct gggtagaatc ctctcggaat cactccgcag 1500
gaaacgttac tcaagactgg attatttgat caatgggatc tatgtggaca tctaaggatg 1560
gaactcgggt tctcttaatt catttagtaa ccagaagccc aaatgcaatg agtttctgct 1620
gacttgctag tcttagcagg aggttggtatt ttgaagacag gaaaatgccc cttctgctt 1680
tctttttttt tttttggaga cagagtcttg ctctgttgcc caggctggag tgcagtagca 1740
cgatctcggc tctcaccgca acctccgtct cctgggttca agcgattctc ctgctcagc 1800
ctcctaagta tctgggatta caggcatgtg ccaccacacc tgggtgattt ttgtattttt 1860
agtagagacg gggtttcacc atgttggtca ggctggtctc aaactcctga cctagtgate 1920
caccctcttc ggcctcccaa agtgctggga ttacaggcat gagccaccac agctggcccc 1980
cttctgtttt atgtttgggt tttgagaagg aatgaagtgg gaaccaaatt aggtaatatt 2040
gggtaaatct tctctaaaat attagctaaa aacaaagctc tatgtaaagt aataaagtat 2100
aattgccata taaatttcaa aattcaactg gcttttatgc aaagaaacag gttaggacat 2160
ctaggttcca attcattcac attcttgggt ccagataaaa tcaactgttt atatcaattt 2220
ctaattggatt tgcttttctt tttatatgga ttcttttaaa acttattcca gatgtagttc 2280
cttccaatta aatatttgaa taaatctttt gttactcaaa aaaaaaaaaa aaaaaaaaaa 2340
aa

```

2342

&lt;210&gt; 106

&lt;211&gt; 431

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 106

Met Phe Phe Gly Gly Glu Gly Ser Leu Thr Tyr Thr Leu Val Ile Ile  
1 5 10 15

Cys Phe Leu Thr Leu Arg Leu Ser Ala Ser Gln Asn Cys Leu Lys Lys  
 20 25 30  
 Ser Leu Glu Asp Val Val Ile Asp Ile Gln Ser Ser Leu Ser Lys Gly  
 35 40 45  
 Ile Arg Gly Asn Glu Pro Val Tyr Thr Ser Thr Gln Glu Asp Cys Ile  
 50 55 60  
 Asn Ser Cys Cys Ser Thr Lys Asn Ile Ser Gly Asp Lys Ala Cys Asn  
 65 70 75 80  
 Leu Met Ile Phe Asp Thr Arg Lys Thr Ala Arg Gln Pro Asn Cys Tyr  
 85 90 95  
 Leu Phe Phe Cys Pro Asn Glu Glu Ala Cys Pro Leu Lys Pro Ala Lys  
 100 105 110  
 Gly Leu Met Ser Tyr Arg Ile Ile Thr Asp Phe Pro Ser Leu Thr Arg  
 115 120 125  
 Asn Leu Pro Ser Gln Glu Leu Pro Gln Glu Asp Ser Leu Leu His Gly  
 130 135 140  
 Gln Phe Ser Gln Ala Val Thr Pro Leu Ala His His His Thr Asp Tyr  
 145 150 155 160  
 Ser Lys Pro Thr Asp Ile Ser Trp Arg Asp Thr Leu Ser Gln Lys Phe  
 165 170 175  
 Gly Ser Ser Asp His Leu Glu Lys Leu Phe Lys Met Asp Glu Ala Ser  
 180 185 190  
 Ala Gln Leu Leu Ala Tyr Lys Glu Lys Gly His Ser Gln Ser Ser Gln  
 195 200 205  
 Phe Ser Ser Asp Gln Glu Ile Ala His Leu Leu Pro Glu Asn Val Ser  
 210 215 220  
 Ala Leu Pro Ala Thr Val Ala Val Ala Ser Pro His Thr Thr Ser Ala  
 225 230 235 240  
 Thr Pro Lys Pro Ala Thr Leu Leu Pro Thr Asn Ala Ser Val Thr Pro  
 245 250 255  
 Ser Gly Thr Ser Gln Pro Gln Leu Ala Thr Thr Ala Pro Pro Val Thr  
 260 265 270  
 Thr Val Thr Ser Gln Pro Pro Thr Thr Leu Ile Ser Thr Val Phe Thr  
 275 280 285  
 Arg Ala Ala Ala Thr Leu Gln Ala Met Ala Thr Thr Ala Val Leu Thr  
 290 295 300  
 Thr Thr Phe Gln Ala Pro Thr Asp Ser Lys Gly Ser Leu Glu Thr Ile  
 305 310 315 320  
 Pro Phe Thr Glu Ile Ser Asn Leu Thr Leu Asn Thr Gly Asn Val Tyr  
 325 330 335

Asn Pro Thr Ala Leu Ser Met Ser Asn Val Glu Ser Ser Thr Met Asn  
 340 345 350

Lys Thr Ala Ser Trp Glu Gly Arg Glu Ala Ser Pro Gly Ser Ser Ser  
 355 360 365

Gln Gly Ser Val Pro Glu Asn Gln Tyr Gly Leu Pro Phe Glu Lys Trp  
 370 375 380

Leu Leu Ile Gly Ser Leu Leu Phe Gly Val Leu Phe Leu Val Ile Gly  
 385 390 395 400

Leu Val Leu Leu Gly Arg Ile Leu Ser Glu Ser Leu Arg Arg Lys Arg  
 405 410 415

Tyr Ser Arg Leu Asp Tyr Leu Ile Asn Gly Ile Tyr Val Asp Ile  
 420 425 430

<210> 107  
 <211> 3153  
 <212> DNA  
 <213> Homo sapiens

<400> 107  
 agcgccttgc caaaatggag cttgtaagaa ggctcatgcc attgaccctc ttaattctct 60  
 cctgttttggc ggagctgaca atggcggagg ctgaaggcaa tgcaagctgc acagtcagtc 120  
 taggggggtgc caatatggca gagaccaca aagccatgat cctgcaactc aatcccagtg 180  
 agaactgcac ctggacaata gaaagaccag aaaacaaagc atcagaatta tcttttccta 240  
 tgtccagctt gatccagatg gaagctgtga aagtgaanaac attaaagtct ttgacggaac 300  
 ctccagcaat gggcctctgc tagggcaagt ctgcagtaaa aacgactatg ttctctgtatt 360  
 tgaatcatca tccagtacat tgacgtttca aatagttact gactcagcaa gaattcaaag 420  
 aactgtcttt gtctcttact acttcttctc tcttaacatc tgtaagtcct catttacaca 480  
 accttcaccc acatctccta caagcacagg ttacacactc ttctgtccgg gttcttataa 540  
 tcttcagtct gagtatattc cctcagagtc ttctctatct gagagagcag tagttctcta 600  
 acttacatgt gcataagaat catctggaga gcttggttaa tcacagatcc tggaccccat 660  
 cacagatatt gtgattcatt ggtctctggat ggagcacatg atcttgactt tctaactagc 720  
 tcccagatga ggcagagaca aggggctgat cagatggcta catattgagg ggtactgtgt 780  
 tagagcccaa agcggagatt ctgcattcag cctgtttaat agtaaccctg gtatcttctg 840  
 caatgcctct tcaggggactg cactctccta tgctctgatt ccttagatct tggttcgggt 900  
 tgttgaacct tctctgttaa caagtccccc ggatgattct gacaaagaag actggagagt 960  
 gcatgctggg aatctctggg ctccaccattg aaatttaaga ctgtcagggtg attggaggct 1020  
 acatttacag ggctctgcat tcacagcacc tacattccac tgtgatccga agcagaatgc 1080  
 caagaacatc tgcgagtggg ttcatgagga gagctccact gtggatttct ttccaaggcc 1140  
 cagagctgac catgtcactc tctgtctaaa accactgact tcttggtacc agcagatctc 1200  
 cagagtgcag cagtcaaggt tttccacgcg tggacccagg ccaccttttc aagccttgcc 1260  
 tctagctgct gcctggcggtg caccctatgc ttacagccaa ctaaaccatt ttacagctcc 1320  
 aaaaagagct ctctgcattg tttctccaca tttcatctgt ccagctgctt ccttatgtac 1380  
 ttactgggga aaggagagac caagggtgc ttcatgaagt agtggcggat gagactggaa 1440  
 agctgagctg gtgagagtga ggaagagcct tcaagggagg gtgtgacttt attttctatg 1500  
 gcgcaattca ggtaattggga gtctctccag atgcaacaca tgttacagca caactttaat 1560  
 atgattcgtt tggcaatagt ataaagattc cctctctgcc ctgcaccagt ggacagaaat 1620  
 ttcaatgata aaaaaggatg aaaacctcaa tatgtgaccc atgggaaaga tagagatgga 1680  
 atcatactca ggagagatgt gttaaactca atgcacaaca cagaaattaa aagcagtctc 1740  
 tattttgatg ttcatctctg accaccttca catccatttc agctattcca aactgtggcg 1800  
 gttacctgga taccttggaa ggatccttca ccagcccaaa ttacccaaag ccgcatcctg 1860  
 agctggctta ttgtgtgtgg cacatacaag tggagaaaga ttacaagata aaactaaact 1920  
 tcaaagagat tttattatgc caattcttac cggggatttt ctgcttccca cacctcaatt 1980  
 ttatgcagaa aacatcaaca ctacatcttt aacttgctct tctgacagga tgagagtat 2040  
 tataagcaaa tctacacatg aggccttttaa ctctaattgg aataacttgc aactaaaaga 2100

```

cccaacttgc agacaaaaat tatcaaatgt tgtggaattt tctgtccctc ttaatggatg 2160
tggtaacaatc agaaaggtag aagatcagtc aattacttac accaatataa tcaccttttc 2220
tgcatcctca acttctgaag tgatcacccg tcagaaacaa ctccagatta ttgtgaagtg 2280
tgaaatggga cataattcta cagtggagat aatatacata acagaagatg atgtaatata 2340
aagtcaaaat gcaactgggca aatataacac cagcatggct ctttttgaat ccaattcatt 2400
tgaaaagact atacttgaat caccatatta tgtggatttg aacaaaactc tttttgttca 2460
agttagtctg cacacctcag atccaaattt ggtggtgttt ctgtatacct gtagagcctc 2520
tcccacctct gactttgcat ctccaacctc cgacctaatc aagagtggat gtagtcgaga 2580
tgaaacttgt aagggtgata ccttatttgg acactatggg agattccagt ttaatgcctt 2640
taaattcttg agaagtatga gctctgtgta tctgcagtgt aaagttttga tatgtgatag 2700
cagtgaccac cagtctcgct gcaatcaagg ttgtgtctcc agaagcaaac gagacatttc 2760
ttcatataaa tggaaaacag attccatcat aggaccatt cgtctgaaaa gggatcgaag 2820
tgcaagtggc aattcaggat ttcagcatga aacacatgcg gaagaaactc caaaccagcc 2880
tttcaacagt gtgcatctgt tttccttcat ggttctagct ctgaatgtgg tgactgtagc 2940
gacaatcaca gtgaggcatt ttgtaaatca acgggcagac tacaataacc agaagctgca 3000
gaactattaa ctaacaggtc caaccctaag tgagacatgt ttctccagga tgccaaagga 3060
aatgctacct cgtggctaca catattatga ataaatgagg aagggcctga aagtgcacac 3120
caggcctgca aaaaaaaaaa aaaaaaaaaa aaa 3153

```

&lt;210&gt; 108

&lt;211&gt; 102

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 108

```

Met Glu Leu Val Arg Arg Leu Met Pro Leu Thr Leu Leu Ile Leu Ser
  1                      5                      10                      15

```

```

Cys Leu Ala Glu Leu Thr Met Ala Glu Ala Glu Gly Asn Ala Ser Cys
          20                      25                      30

```

```

Thr Val Ser Leu Gly Gly Ala Asn Met Ala Glu Thr His Lys Ala Met
    35                      40                      45

```

```

Ile Leu Gln Leu Asn Pro Ser Glu Asn Cys Thr Trp Thr Ile Glu Arg
    50                      55                      60

```

```

Pro Glu Asn Lys Ala Ser Glu Leu Ser Phe Pro Met Ser Ser Leu Ile
    65                      70                      75                      80

```

```

Gln Met Glu Ala Val Lys Val Lys Thr Leu Lys Ser Leu Thr Glu Pro
          85                      90                      95

```

```

Pro Ala Met Gly Leu Cys
          100

```

&lt;210&gt; 109

&lt;211&gt; 1805

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 109

```

acgttccttc tactctggca ccactctcca ggctgccatg gggcccagca cccctctcct 60
catcttggtc cttttgtcat ggtcgggacc cctccaagga cagcagcacc accttggtga 120
gtacatggaa cgccgactag ctgctttaga ggaacggctg gccagtgcc aggaccagag 180
tagtcggcat gctgctgagc tgccgggactt caagaacaag atgctgccac tgctggagg 240
ggcagagaag gagcgggagg cactcagaac tgaggccgac accatctccg ggagagtgga 300
tcgtctggag cgggaggtag actatctgga gaccagaac ccagctctgc cctgtgtaga 360
gtttgatgag aagggtgactg gaggccttgg gaccaaaggc aagggaagaa ggaatgagaa 420

```

```

gtacgatatg gtgacagact gtggctacac aatctctcaa gtgagatcaa tgaagattct 480
gaagcgattt ggtggcccag ctgggtctatg gaccaaggat ccactggggc aaacagagaa 540
gatctacgtg ttagatggga cacagaatga cacagccttt gtcttcccaa ggctgcgtga 600
cttcaccctt gccatggctg cccggaaaagc ttcccgagtc cgggtgccct tccctgggt 660
aggcacaggg cagctgggtat atgggtggctt tctttatttt gctcggaggc ctcttggaag 720
acctggtgga ggtggtgaga tggagaacac tttgcagcta atcaaattcc acctggcaaa 780
ccgaacagtg gtggacagct cagtattccc agcagagggg ctgatcccc cctacggctt 840
gacagcagac acctacatcg acctggcagc tgatgaggaa ggtctttggg ctgtctatgc 900
caccggggag gatgacaggc acttgtgtct ggccaagtta gatccacaga cactggacac 960
agagcagcag tgggacacac catgtcccag agagaatgct gaggtgcct ttgtcatctg 1020
tgggaccctc tatgtcgtct ataacaccg tcctgccagt cgggcccgca tccagtgtct 1080
ctttgatgcc agcgccacc tgacctga acgggcagca ctcccttatt tcccccag 1140
atatggtgcc catgccagcc tccgtataa ccccgagaa cgcagctct atgcctggga 1200
tgatggctac cagattgtct ataagctgga gatgaggaag aaagaggagg aggtttgagg 1260
agctagcctt gtttttgc tctttctcac tccatacat ttatattata tccccactaa 1320
atttctgtt cctcattctt caaatgtggg ccagttgtg ctcaaactct ctatattttt 1380
agccaatggc aatcaaattc tttcagctcc tttgtttcat acggaactcc agatcctgag 1440
taatcctttt agagcccgaa gagtcaaaac cctcaatgtt cctcctgct ctcctgccc 1500
atgtcaacaa atttcaggct aaggatgcc cagaccag gctctaacct tgtatgcggg 1560
caggccagg gagcaggcag cagtgttctt cccctcagag tgacttggg agggagaaat 1620
aggaggagac gtccagctct gtccctctct cctcactcct ccttcagtgc tctgaggaa 1680
caggactttc tccacattgt tttgtattgc aacattttgc attaaaagga aaatccactg 1740
ctaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1800
aaaaa 1805

```

&lt;210&gt; 110

&lt;211&gt; 406

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 110

```

Met Gly Pro Ser Thr Pro Leu Leu Ile Leu Phe Leu Leu Ser Trp Ser
  1             5             10            15

Gly Pro Leu Gln Gly Gln Gln His His Leu Val Glu Tyr Met Glu Arg
      20             25            30

Arg Leu Ala Ala Leu Glu Glu Arg Leu Ala Gln Cys Gln Asp Gln Ser
      35             40            45

Ser Arg His Ala Ala Glu Leu Arg Asp Phe Lys Asn Lys Met Leu Pro
      50             55            60

Leu Leu Glu Val Ala Glu Lys Glu Arg Glu Ala Leu Arg Thr Glu Ala
      65             70            75            80

Asp Thr Ile Ser Gly Arg Val Asp Arg Leu Glu Arg Glu Val Asp Tyr
      85             90            95

Leu Glu Thr Gln Asn Pro Ala Leu Pro Cys Val Glu Phe Asp Glu Lys
      100            105            110

Val Thr Gly Gly Pro Gly Thr Lys Gly Lys Gly Arg Arg Asn Glu Lys
      115            120            125

Tyr Asp Met Val Thr Asp Cys Gly Tyr Thr Ile Ser Gln Val Arg Ser
      130            135            140

Met Lys Ile Leu Lys Arg Phe Gly Gly Pro Ala Gly Leu Trp Thr Lys
      145            150            155            160

```

Asp Pro Leu Gly Gln Thr Glu Lys Ile Tyr Val Leu Asp Gly Thr Gln  
 165 170 175  
 Asn Asp Thr Ala Phe Val Phe Pro Arg Leu Arg Asp Phe Thr Leu Ala  
 180 185 190  
 Met Ala Ala Arg Lys Ala Ser Arg Val Arg Val Pro Phe Pro Trp Val  
 195 200 205  
 Gly Thr Gly Gln Leu Val Tyr Gly Gly Phe Leu Tyr Phe Ala Arg Arg  
 210 215 220  
 Pro Pro Gly Arg Pro Gly Gly Gly Glu Met Glu Asn Thr Leu Gln  
 225 230 235 240  
 Leu Ile Lys Phe His Leu Ala Asn Arg Thr Val Val Asp Ser Ser Val  
 245 250 255  
 Phe Pro Ala Glu Gly Leu Ile Pro Pro Tyr Gly Leu Thr Ala Asp Thr  
 260 265 270  
 Tyr Ile Asp Leu Ala Ala Asp Glu Glu Gly Leu Trp Ala Val Tyr Ala  
 275 280 285  
 Thr Arg Glu Asp Asp Arg His Leu Cys Leu Ala Lys Leu Asp Pro Gln  
 290 295 300  
 Thr Leu Asp Thr Glu Gln Gln Trp Asp Thr Pro Cys Pro Arg Glu Asn  
 305 310 315 320  
 Ala Glu Ala Ala Phe Val Ile Cys Gly Thr Leu Tyr Val Val Tyr Asn  
 325 330 335  
 Thr Arg Pro Ala Ser Arg Ala Arg Ile Gln Cys Ser Phe Asp Ala Ser  
 340 345 350  
 Gly Thr Leu Thr Pro Glu Arg Ala Ala Leu Pro Tyr Phe Pro Arg Arg  
 355 360 365  
 Tyr Gly Ala His Ala Ser Leu Arg Tyr Asn Pro Arg Glu Arg Gln Leu  
 370 375 380  
 Tyr Ala Trp Asp Asp Gly Tyr Gln Ile Val Tyr Lys Leu Glu Met Arg  
 385 390 395 400  
 Lys Lys Glu Glu Glu Val  
 405

&lt;210&gt; 111

&lt;211&gt; 2824

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 111

gtcccgagcagg ggctggagcc ctgggcgctg caaagcgtgt cccgccgggt ccccgagcgt 60  
 cccgcgcctt cgcgcgcctt tgctcctgct gctggggctg tgccctggggc tgcctcctgtg 120  
 tgtggggctg caggaagagg cgcagagctg gggccactct tcggagcagg atggactcag 180  
 ggtcccaggg caagtcagac tgttcagag gctgaaaacc aaacctttga tgacagaatt 240



```

ctcagtgaag tctaccatca tttcccggtta tgccttcact acggttttccct gcagaatgct 300
gaacagagct tctgaagacc aggacattga gttccagatg cagattccag ctgcagcttt 360
catcaccaac ttcactatgc ttattggaga caaggtgtat cagggcgaaa ttacagagag 420
agaaaagaag agtgggtgata gggtaaaaga gaaaagggaat aaaaccacag aagaaaatgg 480
agagaaggggg actgaaatat tcagagcttct tgcaagtatt cccagcaaaag acaaagccgc 540
ctttttctctg agttatgagg agcttctgca gaggcgcttg ggcaagtacg agcacagcat 600
cagcgtgcgg ccccagcagc tgtccgggag gctgagcgtg gacgtgaata tcctggagag 660
cgcgggcatc gcatccctgg aggtgctgcc gcttcacaac agcaggcaga ggggcagtgg 720
gcgcgggggaa gatgattctg ggcctcccc atctactgtc attaaccaaa atgaaacatt 780
tgccaacata atttttaaac ctactgtagt acaacaagcc aggattgccc agaattggaat 840
tttgggagac tttatcatta gatatgacgt caatagagaa cagagcattg gggacatcca 900
ggttctaaaat ggcatttttg tgcactactt tgctcctaaa gaccttcctc ctttacccaa 960
gaatgtggta ttcgtgcttg acagcagtgc ttctatgggtg ggaaccaaac tccggcagac 1020
caaggatgcc ctcttcacaa ttctccatga cctccgaccc caggaccgtt tcagtatcat 1080
tggtttttcc aaccggatca aagtatggaa ggaccacttg atatcagtca ctccagacag 1140
catcagggat gggaaagtgt acattcacca tatgtcaccc actggaggta aagatgcac 1200
cttcttttca cattggcttg gctttgaaat aatgttttct ttttttgtt tttttttttg 1260
tttttttgcc aagagataag gatttgaaag ctgggggtga tggcaggaag ggagactgca 1320
ggcgagagct ttctcagcag cttaggttgc tcggaatacc aaatccgtac tctccttttc 1380
cagtctacaa gtgaagctgg tgaagtcttg ggggaaaata acctgatagg tgtttggagg 1440
cttgccctggt tcatctcatg gatcctgtta gaaaaagtct tgctgataca aaaacaatac 1500
tgtttttgta aatggttttg ggggtggagaa ttacagccac gtggaccagc cagtgtttcg 1560
tgcgcacctt gggcaggggc aggtggaacg ctttgcaaaa caaggagccc tcttggtaga 1620
gctgcagcag agtgaaacag gtatcctggt ggcccgggag ggttcgagg gtaaaatgtg 1680
atggggagga tgtgggctct cagccactgc acgcttgtct tgggtgttgc ccaggtaagg 1740
gtgcagaatg ttaagagagg agagaaaaga tcataaaaat tctgagaaga gtggagaatc 1800
agagcagtgt gttaccagct gtcctagatg ttaagtcttc ccaactcagg acctgcgcct 1860
tactcttgcc attccctgta tcaacagtag agtggacaca gaagaggtgc tcagtattgg 1920
agcatttagc agaacttgac ctactccctc ccccttcctt tcaggctctg tacagagaca 1980
ccacctctg gttaattcca cttcagttat ctggaaacat acaaaacact ctgttcagtc 2040
ttctcccatc tgggattttc catctttctt atattcatgg tttccagtc caagacaaag 2100
ccctcatttc ccaaatgaac ttaatgtgga acttgatctt gctgggccc ggtactttcc 2160
atttacctg ggatttccta ctagcctccc aaacagtata atctaggaac tgttttgggtg 2220
tttgatggga gaaatagaag attagattta atatgatagt gatgtctac ggcagggaag 2280
ccacaacacc ataaaattct tgaaggaaca gatttctct catgtttaca tagctcctac 2340
ctccatgcac agttcctaac acagaggaca tgctcaatgg atattttag attggattta 2400
ttttaatggc cctgcttact agcagaatga tgtgtgtgtg ctggtggagt gggggccatg 2460
gagtaataaa cagcttttaga aaatggctgg gccgggtgtg gtggctcaca cttgtaatcc 2520
cagcactttg ggaggccgag gcaggtggat cacctgaggt caggagttca agaccagcct 2580
ggccaacatg gcgaaacccc atctctacta aaaatacaaa aagttagccg ggtgtgggtg 2640
cgggtgcctg taatcccagc tacttgggag gctgaggcag gagaattgct tgaacctggg 2700
aggcagaggt tgcagtgagc caagatcatg ccagtgcact ccagcctggg tgacagcgag 2760
actctgtctc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2820
aaaa
2824

```

&lt;210&gt; 112

&lt;211&gt; 399

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 112

Met Leu Leu Leu Leu Gly Leu Cys Leu Gly Leu Ser Leu Cys Val Gly  
1 5 10 15

Ser Gln Glu Glu Ala Gln Ser Trp Gly His Ser Ser Glu Gln Asp Gly  
20 25 30

Leu Arg Val Pro Arg Gln Val Arg Leu Leu Gln Arg Leu Lys Thr Lys  
35 40 45

Pro Leu Met Thr Glu Phe Ser Val Lys Ser Thr Ile Ile Ser Arg Tyr  
 50 55 60  
 Ala Phe Thr Thr Val Ser Cys Arg Met Leu Asn Arg Ala Ser Glu Asp  
 65 70 75 80  
 Gln Asp Ile Glu Phe Gln Met Gln Ile Pro Ala Ala Ala Phe Ile Thr  
 85 90 95  
 Asn Phe Thr Met Leu Ile Gly Asp Lys Val Tyr Gln Gly Glu Ile Thr  
 100 105 110  
 Glu Arg Glu Lys Lys Ser Gly Asp Arg Val Lys Glu Lys Arg Asn Lys  
 115 120 125  
 Thr Thr Glu Glu Asn Gly Glu Lys Gly Thr Glu Ile Phe Arg Ala Ser  
 130 135 140  
 Ala Val Ile Pro Ser Lys Asp Lys Ala Ala Phe Phe Leu Ser Tyr Glu  
 145 150 155 160  
 Glu Leu Leu Gln Arg Arg Leu Gly Lys Tyr Glu His Ser Ile Ser Val  
 165 170 175  
 Arg Pro Gln Gln Leu Ser Gly Arg Leu Ser Val Asp Val Asn Ile Leu  
 180 185 190  
 Glu Ser Ala Gly Ile Ala Ser Leu Glu Val Leu Pro Leu His Asn Ser  
 195 200 205  
 Arg Gln Arg Gly Ser Gly Arg Gly Glu Asp Asp Ser Gly Pro Pro Pro  
 210 215 220  
 Ser Thr Val Ile Asn Gln Asn Glu Thr Phe Ala Asn Ile Ile Phe Lys  
 225 230 235 240  
 Pro Thr Val Val Gln Gln Ala Arg Ile Ala Gln Asn Gly Ile Leu Gly  
 245 250 255  
~~Asp Phe Ile Ile Arg Tyr Asp Val Asn Arg Glu Gln Ser Ile Gly Asp~~  
~~260 265 270~~  
 Ile Gln Val Leu Asn Gly Tyr Phe Val His Tyr Phe Ala Pro Lys Asp  
 275 280 285  
 Leu Pro Pro Leu Pro Lys Asn Val Val Phe Val Leu Asp Ser Ser Ala  
 290 295 300  
 Ser Met Val Gly Thr Lys Leu Arg Gln Thr Lys Asp Ala Leu Phe Thr  
 305 310 315 320  
 Ile Leu His Asp Leu Arg Pro Gln Asp Arg Phe Ser Ile Ile Gly Phe  
 325 330 335  
 Ser Asn Arg Ile Lys Val Trp Lys Asp His Leu Ile Ser Val Thr Pro  
 340 345 350  
 Asp Ser Ile Arg Asp Gly Lys Val Tyr Ile His His Met Ser Pro Thr  
 355 360 365

Gly Gly Lys Asp Asp Thr Phe Phe Ser His Trp Leu Gly Phe Glu Ile  
370 375 380

Met Phe Ser Phe Phe Val Phe Phe Phe Cys Phe Phe Ala Lys Arg  
385 390 395

<210> 113  
<211> 1711  
<212> DNA  
<213> Homo sapiens

<400> 113  
aggaaatgtc agaggaacca gaactcagaa ctcttctctc tggaccttcc ttcccttccc 60  
ttggagggtat ccctttgaat caggcctctc tcttctcatc agtctgtagc ttcccccttt 120  
gtataacctg ctttcttttt tacattttatt aaaagtggat tttgtaaaag catttcattg 180  
acacgcgacc tatcacagac aatggaattc gtcagtgggtg gtaagactga aatcctgatg 240  
cttttcacac ttcttgcttc ttgctatgta ttctgcctc tagccttgcc atgttttgcc 300  
tttttttttc tttttggcca attccttttt atatgtgccc acaacagagg tggggagaca 360  
cggagcacc cgggtccttc ccagcgctgc tgggcaggcc ccgtctccag gccccagctg 420  
ttgaaacttt gaagggaac aaacaacat ccacactgcc ggaccctagg ctgttcaggg 480  
aggcagctca tttccacccc ggccccagga cccccagcct gtccccaca aggatctctc 540  
taaatgggag ggattgaggc tactttttctg ccaagcccta ttaagtagta atgtggggaa 600  
acccactgtg tcagtgcagg aagccctaga caaatgtttt caaataaatt tcaactgccc 660  
gcctgcacag atttccattt gaagtacttc ccattccacc tgacacccaa aggggttttt 720  
ttgttttgtt ttgtttttga gacaggtctt tgctttgttg ccaggtgg agtgacagtga 780  
cgtggtcata gctcactgca gcctcaacct cctgggtcga agtgacctc ctgcctcagc 840  
ctcccaaagt tctgagatga taggcattgag ccattgtgcc tagcctattt tgattttttt 900  
cttagagtca aggtcttgct ctgttgccca ggctgatctt ggacttgcca gccaccatgc 960  
ctggctgggt tttttaaaaa tagaatctca ctgatagcct gcaagaaaca gatgcagtgc 1020  
ctgcttcctg atcagtcaca ggagccctcg tgtttgccac ctttaccttt gaacctcccc 1080  
ctgcctccct gcctgtgtcc gcttttgtag ctcaatgcag ccatgacaag gaaagaaaag 1140  
acaaaggaag gccagagagc cgcgcatgtc tctgcaggtg cagatgcagg cagtggaggt 1200  
ggcctgagca ggcagaagga caccaagcgc cctatgttgc ttgtcattca tgacgtgggtc 1260  
ttggagcttc tgactagtcc agactgccac gccaacccca gaaaataccc cacatgccag 1320  
aaaagtgaag tcctaggtgt ttccatctat gtttcaatct gtccatctac caggcctcgc 1380  
gataaaaaca aaacaaaaaa acgctgccag gttttagaag cagttctggt ctcaaaaaca 1440  
tcaggatcct gccaccaggg ttcttttgaa atagtaccac atgtaaaagg gaatttggtc 1500  
ttcacttcat ctaataactg aattgtcagg ctttgattga taattgtaga aataagtagc 1560  
cttctgttgt gggaataagt tataatcagt attcatctct ttgttttttg tcaactcttt 1620  
ctctctaatt gtgtcatttg tactgtttga aaaatatctt ttctataaaa ttaactaac 1680  
ctgccttaaa aaaaaaaaaa aaaaaaaaaa a 1711

<210> 114  
<211> 76  
<212> PRT  
<213> Homo sapiens

<400> 114  
Met Glu Phe Val Ser Gly Gly Lys Thr Glu Ile Leu Met Leu Phe Thr  
1 5 10 15  
Leu Leu Val Ser Cys Tyr Val Phe Leu Pro Leu Ala Leu Pro Cys Phe  
20 25 30  
Ala Phe Phe Phe Leu Phe Gly Gln Phe Leu Phe Ile Cys Ala His Asn  
35 40 45  
Arg Gly Gly Glu Thr Arg Ser Thr Leu Gly Pro Ser Gln Arg Cys Trp  
50 55 60

Ala Gly Pro Val Ser Arg Pro Gln Leu Leu Lys Leu  
65 70 75

<210> 115  
<211> 2116  
<212> DNA  
<213> Homo sapiens

<400> 115  
agtgttgggg ttgcaggaga cctaaacaca gtcaccatga agctgggctg tgtcctcatg 60  
gcctgggccc tctaccttcc ccttgggtgtg ctctgggtgg cccagatgct actggctgcc 120  
agttttgaga cgctgcagtg tgaggagacct gtctgcactg aggagagcag ctgccacacg 180  
gaggatgact tgactgatgc aagggaagct ggcttccagg tcaaggccta cactttcagt 240  
gaaccttccc acctgattgt gtcctatgac tggctgatcc tccaaggctc agccaagcca 300  
gtttttgaag gggacctgct ggttctgcgc tgccaggcct ggcaagactg gccactgact 360  
caggtgacct tctaccgaga tggctcagct ctgggtcccc ccgggcctaa cagggaattc 420  
tccatcaccg tggtaaaaaa ggcagacagc gggcactacc actgcagtgg catcttccag 480  
agccctggtc ctgggatccc agaaacagca tctgttgtgg ctatcacagt ccaagaactg 540  
tttccagcgc caattctcag agctgtacct tcagctgaac cccaagcagg agggcccatg 600  
acctgtgatt gtcagacaaa gttgccctcg cagaggtcag ctgcccgcct cctcttctcc 660  
ttctacaagg atggaaggat agtgcaaaag agggggctct cctcagaatt ccagatcccc 720  
acagcttcag aagatcactc cgggtcatac tgggtgtgagg cagccactga ggacaaccaa 780  
gtttggaaac agagccccca gctagagatc agagtgcagg gtgcttccag ctctgctgca 840  
cctcccacat tgaatccagc tcctcagaaa tcagctgctc caggaactgc tcctgaggag 900  
gcccctgggc ctctgcctcc gccgccaaac ccattctctg aggatccagg cttttcttct 960  
cctctgggga tgcagatcc tcattctgtat caccagatgg gccttcttct caaacacatg 1020  
caggatgtga gagtccctct cggtcacctg ctcatggagt tgagggaatt atctggccac 1080  
cggaagcctg ggaccacaaa ggctactgct gaatagaagt aaacagttca tccatgatct 1140  
cacttaacca cccaataaaa tctgattctt tattttctct tcctgtcctg cacatatgca 1200  
taagtacttt tacaagttgt ccagtggtt tgttagaata atgtagttag gtgagtgtaa 1260  
ataaatttat ataaagttag aattagagtt tagctataat tgtgtattct ctcttaacac 1320  
aacagaattc tgctgtctag atcaggaatt tctatctgtt atatcgacca gaatgttggt 1380  
atttaaagag aactaatgga agtggattga atacagcagt ctcaactggg ggcaattttg 1440  
ccccccagag gacattgggc aatgtttgga gacattttgg tcattatact tgggggggtg 1500  
ggggatggtg ggtgtgtgt gctactggca tccagtaaat agaagccagg ggtgccgcta 1560  
aacatcctat aatgcacagg gcagtacccc acaacgaaaa ataactctggc ccaaaatgtc 1620  
agttgtactg agtttgagaa accccagcct aatgaaaccc taggtgttgg gctctggaat 1680  
gggactttgt ccttctctaat tattatctct tccagcctc attcagctat tcttactgac 1740  
ataccagtct ttagtgtgtg ctatggtctg ttcttttagtt ctagtgttga tcccctcaaa 1800  
agccattatg ttgaaatcct aatccccaag gtgatggcat taagaagtgg gcctttggga 1860  
agtgattaga tcaggagtgc agagccctca tgattaggat tagtgccctt atttaaaaag 1920  
gccccagaga gctaactcac ccttccacca tatgaggacg tggcaagaag atgacatgta 1980  
tgagaaccaa aaaacagctg tcgccaacaa ccgactctgt cgttgccttg atcttgaact 2040  
tccagcctcc agaactatga gaaataaaat tctgttgttt gtaagctaaa aaaaaaaaaa 2100  
aaaaaaaaaa aaaaaa 2116

<210> 116  
<211> 359  
<212> PRT  
<213> Homo sapiens

<400> 116  
Met Lys Leu Gly Cys Val Leu Met Ala Trp Ala Leu Tyr Leu Ser Leu  
1 5 10 15  
Gly Val Leu Trp Val Ala Gln Met Leu Leu Ala Ala Ser Phe Glu Thr  
20 25 30

Leu Gln Cys Glu Gly Pro Val Cys Thr Glu Glu Ser Ser Cys His Thr  
 35 40 45  
 Glu Asp Asp Leu Thr Asp Ala Arg Glu Ala Gly Phe Gln Val Lys Ala  
 50 55 60  
 Tyr Thr Phe Ser Glu Pro Phe His Leu Ile Val Ser Tyr Asp Trp Leu  
 65 70 75 80  
 Ile Leu Gln Gly Pro Ala Lys Pro Val Phe Glu Gly Asp Leu Leu Val  
 85 90 95  
 Leu Arg Cys Gln Ala Trp Gln Asp Trp Pro Leu Thr Gln Val Thr Phe  
 100 105 110  
 Tyr Arg Asp Gly Ser Ala Leu Gly Pro Pro Gly Pro Asn Arg Glu Phe  
 115 120 125  
 Ser Ile Thr Val Val Gln Lys Ala Asp Ser Gly His Tyr His Cys Ser  
 130 135 140  
 Gly Ile Phe Gln Ser Pro Gly Pro Gly Ile Pro Glu Thr Ala Ser Val  
 145 150 155 160  
 Val Ala Ile Thr Val Gln Glu Leu Phe Pro Ala Pro Ile Leu Arg Ala  
 165 170 175  
 Val Pro Ser Ala Glu Pro Gln Ala Gly Gly Pro Met Thr Leu Ser Cys  
 180 185 190  
 Gln Thr Lys Leu Pro Leu Gln Arg Ser Ala Ala Arg Leu Leu Phe Ser  
 195 200 205  
 Phe Tyr Lys Asp Gly Arg Ile Val Gln Ser Arg Gly Leu Ser Ser Glu  
 210 215 220  
 Phe Gln Ile Pro Thr Ala Ser Glu Asp His Ser Gly Ser Tyr Trp Cys  
 225 230 235 240  
~~Glu Ala Ala Thr Glu Asp Asn Gln Val Trp Lys Gln Ser Pro Gln Leu~~  
~~245 250 255~~  
 Glu Ile Arg Val Gln Gly Ala Ser Ser Ser Ala Ala Pro Pro Thr Leu  
 260 265 270  
 Asn Pro Ala Pro Gln Lys Ser Ala Ala Pro Gly Thr Ala Pro Glu Glu  
 275 280 285  
 Ala Pro Gly Pro Leu Pro Pro Pro Pro Thr Pro Ser Ser Glu Asp Pro  
 290 295 300  
 Gly Phe Ser Ser Pro Leu Gly Met Pro Asp Pro His Leu Tyr His Gln  
 305 310 315 320  
 Met Gly Leu Leu Leu Lys His Met Gln Asp Val Arg Val Leu Leu Gly  
 325 330 335  
 His Leu Leu Met Glu Leu Arg Glu Leu Ser Gly His Arg Lys Pro Gly  
 340 345 350

Thr Thr Lys Ala Thr Ala Glu  
355

<210> 117  
<211> 1391  
<212> DNA  
<213> Homo sapiens

<400> 117  
atccttggaa aagaaaatgt ttatgttgca gggatttgca tggtcacgag tgagggcagg 60  
cccttgggga cacatctgcc cacagctgca caggccaggg cgcaggcaca tctgttgggt 120  
ctcaggcctc agataaaacc atctccgcat catatggcca gtgaccgctt tctcccttca 180  
agaaaattct gtggctgtgc agtactttga agttttaatt attaacctgc tttaattaaa 240  
gcagtttctt ttcttataaa gtggaatcac caaatcttat cacacagagc acagtccctgt 300  
agttaccag cccgctccag cagtgcggga gattgtaagg aagcgggtggc ggctggtgaa 360  
gcaagtctca catgtcggcg ttcttggcca atggatacaa agataaagaa aatgttgctt 420  
tttcttagga actgtcagaa atcctcatgc ctttcaagac ttctgtgaat gacttgaatt 480  
ttttattccc tgcctagggt ctgtgaacga ggctgtctc ttccctgggg tttctttcca 540  
tggccttctt ttctctctt ccagtgggag ttttgcaggc tcttctctgt ggaaacttca 600  
cgagcgttgg ctgggcctcg gcttcgctgg agtgactcc aggggtgaagg cagagtggga 660  
tttgagacc aggttaggca cgaccaggc tgagaaggga cgtttccatc attcacagt 720  
ccctccccac agcactacct caccgccacc cccaccctca ctctacccc accccgcgat 780  
cgtcagggtg gccacgggtg gccggagggt gccggctctg gctgtccctg tgccgggtccc 840  
tcacaaacct ctcccccttt gaaactcaag cacagctgcg aggagggcag cgaggaggga 900  
cccctctctc atggttgtct ctttccccg ctatgtcata ggtagtggag gaagcgaagg 960  
aagtgaacgc tgaatgtgac gcatttctga agagctcagc tgtcaccggg catagcctgg 1020  
aagccccaag tctgttctga ctttgcctgg ctgtctcctt gaccgcctc ctatgcatt 1080  
gtccttgatg tccaggctgg gtcattttaa atagagatgc aatcaggaag gttgggggac 1140  
ttgggactgt ggctgaattg agacctgtct gatgtattca tgtcagcacc tgagtcacag 1200  
cccagggtgc cggaagcagc ctcttcgcat aggcagtgat ttgcgattac tttaaagctc 1260  
accttttttc tteccctctc tgttcgctgc tgtcagcata atgattgtgt tccttcccta 1320  
tgggatccat ctgttttga aacaataaag cgtctgaggg agtgtaaaaa aaaaaaaaaa 1380  
aaaaaaaaa a 1391

<210> 118  
<211> 56  
<212> PRT  
<213> Homo sapiens

<400> 118  
Met Val Thr Ser Glu Gly Arg Pro Leu Gly Thr His Leu Pro Thr Ala  
1 5 10 15  
Ala Gln Ala Arg Ala Gln Ala His Leu Leu Val Leu Arg Pro Gln Ile  
20 25 30  
Lys Pro Ser Pro His His Met Ala Ser Asp Arg Phe Leu Pro Ser Arg  
35 40 45  
Lys Phe Cys Gly Cys Ala Val Leu  
50 55

<210> 119  
<211> 21  
<212> DNA  
<213> Artificial Sequence  
  
<220>

<223> oligonucleotide

<400> 119

cttccacaga acacaagcca c

21

<210> 120

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 120

acgctcaact ccacctcc

18

<210> 121

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 121

cttggaacat agcaccactc c

21

<210> 122

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 122

ccattccaga cttccctgtc

20

<210> 123

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 123

gatgcagggt gtctcctgg

19

<210> 124

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 124

ctgtggacta cggaagggtg

20

<210> 125  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 125  
gaacagatgg actctccccc 20

<210> 126  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 126  
tggaggcatt gctatgtgg 19

<210> 127  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 127  
tcagggagaa tgagcacatc t 21

<210> 128  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

---

<400> 128  
ccgatcaatt ttacacaaca a 21

<210> 129  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 129  
ggacacaaga agaggagagc a 21

<210> 130  
<211> 20  
<212> DNA  
<213> Artificial Sequence



<220>  
<223> oligonucleotide  
  
<400> 130  
tcacctcaga tgagtgtggc  
  
<210> 131  
<211> 21  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> oligonucleotide  
  
<400> 131  
acagatggat gatctgtgaa c  
  
<210> 132  
<211> 21  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> oligonucleotide  
  
<400> 132  
ggagactcac tatgaatccc t  
  
<210> 133  
<211> 21  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> oligonucleotide  
  
<400> 133  
tttaaacaca ttccctgact c  
  
<210> 134  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> oligonucleotide  
  
<400> 134  
ccttgacacag cacttgacat  
  
<210> 135  
<211> 21  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> oligonucleotide  
  
<400> 135  
tgggtctcag ttaccatttg g

20

21

21

21

20

21

<210> 136  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 136  
gtgaattagt gaagagccag c

21

<210> 137  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 137  
ttctctgaaa ctgagtcgcc t

21

<210> 138  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 138  
tgggagtcgc ttagcctatc

20

<210> 139  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 139  
agtcatgaat ggcacctggt

20

<210> 140  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 140  
cataaaacag ctttccccca

20

<210> 141  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 141  
aggagtttcc agggcagttt 20

<210> 142  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 142  
ggctcagata tagttcaggc a 21

<210> 143  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 143  
aggcttatac tacggcgggt 20

<210> 144  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 144  
gggagggaga gtttgcctc 20

<210> 145  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 145  
gcctcaccct ttggttatga 20

<210> 146  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 146

aacaggcact ttgaagtcag c

21

&lt;210&gt; 147

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;400&gt; 147

tggttgaga tgaacatccc

20

&lt;210&gt; 148

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;400&gt; 148

cctgaagatc cagcatgact t

21

&lt;210&gt; 149

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;400&gt; 149

ggcaaactgt ctaaaaagtg a

21

&lt;210&gt; 150

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;400&gt; 150

atattgcaaa tgctgcacca

20

&lt;210&gt; 151

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;400&gt; 151

cagctgcctc ctttaacagc

20

&lt;210&gt; 152

&lt;211&gt; 20

&lt;212&gt; DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 152

tcatcacacc atccatcctg

20

<210> 153

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 153

ggatccctag gctctgttcc

20

<210> 154

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 154

tggaaaaccg ttatagaccc a

21

<210> 155

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 155

ggctcaggta aacaaagatt g

21

<210> 156

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 156

aagagatcaa cgtcgggatg

20

<210> 157

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 157  
ggggatttca gtttcagcaa 20

<210> 158  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 158  
tcattcaaca accagaacgt g 21

<210> 159  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 159  
gggctatcac tgtggctatg a 21

<210> 160  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 160  
tttaattgga agagtgggcg 20

<210> 161  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 161  
tacctcacgc ctgtaatccc 20

<210> 162  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 162  
gaggagctat ggacgtctgc 20

<210> 163  
<211> 21

<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 163  
agttcattca gccttataca a

21

<210> 164  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 164  
ctaggttctg aagaggggcc

20

<210> 165  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 165  
ctgaggccag ttgtttccat

20

<210> 166  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 166  
ggatcagcag gattacttgc a

21

<210> 167  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 167  
ttcacgcatt cttcaagcag

20

<210> 168  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

&lt;400&gt; 168

cctgaaatct ttggccttga

20

&lt;210&gt; 169

&lt;211&gt; 113

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 169

Met Val Leu Thr Leu Trp Cys Asn Leu Cys Ser Arg Ala Ser Ser Trp  
 1 5 10 15

Val Arg Gln Lys His Val Ser Cys Cys Val His Asn Tyr Thr Gln Pro  
 20 25 30

Phe Leu Leu Ile Gln Ser Ser Phe Trp Ala Met Ser Ser Glu Thr Lys  
 35 40 45

Pro Lys Ala Leu Ser Lys Asp Tyr Leu Cys Ile Ser Tyr Arg Ser Pro  
 50 55 60

His Ser Thr Pro Thr His Arg His Ser Ser Asn Ser Ser Tyr Asp Leu  
 65 70 75 80

Pro Val Glu Ala Gln Ala Ser Tyr Leu Asp Ile Lys Ser Leu His Gly  
 85 90 95

Gln Ser Gly Leu Cys Leu Ser Arg Phe Ile Phe His Tyr His Thr Pro  
 100 105 110

Tyr

&lt;210&gt; 170

&lt;211&gt; 321

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 170

Met Ala Val Ser Glu Arg Arg Gly Leu Gly Arg Gly Ser Pro Ala Glu  
 1 5 10 15

Trp Gly Gln Arg Leu Leu Leu Val Leu Leu Gly Gly Cys Ser Gly  
 20 25 30

Arg Ile His Arg Leu Ala Leu Thr Gly Glu Lys Arg Ala Asp Ile Gln  
 35 40 45

Leu Asn Ser Phe Gly Phe Tyr Thr Asn Gly Ser Leu Glu Val Glu Leu  
 50 55 60

Ser Val Leu Arg Leu Gly Leu Arg Glu Ala Glu Glu Lys Ser Leu Leu  
 65 70 75 80

Val Gly Phe Ser Leu Ser Arg Val Arg Ser Gly Arg Val Arg Ser Tyr  
 85 90 95

Ser Thr Arg Asp Phe Gln Asp Cys Pro Leu Gln Lys Asn Ser Ser Ser



100	105	110
Phe Leu Val Leu Phe Leu Ile Asn Thr Lys Asp Leu Gln Val Gln Val		
115	120	125
Arg Lys Tyr Gly Glu Gln Lys Thr Leu Phe Ile Phe Pro Gly Leu Leu		
130	135	140
Pro Glu Ala Pro Ser Lys Pro Gly Leu Pro Lys Pro Gln Ala Thr Val		
145	150	155
Pro Arg Lys Val Asp Gly Gly Gly Thr Ser Ala Ala Ser Lys Pro Lys		
165	170	175
Ser Thr Pro Ala Val Ile Gln Gly Pro Ser Gly Lys Asp Lys Asp Leu		
180	185	190
Val Leu Gly Leu Ser His Leu Asn Asn Ser Tyr Asn Phe Ser Phe His		
195	200	205
Val Val Ile Gly Ser Gln Ala Glu Glu Gly Gln Tyr Ser Leu Asn Phe		
210	215	220
His Asn Cys Asn Asn Ser Val Pro Gly Lys Glu His Pro Phe Asp Ile		
225	230	235
Thr Val Met Ile Arg Glu Lys Asn Pro Asp Gly Phe Leu Ser Ala Ala		
245	250	255
Glu Met Pro Leu Phe Lys Leu Tyr Met Val Met Ser Ala Cys Phe Leu		
260	265	270
Ala Ala Gly Ser Gly Cys Thr Ser Ser Trp Trp Arg Ala Pro Pro Trp		
275	280	285
Pro Ser Ser Cys Ser Arg Ala Thr Ser Ser Ser Pro Gln Glu Thr Thr		
290	295	300
Arg Thr Cys Ser Cys Pro Arg Arg Thr Arg Arg Met Phe Arg Trp Ser		
305	310	315
		320

Lys

&lt;210&gt; 171

&lt;211&gt; 39

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 171

Met Gln Arg Val Glu Val Phe Ser Thr Gln Glu Leu Ala Asp Val Asn
1 5 10 15

Glu Val Leu Arg Met Gly Pro Ser Pro Ile Ser Val Ala Ser Thr Glu
20 25 30

Phe Cys Tyr Pro Ser Phe Arg
35

&lt;210&gt; 172

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 172

Gly Trp Gly His Leu Leu Phe Leu Trp Pro Val Leu Ser Phe Val Ile  
 1 5 10 15

Leu Pro Leu Gly Lys Glu Cys Gln Trp Thr Asp Ala Cys Leu Ser His  
 20 25 30

Pro Cys Ala Asn Gly Ser Thr Cys Thr Thr Val Ala Asn Gln Phe Ser  
 35 40 45

Cys Lys Cys Leu Thr Gly Phe Thr Gly Gln Lys Cys Glu Thr Asp Val  
 50 55 60

Asn Glu Cys Asp Ile Pro Gly His Cys Gln His Gly Gly Thr Cys Leu  
 65 70 75 80

Asn Leu Pro Gly Ser Tyr Gln Cys Gln Cys Leu Gln Gly Phe Thr Gly  
 85 90 95

Gln Tyr Cys Asp Ser Leu Tyr Val Pro Cys Ala Pro Ser Pro Cys Val  
 100 105 110

Asn Gly Gly Thr Cys Arg Gln Thr Gly Asp Phe Thr Phe Glu Cys Asn  
 115 120 125

Cys Leu Pro Gly Lys Glu Leu Pro Ser Val Pro Gly Leu Gly Asp Lys  
 130 135 140

Pro Leu Ala Gln Glu Val Val Gly Val Ala Gln Leu Phe Phe Leu Gly  
 145 150 155 160

Ser Ala Arg Lys Lys Gly Ser Glu Asn Phe Val Gly Gly Gly Leu Leu  
 165 170 175

Val Arg Glu Glu Phe Tyr Gly Pro Thr Val Val His Lys Leu Ser Arg  
 180 185 190

Gly

&lt;210&gt; 173

&lt;211&gt; 72

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 173

Met Pro Ala Cys Leu Ile Pro Val Gln Met Glu Val Pro Val Pro Leu  
 1 5 10 15

Trp Pro Thr Ser Ser Pro Ala Asn Ala Ser Gln Ala Ser Gln Gly Arg  
 20 25 30

Ser Val Arg Leu Met Ser Met Ser Val Thr Phe Gln Asp Leu Pro Ala

35 40 45  
 Trp Trp His Leu Pro Gln Pro Ala Trp Phe Leu Pro Val Pro Val Pro  
 50 55 60  
 Ser Gly Leu His Arg Pro Val Leu  
 65 70

<210> 174  
 <211> 73  
 <212> PRT  
 <213> Homo sapiens

<400> 174  
 Met Leu Arg Ala Gly Ala Ala Gln Thr Cys Ser Ala Gly Leu Gln Val  
 1 5 10 15  
 Leu Lys Pro Tyr Trp Gly Trp Val Gly Ser Gly Ala Ala Ala Phe Ala  
 20 25 30  
 Thr Leu Arg Ile Gly Ala Lys Ala Thr Asp Val Tyr Leu Thr Val Thr  
 35 40 45  
 Leu His Trp Val Leu Lys Glu Ile Ile Ser Arg Cys Asn Tyr Asn Tyr  
 50 55 60  
 Cys Leu Leu Arg Lys Ile Trp Glu Phe  
 65 70

<210> 175  
 <211> 78  
 <212> PRT  
 <213> Homo sapiens

<400> 175  
 Met Val Leu Val Ser Ser Phe Phe Val Phe Tyr Ser Val His Ser Phe  
 1 5 10 15  
 Leu Thr Ile Trp Thr Thr Val Val Ala Asn Pro Gly Gln Trp Ile Val  
 20 25 30  
 Thr Asn Ser Val Leu Val Ala Ser Cys Phe Pro Ala Arg Ser Pro Phe  
 35 40 45  
 Val Leu Ile Met Ser Asp Thr His Ile Ser Gln Phe Cys Phe Ala Cys  
 50 55 60  
 Arg Thr Arg Lys Thr Leu Phe Pro Asn Leu Val Val Met Pro  
 65 70 75

<210> 176  
 <211> 249  
 <212> PRT  
 <213> Homo sapiens

<400> 176  
 Met Trp Arg Lys Asn Gln Tyr Val Ser Asn Gly Leu Arg Asp Phe Ala

```
<210> 177
<211> 191
<212> PRT
<213> Homo sapiens
```

```

<400> 177
Met Ile Thr Val Asp Ile Ile Pro Ser Gly Trp Asn Ser Ala Asp Gly
 1             5             10             15

Lys Ser Asp Lys Thr Lys Ser Ala Pro Ser Arg Asp Pro Glu Arg Leu
          20             25             30

```

Gln Lys Ile Lys Glu Ser Leu Leu Leu Glu Asp Ser Glu Glu Glu Glu  
 35 40 45  
 Gly Asp Leu Cys Arg Ile Cys Gln Met Ala Ala Ala Ser Ser Ser Asn  
 50 55 60  
 Leu Leu Ile Glu Pro Cys Lys Cys Thr Gly Ser Leu Gln Tyr Val His  
 65 70 75 80  
 Gln Asp Cys Met Lys Lys Trp Leu Gln Ala Lys Ile Asn Ser Gly Ser  
 85 90 95  
 Ser Leu Glu Ala Val Thr Thr Cys Glu Leu Cys Lys Glu Lys Leu Glu  
 100 105 110  
 Leu Asn Leu Glu Asp Phe Asp Ile His Glu Leu His Arg Ala His Ala  
 115 120 125  
 Asn Glu Gln Ala Glu Tyr Glu Phe Ile Ser Ser Gly Leu Tyr Leu Val  
 130 135 140  
 Val Leu Leu His Leu Cys Glu Gln Ser Phe Ser Asp Met Met Gly Asn  
 145 150 155 160  
 Thr Asn Glu Pro Ser Thr Arg Val Arg Leu Gln Arg Met Ile Pro Lys  
 165 170 175  
 Lys Thr Glu Thr Ile Thr Gly His Leu Ile Leu Pro Asn Phe Ile  
 180 185 190

&lt;210&gt; 178

&lt;211&gt; 80

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 178

Met Phe Leu Ala Cys Leu Cys Leu Glu Asn Trp Ser Ser Gln Ala Pro  
 1 5 10 15  
 Leu Ala Ala Thr Ser Pro Cys Trp Ala Ser Glu Thr Ser Leu Cys Leu  
 20 25 30  
 Val Ser Tyr Tyr Ala Leu Ser Phe Ala Met Thr Thr Thr Lys Ser Lys  
 35 40 45  
 Pro Val Gly Thr Pro Val Gly Pro Leu Asp Leu Pro Thr Ser Pro Gly  
 50 55 60  
 Ala Cys Arg Arg Ser Pro Thr Phe Thr Ala Pro Ser Ser Asp Thr Leu  
 65 70 75 80

&lt;210&gt; 179

&lt;211&gt; 62

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 179

Met Pro Gly Phe Ala Gly Phe Ile Cys Leu Ile Leu Phe Cys Val Phe

1                      5                      10                      15  
 Ser Trp Leu Phe Gly Ser Phe Pro Gly Thr Leu Asp Gly Ser Ile Pro  
                     20                      25                      30  
 Arg His Leu Val Ile Lys Gln Leu Ser Pro Thr Pro Tyr His Gly Lys  
                     35                      40                      45  
 Arg Gly Arg Asn Ile Ala Pro Ser Leu Ile Thr Tyr His Leu  
                     50                      55                      60

<210> 180  
 <211> 61  
 <212> PRT  
 <213> Homo sapiens

<400> 180  
 Met Leu Gly Ser Leu Gly Asp Ala Arg Phe Cys Gly Phe Tyr Leu Phe  
   1                      5                      10                      15  
 Asn Phe Ile Leu Cys Phe Leu Leu Ala Leu Trp Val Phe Pro Gly Tyr  
                     20                      25                      30  
 Thr Arg Trp Leu His Pro Lys Ala Ser Cys His Lys Thr Ala Phe Pro  
                     35                      40                      45  
 His Pro Ile Ser Trp Glu Lys Gly Glu Lys Tyr Ser Pro  
                     50                      55                      60

<210> 181  
 <211> 60  
 <212> PRT  
 <213> Homo sapiens

<400> 181  
 Met Met Ile Ser Leu His Thr Val Gln Ser His Asn Leu Lys Ile Lys  
   1                      5                      10                      15  
 Leu Ser Trp Leu Cys Phe Leu Cys Ser Cys Gln Asn Ile Gly Thr Ile  
                     20                      25                      30  
 Gly Arg Ser Lys Thr Phe Ile Leu Leu Leu Gln Val Tyr Leu Gly Thr  
                     35                      40                      45  
 Phe Thr Cys Val Phe Lys Gly Ile Ser Phe Gln Gln  
                     50                      55                      60

<210> 182  
 <211> 227  
 <212> PRT  
 <213> Homo sapiens

<400> 182  
 Met Met Gly Ser Glu Ala Ala Gly Arg Gly Ser Gln Glu Leu Leu Val  
   1                      5                      10                      15  
 Val Gln Pro Val Leu Pro Ser Glu Ala Leu Leu Phe Pro Gly Leu Pro

```
<210> 183
<211> 97
<212> PRT
<213> Homo sapiens
```

Met Leu Ser Cys Thr Leu Gly Leu Thr Val Cys Pro Leu Ser Pro Ala  
1 5 10 15

Pro Ser Val Thr Leu Ala Val Ala Leu Asn Gly Gln Leu Arg Arg Pro  
20 25 30

Leu Cys Cys Ser Ser Ala Phe Pro Glu Val Gly Glu Pro Ala Trp Pro  
35 40 45

Arg Pro Leu Ser Ser Asp Gln Ala Leu Ser Pro Arg Ser Tyr Gly Arg  
50 55 60

Pro Gly Ser Gly Val Gly Thr His Gly Pro Gly Trp Gly Gly Ala Gln  
65 70 75 80

Ser Asp Val Asn Phe Phe Pro Cys Val Asp Met Tyr Ser Gln Arg Val  
85 90 95

Val

<210> 184

<211> 68

<212> PRT

<213> Homo sapiens

<400> 184

Met Cys Phe Leu Leu Phe Gly Ser Leu Cys Ile Tyr Tyr Phe Ser Leu  
1 5 10 15

Phe Leu Val Phe Phe Phe Ser Cys Phe Cys Phe Val Trp Cys Phe Val  
20 25 30

Pro Val Phe Ile Val Ser Gly Ile Ser Leu Pro Leu Trp Ile Pro His  
35 40 45

Gly Leu Asp Arg Asp Gly Pro Val Met Pro Ser Ser Phe Leu Leu Leu  
50 55 60

Leu Leu Leu Trp  
65

<210> 185

<211> 142

<212> PRT

<213> Homo sapiens

<400> 185

Met Phe Ser Cys Asn Glu Asn Ser Ile Phe Phe Arg Ile Gly Phe Val  
1 5 10 15

Phe Ile Leu Leu Ser Phe Ile Ser Ser Cys Gln Thr Leu Asn Gly Tyr  
20 25 30

Val Cys Ile Leu Ile Thr Leu Phe Ser Leu Leu Trp Lys Arg Arg Thr  
35 40 45

Arg Glu Gln Met Leu Leu Arg Ala Gly Val Ser Glu Lys Asn Leu Ser  
50 55 60

Met Leu Phe Asn Val Phe Leu Pro Leu Pro His Ser Val Cys Val Thr  
65 70 75 80

Phe Tyr Asn Ile Lys Lys Tyr Tyr Asn Ile Ser Arg Ile Trp Asn Cys  
85 90 95

His His Asp Glu Trp Pro Phe Gln Cys Ile Val Thr Glu Ile Pro Glu  
100 105 110

Asp Ser Pro Gly Leu Gln Phe His Trp Phe Leu Leu Gln Phe Leu Val



115 120 125

Ala Val Ile Val Ala Val Ser Ser Leu Lys Asp Leu Leu Trp  
 130 135 140

<210> 186  
 <211> 111  
 <212> PRT  
 <213> Homo sapiens

<400> 186

Met Ser Cys Pro Leu Pro Leu Leu Ile Ser Ala Ile Ala Val Gly  
 1 5 10 15

Ser Ser Met Gln Thr His Ala Arg Ala Ser Phe Ala Ala Gly Pro Ser  
 20 25 30

Gln Glu Asp Phe Ser Ala His Leu Ala Gln Asp Gln His Ser Pro Glu  
 35 40 45

Val Gln Gly His Tyr His Ala Arg Gly Asn Pro Pro Ala Val Gly Asp  
 50 55 60

Thr Ser Leu Trp Met Lys Val Pro Thr Ser His His Ser Asp Glu Lys  
 65 70 75 80

His Gln Glu Ala Ser Cys Thr Phe Leu Lys Arg Pro Gln Gln Asp Gln  
 85 90 95

Ser Pro Ile Ala His Ser Ser His Leu Asn Asn Ala Pro Phe Tyr  
 100 105 110

<210> 187  
 <211> 72  
 <212> PRT  
 <213> Homo sapiens

<400> 187

Met Phe Gly Met Pro His Thr Met Ser Cys Pro Leu Pro Leu Leu Ile  
 1 5 10 15

Ser Ala Ile Ala Ala Val Gly Ser Ser Met Gln Thr His Ala Arg Ala  
 20 25 30

Ser Phe Ala Ala Gly Pro Ser Gln Lys Thr Ser Gln Pro Ile Trp Ser  
 35 40 45

Arg Ile Phe Leu Pro Leu Lys Val Thr Ala Pro Lys Ser Cys Pro Met  
 50 55 60

Phe Tyr Phe Gln Glu Phe Pro Asn  
 65 70

<210> 188  
 <211> 109  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 188

Met Asp Ala Arg Trp Trp Ala Val Val Val Leu Ala Ala Phe Pro Ser  
 1 5 10 15

Leu Gly Ala Gly Gly Glu Thr Pro Glu Ala Pro Pro Glu Ser Trp Thr  
 20 25 30

Gln Leu Trp Phe Phe Arg Phe Val Val Asn Ala Ala Gly Tyr Ala Ser  
 35 40 45

Phe Met Val Pro Gly Tyr Leu Leu Val Gln Tyr Phe Arg Arg Lys Asn  
 50 55 60

Tyr Leu Glu Thr Gly Met Gly Leu Cys Phe Pro Leu Val Lys Ala Cys  
 65 70 75 80

Val Phe Gly Asn Glu Pro Lys Ala Ser Asp Glu Val Pro Leu Arg Pro  
 85 90 95

Gln Gln Arg Arg Gln Arg Pro Pro Arg Cys Gly Arg Pro  
 100 105

&lt;210&gt; 189

&lt;211&gt; 76

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 189

Met Trp Pro Ala Leu His Leu Leu His His Trp Ala Val Trp Gly Cys  
 1 5 10 15

Arg Leu His His His His Asp Pro Pro Pro Gly Leu Cys His Pro Ser  
 20 25 30

Phe Leu Pro Ser Leu Trp Pro His Cys His Cys Gly Gly Arg Ala Gly  
 35 40 45

Gly Gly Cys Gly Leu Cys Cys Pro Pro Ala Gln Ser Leu Arg Ala Gly  
 50 55 60

Pro Ser Lys Ala Thr Gly Lys Glu Gly Cys Ala Cys  
 65 70 75

&lt;210&gt; 190

&lt;211&gt; 168

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 190

Leu Cys Arg Ala Leu Ile Lys Arg Ile Gln Ala Met Ile Pro Lys Gly  
 1 5 10 15

Ala Leu Ala Val Ala Val Ala Gln Val Cys Arg Val Val Pro Leu Val  
 20 25 30

Ala Gly Gly Ile Cys Gln Cys Leu Ala Glu Arg Tyr Ser Val Ile Leu  
 35 40 45

Leu Asp Thr..Leu Leu Gly Arg Met Leu Pro Gln Leu Val Cys Arg Leu  
 50 55 60  
 Val Leu Arg Cys Ser Met Asp Asp Ser Ala Gly Pro Arg Glu Trp Leu  
 65 70 75 80  
 Pro Arg Asp Ser Glu Cys His Leu Cys Met Ser Val Thr Thr Gln Ala  
 85 90 95  
 Gly Asn Ser Ser Glu Gln Ala Ile Pro Gln Ala Met Leu Gln Ala Cys  
 100 105 110  
 Val Gly Ser Trp Leu Asp Arg Glu Lys Cys Lys Gln Phe Val Glu Gln  
 115 120 125  
 His Thr Pro Gln Leu Leu Thr Leu Val Pro Arg Gly Trp Asp Ala His  
 130 135 140  
 Thr Thr Cys Gln Ala Leu Gly Val Cys Gly Thr Met Ser Ser Pro Leu  
 145 150 155 160  
 Gln Cys Ile His Ser Pro Asp Leu  
 165

<210> 191  
 <211> 272  
 <212> PRT  
 <213> Homo sapiens

<400> 191  
 Met Ala Glu Ser His Leu Leu Gln Trp Leu Leu Leu Leu Leu Pro Thr  
 1 5 10 15  
 Leu Cys Gly Pro Gly Thr Ala Ala Trp Thr Thr Ser Ser Leu Ala Cys  
 20 25 30  
 Ala Gln Gly Pro Glu Phe Trp Cys Gln Ser Leu Glu Gln Ala Leu Gln  
 35 40 45  
 Cys Arg Ala Leu Gly His Cys Leu Gln Glu Val Trp Gly His Val Gly  
 50 55 60  
 Ala Asp Asp Leu Cys Gln Glu Cys Glu Asp Ile Val His Ile Leu Asn  
 65 70 75 80  
 Lys Met Ala Lys Glu Ala Ile Phe Gln Asp Leu Ser Glu Gln Gln Phe  
 85 90 95  
 Pro Ile Pro Leu Pro Tyr Cys Trp Leu Cys Arg Ala Leu Ile Lys Arg  
 100 105 110  
 Ile Gln Ala Met Ile Pro Lys Gly Ala Leu Ala Val Ala Val Ala Gln  
 115 120 125  
 Val Cys Arg Val Val Pro Leu Val Ala Gly Gly Ile Cys Gln Cys Leu  
 130 135 140  
 Ala Glu Arg Tyr Ser Val Ile Leu Leu Asp Thr Leu Leu Gly Arg Met

145                      150                      155                      160  
 Leu Pro Gln Leu Val Cys Arg Leu Val Leu Arg Cys Ser Met Asp Asp  
                                  165                      170                      175  
 Ser Ala Gly Pro Arg Glu Trp Leu Pro Arg Asp Ser Glu Cys His Leu  
                                  180                      185                      190  
 Cys Met Ser Val Thr Thr Gln Ala Gly Asn Ser Ser Glu Gln Ala Ile  
                                  195                      200                      205  
 Pro Gln Ala Met Leu Gln Ala Cys Val Gly Ser Trp Leu Asp Arg Glu  
                                  210                      215                      220  
 Lys Cys Lys Gln Phe Val Glu Gln His Thr Pro Gln Leu Leu Thr Leu  
 225                                   230                                   235                                   240  
 Val Pro Arg Gly Trp Asp Ala His Thr Thr Cys Gln Ala Leu Gly Val  
                                  245                                   250                                   255  
 Cys Gly Thr Met Ser Ser Pro Leu Gln Cys Ile His Ser Pro Asp Leu  
                                  260                                   265                                   270

<210> 192  
 <211> 60  
 <212> PRT  
 <213> Homo sapiens

<400> 192  
 Met Pro Pro Ser Ala Phe Leu Phe Phe Phe Trp Arg Gln Ser Leu Ala  
   1                                  5                                  10                                  15  
 Leu Leu Pro Arg Leu Glu Cys Ser Ser Thr Ile Ser Ala Leu Thr Ala  
                                   20                                  25                                  30  
 Thr Ser Val Ser Trp Val Gln Ala Ile Leu Leu Pro Gln Pro Pro Lys  
                                   35                                  40                                  45  
 Tyr Leu Gly Leu Gln Ala Cys Ala Thr Thr Pro Gly  
                                   50                                  55                                  60

<210> 193  
 <211> 357  
 <212> PRT  
 <213> Homo sapiens

<400> 193  
 Met Pro Ile Leu Thr Gly Asp Phe Leu Leu Pro Thr Pro Gln Phe Tyr  
   1                                  5                                  10                                  15  
 Ala Glu Asn Ile Asn Thr Thr Ser Leu Thr Cys Ser Ser Asp Arg Met  
                                   20                                  25                                  30  
 Arg Val Ile Ile Ser Lys Ser Tyr Leu Glu Ala Phe Asn Ser Asn Gly  
                                   35                                  40                                  45  
 Asn Asn Leu Gln Leu Lys Asp Pro Thr Cys Arg Pro Lys Leu Ser Asn  
                                   50                                  55                                  60

Val Val Glu Phe Ser Val Pro Leu Asn Gly Cys Gly Thr Ile Arg Lys  
 65 70 75 80  
 Val Glu Asp Gln Ser Ile Thr Tyr Thr Asn Ile Ile Thr Phe Ser Ala  
 85 90 95  
 Ser Ser Thr Ser Glu Val Ile Thr Arg Gln Lys Gln Leu Gln Ile Ile  
 100 105 110  
 Val Lys Cys Glu Met Gly His Asn Ser Thr Val Glu Ile Ile Tyr Ile  
 115 120 125  
 Thr Glu Asp Asp Val Ile Gln Ser Gln Asn Ala Leu Gly Lys Tyr Asn  
 130 135 140  
 Thr Ser Met Ala Leu Phe Glu Ser Asn Ser Phe Glu Lys Thr Ile Leu  
 145 150 155 160  
 Glu Ser Pro Tyr Tyr Val Asp Leu Asn Gln Thr Leu Phe Val Gln Val  
 165 170 175  
 Ser Leu His Thr Ser Asp Pro Asn Leu Val Val Phe Leu Asp Thr Cys  
 180 185 190  
 Arg Ala Ser Pro Thr Ser Asp Phe Ala Ser Pro Thr Tyr Asp Leu Ile  
 195 200 205  
 Lys Ser Gly Cys Ser Arg Asp Glu Thr Cys Lys Val Tyr Pro Leu Phe  
 210 215 220  
 Gly His Tyr Gly Arg Phe Gln Phe Asn Ala Phe Lys Phe Leu Arg Ser  
 225 230 235 240  
 Met Ser Ser Val Tyr Leu Gln Cys Lys Val Leu Ile Cys Asp Ser Ser  
 245 250 255  
 Asp His Gln Ser Arg Cys Asn Gln Gly Cys Val Ser Arg Ser Lys Arg  
 260 265 270  
 Asp Ile Ser Ser Tyr Lys Trp Lys Thr Asp Ser Ile Ile Gly Pro Ile  
 275 280 285  
 Arg Leu Lys Arg Asp Arg Ser Ala Ser Gly Asn Ser Gly Phe Gln His  
 290 295 300  
 Glu Thr His Ala Glu Glu Thr Pro Asn Gln Pro Phe Asn Ser Val His  
 305 310 315 320  
 Leu Phe Ser Phe Met Val Leu Ala Leu Asn Val Val Thr Val Ala Thr  
 325 330 335  
 Ile Thr Val Arg His Phe Val Asn Gln Arg Ala Asp Tyr Lys Tyr Gln  
 340 345 350  
 Lys Leu Gln Asn Tyr  
 355

&lt;210&gt; 194

&lt;211&gt; 169

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 194

Met Gln Cys Leu Leu Pro Tyr Gln Ser Lys Glu Pro Ser Cys Leu Pro  
 1 5 10 15

Pro Leu Pro Leu Asn Leu Pro Leu Pro Pro Cys Leu Cys Pro Leu Leu  
 20 25 30

Gln Leu Asn Ala Ala Met Thr Arg Lys Glu Lys Thr Lys Glu Gly Gln  
 35 40 45

Arg Ala Ala Gln Phe Ser Ala Gly Ala Asp Ala Gly Ser Gly Gly Gly  
 50 55 60

Leu Ser Arg Gln Lys Asp Thr Lys Arg Pro Met Leu Leu Val Ile His  
 65 70 75 80

Asp Val Val Leu Glu Leu Leu Thr Ser Ser Asp Cys His Ala Asn Pro  
 85 90 95

Arg Lys Tyr Pro Thr Cys Gln Lys Ser Glu Val Leu Gly Val Ser Ile  
 100 105 110

Tyr Val Ser Ile Cys Pro Ser Thr Arg Pro Arg Asp Lys Asn Lys Thr  
 115 120 125

Lys Lys Arg Cys Gln Val Leu Glu Ala Val Leu Val Ser Lys Pro Ser  
 130 135 140

Gly Ser Cys His Gln Gly Ser Phe Glu Ile Val Pro His Val Lys Gly  
 145 150 155 160

Asn Leu Ala Phe Thr Ser Ser Asn Asn  
 165

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/07285

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : C12Q 1/68; C12N 15/00, 15/09, 15/63, 15/86 US CL : 435/6, 69.1, 69.3, 320.1, 455, 471; 530/300, 350; 424/189.1 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6, 69.1, 69.3, 320.1, 455, 471; 530/300, 350; 424/189.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EAST, USPATFULL		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	Database EST on STN. Hudson et al. AN X11582. 'New isolated nucleic acid segments from the human genome - used for determining polymorphic forms for use in e.g. forensics, paternity testing or phenotypic typing for disease' WO 98/20165 see, bases 1-67, which would hybridize to SEQ ID NO: 1.	1 -- 2-11
X -- Y	Database EST on STN. Hillier et al. AN W51776. 'The WashU-Merck EST Project' 11 October 1996. See Sequence Alignment (attached) disclosing 85% similarity to SEQ ID NO: 1, and would hybridize to SEQ ID NO: 1	1 -- 2-11
X -- Y	Database EST on STN. 'NCI-CGAP' AN AA568724. '09 September 1997. See Sequence Alignment (attached) which discloses a polynucleotide with 88% similarity to SEQ ID NO: 1 and would hybridize to SEQ ID NO: 1.	1 -- 2-11
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* *A* *E* *L* *O* *P*	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* *X* *Y* *A* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
Date of the actual completion of the international search 27 JUNE 2000		Date of mailing of the international search report 19 JUL 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer MARY K ZEMAN Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/07285

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P --- Y, P	Database Gen EMBL. AN AC009651. Birren et al. 'Homo sapiens chromosome 11, clone' 29 September 1999. See Sequence Alignment (attached) which discloses a polynucleotide having up to 98% identity to SEQ ID NO: 1, and could encode SEQ ID NO: 2.	1 -- 2-11



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/07285

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-11, SEQ ID NO: 1 and 2

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/07285

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-11, drawn to polynucleotides (SEQ ID NO: 1) and encoded polypeptides (SEQ ID NO: 2), and methods of making the recombinant polypeptides.

Group II, claim(s) 12-13, drawn to polynucleotides (SEQ ID NO: 3) and encoded polypeptides (SEQ ID NO: 4).

Group III, claim(s) 14-15, drawn to polynucleotides (SEQ ID NO: 5) and encoded polypeptides (SEQ ID NO: 6).

Group IV, claim(s) 16-17, drawn to polynucleotides (SEQ ID NO: 7) and encoded polypeptides (SEQ ID NO: 8).

Group V, claim(s) 18-19, drawn to polynucleotides (SEQ ID NO: 9) and encoded polypeptides (SEQ ID NO: 10).

Group VI, claim(s) 20-21, drawn to polynucleotides (SEQ ID NO: 11) and encoded polypeptides (SEQ ID NO: 12).

Group VII, claim(s) 22-23, drawn to polynucleotides (SEQ ID NO: 13) and encoded polypeptides (SEQ ID NO: 14).

Group VIII, claim(s) 24-25, drawn to polynucleotides (SEQ ID NO: 15) and encoded polypeptides (SEQ ID NO: 16).

Group IX, claim(s) 26-27, drawn to polynucleotides (SEQ ID NO: 17) and encoded polypeptides (SEQ ID NO: 18).

Group X, claim(s) 28-29, drawn to polynucleotides (SEQ ID NO: 19) and encoded polypeptides (SEQ ID NO: 20).

Group XI, claim(s) 30-31, drawn to polynucleotides (SEQ ID NO: 21) and encoded polypeptides (SEQ ID NO: 22).

Group XII, claim(s) 32-33, drawn to polynucleotides (SEQ ID NO: 23) and encoded polypeptides (SEQ ID NO: 24).

Group XIII, claim(s) 34-35, drawn to polynucleotides (SEQ ID NO: 25) and encoded polypeptides (SEQ ID NO: 26).

Group XIV, claim(s) 36-37, drawn to polynucleotides (SEQ ID NO: 27) and encoded polypeptides (SEQ ID NO: 28).

Group XV, claim(s) 38-39, drawn to polynucleotides (SEQ ID NO: 29) and encoded polypeptides (SEQ ID NO: 30).

Group XVI, claim(s) 40-41, drawn to polynucleotides (SEQ ID NO: 31) and encoded polypeptides (SEQ ID NO: 32).

Group XVII, claim(s) 42-43, drawn to polynucleotides (SEQ ID NO: 33) and encoded polypeptides (SEQ ID NO: 34).

Group XVIII, claim(s) 44-45, drawn to polynucleotides (SEQ ID NO: 35) and encoded polypeptides (SEQ ID NO: 36).

Group XIX, claim(s) 46-47, drawn to polynucleotides (SEQ ID NO: 37) and encoded polypeptides (SEQ ID NO: 38).

Group XX, claim(s) 48-49, drawn to polynucleotides (SEQ ID NO: 39) and encoded polypeptides (SEQ ID NO: 40).

Group XXI, claim(s) 50-51, drawn to polynucleotides (SEQ ID NO: 41) and encoded polypeptides (SEQ ID NO: 42).

Group XXII, claim(s) 52-53, drawn to polynucleotides (SEQ ID NO: 43) and encoded polypeptides (SEQ ID NO: 44).

Group XXIII, claim(s) 54-55, drawn to polynucleotides (SEQ ID NO: 45) and encoded polypeptides (SEQ ID NO: 46).

Group XXIV, claim(s) 56-57, drawn to polynucleotides (SEQ ID NO: 47) and encoded polypeptides (SEQ ID NO: 48).

Group XXV, claim(s) 58-59, drawn to polynucleotides (SEQ ID NO: 49) and encoded polypeptides (SEQ ID NO: 50).

Group XXVI, claim(s) 60-61, drawn to polynucleotides (SEQ ID NO: 51) and encoded polypeptides (SEQ ID NO: 52).

Group XXVII, claim(s) 62-63, drawn to polynucleotides (SEQ ID NO: 53) and encoded polypeptides (SEQ ID NO: 54).

Group XXVIII, claim(s) 64-65, drawn to polynucleotides (SEQ ID NO: 55) and encoded polypeptides (SEQ ID NO: 56).

Group XXIX, claim(s) 66-67, drawn to polynucleotides (SEQ ID NO: 57) and encoded polypeptides (SEQ ID NO: 58).

Group XXX, claim(s) 68-69, drawn to polynucleotides (SEQ ID NO: 59) and encoded polypeptides (SEQ ID NO: 60).

Group XXXI, claim(s) 70-71, drawn to polynucleotides (SEQ ID NO: 61) and encoded polypeptides (SEQ ID NO: 62).

Group XXXII, claim(s) 72-73, drawn to polynucleotides (SEQ ID NO: 63) and encoded polypeptides (SEQ ID NO: 64).

Group XXXIII, claim(s) 74-75, drawn to polynucleotides (SEQ ID NO: 65) and encoded polypeptides (SEQ ID NO: 66).

Group XXXIV, claim(s) 76-77, drawn to polynucleotides (SEQ ID NO: 67) and encoded polypeptides (SEQ ID NO: 68).

Group XXXV, claim(s) 78-79, drawn to polynucleotides (SEQ ID NO: 69) and encoded polypeptides (SEQ ID NO: 70).

Group XXXVI, claim(s) 80-81, drawn to polynucleotides (SEQ ID NO: 71) and encoded polypeptides (SEQ ID NO: 72).

Group XXXVII, claim(s) 82-83, drawn to polynucleotides (SEQ ID NO: 73) and encoded polypeptides (SEQ ID NO: 74).

Group XXXVIII, claim(s) 84-85, drawn to polynucleotides (SEQ ID NO: 75) and encoded polypeptides (SEQ ID NO: 76).

Group XXXIX, claim(s) 86-87, drawn to polynucleotides (SEQ ID NO: 77) and encoded polypeptides (SEQ ID NO: 78).

Group XXXX, claim(s) 88-89, drawn to polynucleotides (SEQ ID NO: 79) and encoded polypeptides (SEQ ID NO: 80).

Group XXXXI, claim(s) 90-91, drawn to polynucleotides (SEQ ID NO: 81) and encoded polypeptides (SEQ ID NO: 82).

Group XXXXII, claim(s) 92-93, drawn to polynucleotides (SEQ ID NO: 83) and encoded polypeptides (SEQ ID NO: 84).

Group XXXXIII, claim(s) 94-95, drawn to polynucleotides (SEQ ID NO: 85) and encoded polypeptides (SEQ ID NO: 86).

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/07285

86).  
Group XXXXIV, claim(s) 96-97, drawn to polynucleotides (SEQ ID NO: 87) and encoded polypeptides (SEQ ID NO: 88).  
Group XXXXV, claim(s) 98-99, drawn to polynucleotides (SEQ ID NO: 89) and encoded polypeptides (SEQ ID NO: 90).  
Group XXXXVI, claim(s) 100-101, drawn to polynucleotides (SEQ ID NO: 91) and encoded polypeptides (SEQ ID NO: 92).  
Group XXXXVII, claim(s) 102-103, drawn to polynucleotides (SEQ ID NO: 93) and encoded polypeptides (SEQ ID NO: 94).  
Group XXXXVIII, claim(s) 104-105, drawn to polynucleotides (SEQ ID NO: 95) and encoded polypeptides (SEQ ID NO: 96).  
Group XXXXIX, claim(s) 106-107, drawn to polynucleotides (SEQ ID NO: 97) and encoded polypeptides (SEQ ID NO: 98).  
Group L, claim(s) 108-109, drawn to polynucleotides (SEQ ID NO: 99) and encoded polypeptides (SEQ ID NO: 100).  
Group LI, claim(s) 101-111, drawn to polynucleotides (SEQ ID NO: 101) and encoded polypeptides (SEQ ID NO: 102).  
Group LII, claim(s) 112-113, drawn to polynucleotides (SEQ ID NO: 103) and encoded polypeptides (SEQ ID NO: 104).  
Group LIII, claim(s) 114-115, drawn to polynucleotides (SEQ ID NO: 105) and encoded polypeptides (SEQ ID NO: 106).  
Group LIV, claim(s) 116-117, drawn to polynucleotides (SEQ ID NO: 107) and encoded polypeptides (SEQ ID NO: 108).  
Group LV, claim(s) 118-119, drawn to polynucleotides (SEQ ID NO: 109) and encoded polypeptides (SEQ ID NO: 110).  
Group LVI, claim(s) 120-121, drawn to polynucleotides (SEQ ID NO: 111) and encoded polypeptides (SEQ ID NO: 112).  
Group LVII, claim(s) 122-123, drawn to polynucleotides (SEQ ID NO: 113) and encoded polypeptides (SEQ ID NO: 114).  
Group LVIII, claim(s) 124-125, drawn to polynucleotides (SEQ ID NO: 115) and encoded polypeptides (SEQ ID NO: 116).  
Group LIX, claim(s) 126-127, drawn to polynucleotides (SEQ ID NO: 117) and encoded polypeptides (SEQ ID NO: 118).

The inventions listed as Groups ONE (I) to FIFTY NINE (LIX) do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Each polynucleotide and corresponding polypeptide do not share any sequence homology, similar structure or other feature which could be considered a special technical feature. Each polynucleotide sequence and corresponding polypeptide sequence is a separate and distinct invention, having no obvious shared features, and thus, lack unity under PCT Rule 13.2.